



INSTITUT
POLYTECHNIQUE
DE PARIS

*J*ournal de l'École polytechnique *Mathématiques*

Stéphane FISCHLER & Tanguy RIVOAL

Values of E -functions are not Liouville numbers

Tome II (2024), p. 1-18.

<https://doi.org/10.5802/jep.249>

© Les auteurs, 2024.



Cet article est mis à disposition selon les termes de la licence
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.
<https://creativecommons.org/licenses/by/4.0/>

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org
e-ISSN : 2270-518X

VALUES OF E -FUNCTIONS ARE NOT LIOUVILLE NUMBERS

BY STÉPHANE FISCHLER & TANGUY RIVOAL

ABSTRACT. — Shidlovskii has given a linear independence measure of values of E -functions with rational Taylor coefficients at a rational point, not a singularity of the underlying differential system satisfied by these E -functions. Recently, Beukers has proved a qualitative linear independence theorem for the values at an algebraic point of E -functions with arbitrary algebraic Taylor coefficients. In this paper, we obtain an analogue of Shidlovskii's measure for values of arbitrary E -functions at algebraic points. This enables us to solve a long standing problem by proving that the value of an E -function at an algebraic point is never a Liouville number. We also prove that values at rational points of E -functions with rational Taylor coefficients are linearly independent over $\overline{\mathbb{Q}}$ if and only if they are linearly independent over \mathbb{Q} . Our methods rest upon improvements of results obtained by André and Beukers in the theory of E -operators.

RÉSUMÉ (Les valeurs des E -fonctions ne sont pas des nombres de Liouville)

Shidlovskii a donné une mesure d'indépendance linéaire de valeurs de E -fonctions à coefficients de Taylor rationnels en un point rationnel qui n'est pas une singularité du système différentiel sous-jacent vérifié par ces E -fonctions. Récemment, Beukers a prouvé un théorème d'indépendance linéaire qualitatif pour les valeurs en un point algébrique de E -fonctions à coefficients de Taylor algébriques arbitraires. Dans cet article, nous obtenons un analogue de la mesure de Shidlovskii pour des valeurs de E -fonctions arbitraires en des points algébriques. Cela nous permet de résoudre un problème longtemps ouvert : la valeur d'une E -fonction en un point algébrique n'est jamais un nombre de Liouville. Nous prouvons également que des valeurs aux points rationnels de E -fonctions à coefficients de Taylor rationnels sont linéairement indépendantes sur $\overline{\mathbb{Q}}$ si et seulement si elles sont linéairement indépendantes sur \mathbb{Q} . Nos méthodes reposent sur des améliorations de résultats obtenus par André et Beukers concernant la théorie des E -opérateurs.

CONTENTS

1. Introduction.....	2
2. Main tools.....	7
3. Proof of Theorem 1.....	9
4. Decomposition of E -functions over a number field.....	15
5. Structure of $\mathbf{E}_{\mathbb{K}}$	16
6. A Galois action on values of E -functions.....	16
References.....	17

MATHEMATICAL SUBJECT CLASSIFICATION (2020). — 11J82, 11J91.

KEYWORDS. — E -functions, André-Beukers theorems, linear independence measures, irrationality measures, transcendence measures, Liouville numbers, Shidlovskii's theorem.

1. INTRODUCTION

Siegel [25] defined in 1929 the class of E -functions in order to generalize the Diophantine properties of the exponential function (namely the Lindemann-Weierstrass Theorem) to other special functions such as Bessel's function $J_0(z) := \sum_{n=0}^{\infty} (-1)^n (z/2)^{2n} / n!^2$ or hypergeometric series ${}_pF_p$ with rational parameters. A power series $\sum_{n=0}^{\infty} a_n z^n / n! \in \mathbb{Q}[[z]]$ is said to be an E -function when it is solution of a linear differential equation over $\overline{\mathbb{Q}}(z)$ (i.e., holonomic), and $|\sigma(a_n)|$ (for any $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$) and the least common denominator of a_0, a_1, \dots, a_n all grow at most exponentially in n . Note that Siegel's original definition of E -functions is more general: see the end of this introduction. Throughout this paper we fix an embedding of $\overline{\mathbb{Q}}$ in \mathbb{C} .

A lot of important qualitative results are known on the arithmetic nature of the values taken by E -functions at algebraic points, amongst which we cite the celebrated Siegel-Shidlovskii Theorem (see [24] for instance). This result is not always strong enough, in particular to deduce linear independence of values of E -functions. It has been improved in [22] and [10] (see Theorem B below for the linear case), but some assumptions still have to be checked (for instance the functions should not be evaluated at a singular point).

Quantitative versions of the Siegel-Shidlovskii Theorem have been proved, such as the measure of algebraic independence of Lang-Galochkin (see [14, p. 238, Th. 5.29 & remarks] and [24, Ch. 11]), but they hold only under certain assumptions on algebraic independence or rationality of the coefficients of the E -functions. In the linear setting, the main result is the following one, due to Shidlovskii [24, p. 358, Th. 1, Eq. (32)].

THEOREM A (Shidlovskii). — *Let $\underline{f} = {}^t(f_1, \dots, f_N) \in \mathbb{Q}[[z]]^N$ be a vector of E -functions which is solution of a differential system $\underline{f}' = A\underline{f}$ for some $A \in M_N(\mathbb{Q}(z))$. Assume that f_1, \dots, f_N are linearly independent over $\mathbb{Q}(z)$ and that $z_0 \in \mathbb{Q}^*$ is not a pole of an entry of A . Then for any $\varepsilon > 0$, there exists $c = c(\varepsilon, z_0, f_1, \dots, f_N) > 0$ such that for all $\lambda_1, \dots, \lambda_N \in \mathbb{Z}$ not all zero, we have*

$$\left| \sum_{j=1}^N \lambda_j f_j(z_0) \right| > c H^{-N+1-\varepsilon} \quad \text{where } H := \max_{1 \leq j \leq N} |\lambda_j|.$$

This theorem holds *verbatim* with \mathbb{Q} replaced by an imaginary quadratic number field and \mathbb{Z} replaced by its ring of integers. However, no such result is known for other number fields \mathbb{K} . The point is that all known quantitative results are based on the Siegel-Shidlovskii method only, which provides linear independence of the full set of the values of E -functions in Theorem A only when \mathbb{K} is either \mathbb{Q} or imaginary quadratic. Even the qualitative part of Theorem A (namely, $\sum_{j=1}^N \lambda_j f_j(z_0) \neq 0$) has been proved only recently by Beukers [10, Cor. 1.4] for arbitrary number fields, using André's theory of E -operators [4].

THEOREM B (Beukers). — *Let $\underline{f} = {}^t(f_1, \dots, f_N) \in \overline{\mathbb{Q}}[[z]]^N$ be a vector of E -functions which is solution of a differential system $\underline{f}' = A\underline{f}$ for some $A \in M_N(\overline{\mathbb{Q}}(z))$. Assume*

f_1, \dots, f_N are linearly independent over $\overline{\mathbb{Q}}(z)$ and that $z_0 \in \overline{\mathbb{Q}}^*$ is not a pole of an entry of A . Then the numbers $f_1(z_0), \dots, f_N(z_0)$ are linearly independent over $\overline{\mathbb{Q}}$.

The purpose of this paper is to prove new Diophantine results using this approach of André and Beukers. Our first main result is the following theorem, where we remove the rationality assumption on the coefficients and also the non-singularity assumption on z_0 . We recall that for a non-zero algebraic number α , its house $|\overline{\alpha}|$ is the maximum of the moduli of α and of all its Galois conjugates over \mathbb{Q} . We also denote by $\mathcal{O}_{\mathbb{K}}$ the ring of integers of a number field \mathbb{K} .

THEOREM 1. — *Let \mathbb{K} be a number field of degree d over \mathbb{Q} , $z_0 \in \mathbb{K}$, and $\underline{f} = {}^t(f_1, \dots, f_N)$ be a vector of E -functions with coefficients in \mathbb{K} such that $\underline{f}' = A\underline{f}$ for some $A \in M_N(\mathbb{K}(z))$. Then for any $\varepsilon > 0$, there exists $c = c(\varepsilon, \mathbb{K}, z_0, f_1, \dots, f_N) > 0$ with the following property.*

For any $\lambda_1, \dots, \lambda_N \in \mathcal{O}_{\mathbb{K}}$, if $\Lambda := \lambda_1 f_1(z_0) + \dots + \lambda_N f_N(z_0)$ is non-zero, then

$$|\Lambda| > cH^{-dN^d+1-\varepsilon} \quad \text{where } H := \max_{1 \leq j \leq N} |\lambda_j|.$$

REMARKS

– Given arbitrary E -functions f_1, \dots, f_N and any $z_0 \in \overline{\mathbb{Q}}$, this theorem applies because there exists a number field \mathbb{K} containing z_0 and all coefficients of f_1, \dots, f_N , and the family (f_1, \dots, f_N) can always be enlarged to satisfy a first-order differential system. This shows, without any assumption on f_1, \dots, f_N and z_0 , the existence of $\kappa, c > 0$ such that $|\Lambda| > cH^{-\kappa}$ provided $\Lambda \neq 0$.

– To our knowledge, Theorem 1 provides the first quantitative version of Beukers' Theorem B, when it is further assumed in Theorem 1 that f_1, \dots, f_N are linearly independent over $\mathbb{K}(z)$ and that $z_0 \in \mathbb{K}^*$ is not a pole of A , ensuring that $\Lambda \neq 0$.

– The constants c in Theorems A and 1, and their equivalents in other results of the paper, are in principle effective by a remark of André [3, footnote on p. 129] and by the multiplicity estimate of Bertrand-Beukers [8] made completely explicit by Bertrand-Chirskii-Yebbou [9]. However, we did not try to compute them because they would certainly be far from best possible and because we do not see any immediate application.

An important consequence of Theorem 1 is the following result, which completely settles the problem of deciding whether (real) values of E -functions can be Liouville numbers or not. We recall that a Liouville number is a real number ξ such that there exist two sequences of rational integers p_n, q_n such that $q_n \geq 2$ and $0 < |q_n \xi - p_n| < 1/q_n^n$ for all sufficiently large integers n (*a fortiori* $p_n q_n \neq 0$).

COROLLARY 1. — *Let f be an E -function, and z_0 be an algebraic number. Then $f(z_0)$ is not a Liouville number.*

The proof of Corollary 1 runs as follows: in Theorem 1, let $z_0 \in \overline{\mathbb{Q}}$, take $f_1 := 1$, $f_2 := f$ and consider a vector ${}^t(f_1, f_2, \dots, f_N)$ of E -functions with coefficients in a number field \mathbb{K} such that $\underline{f}' = A\underline{f}$ for some $A \in M_N(\mathbb{K}(z))$, where \mathbb{K} is large enough

to contain z_0 . If $f(z_0) \in \mathbb{Q}$, then $f(z_0)$ is not a Liouville number. If $f(z_0) \notin \mathbb{Q}$, then $\lambda_1 + \lambda_2 f(z_0) \neq 0$ for all $\lambda_1, \lambda_2 \in \mathbb{Z}$ not both zero, so that Theorem 1 yields $|\lambda_1 + \lambda_2 f(z_0)| > c \max(|\lambda_1|, |\lambda_2|)^{-\kappa}$ for some $c, \kappa > 0$. This rules out the possibility that $f(z_0)$ is a Liouville number.

Of course, Corollary 1 is interesting only when $f(z_0) \in \mathbb{R}$. If we do not assume this, note however that the real and imaginary parts of $f(z_0)$ are values of E -functions (see the remark in Section 2.1 below), so that none of them is a Liouville number.

Amongst other families of classical real numbers already proved not to be Liouville numbers, we find automatic numbers (Adamczewski-Cassaigne [1]) and Mahler numbers, i.e. the values of Mahler series in $\mathbb{Q}[[z]]$ at the inverse of rational integers (Bell-Bugeaud-Coons [7]). It is interesting to observe that the theories of E -functions and of Mahler series share many common properties, see also [2].

Let us also mention another interesting corollary, which is a consequence of Theorem 1 with $N = 2$ and $N = 3$ respectively (recall that $J_0(z)$ is solution of $zy''(z) + y'(z) + zy(z) = 0$).

COROLLARY 2. — *For any algebraic number $\alpha \in \overline{\mathbb{Q}}^*$ of degree d over \mathbb{Q} and any $\varepsilon > 0$, there exists $c = c(\alpha, \varepsilon)$ such that, for all $(p, q) \in \mathbb{Z} \times \mathbb{N}$ with $q \neq 0$,*

$$\left| e^\alpha - \frac{p}{q} \right| > \frac{c}{q^{d2^d + \varepsilon}}, \quad \text{respectively} \quad \left| J_0(\alpha) - \frac{p}{q} \right| > \frac{c}{q^{d3^d + \varepsilon}}.$$

The above mentioned general transcendence measure for E -functions due to Lang and Galochkin [14, p. 238, Th. 5.29 & remarks], refined by Kappe [18] for e^α in the special case of irrationality measures, gives $4d^2 - 2d$ instead of $d2^d$, and $16d^3 + 1$ instead of $d3^d$; see also [24, p. 404]. This is of course much better than Corollary 2 for large d but our bounds turn out to be smaller for $d \in \{2, 3\}$ and $d \in \{2, 3, 4, 5\}$ respectively. Note that Zudilin [26] has obtained the optimal exponent 2 for $J_0(\alpha)$ when $\alpha \in \overline{\mathbb{Q}}^*$.

A less classical example is the following: for any $\alpha \in \overline{\mathbb{Q}}^*$ and any integers $p, q \geq 1$, the value at $z = \alpha$ of $\mathcal{A}_{p,q}(z) := \sum_{n=0}^{\infty} (\sum_{k=0}^n \binom{n}{k}^p \binom{n+k}{n}^q) z^n / n!$ is not a Liouville number; $\mathcal{A}_{p,q}(\alpha)$ is proved to be a transcendental number in [11, §4.6] when $(p, q) \in \{1, 2, 3, 4\}^2$, the situation in general being unknown. More specifically, it is also proved in [11, §4.6, Table 1] that when $(p, q) = (2, 2)$, the minimal inhomogeneous differential equation over $\mathbb{Q}(z)$ satisfied by $\mathcal{A}_{2,2}$ is of order 4 and 0 is its only singularity. Hence, the following holds by Theorem 1 with $N = 5$: for all $\alpha \in \overline{\mathbb{Q}}^*$ of degree d over \mathbb{Q} and all $\varepsilon > 0$, there exists $c = c(\varepsilon, \alpha) > 0$ such that for all $\lambda_j \in \mathbb{Z}$ not all 0, we have

$$\left| \lambda_4 + \sum_{j=0}^3 \lambda_j \mathcal{A}_{2,2}^{(j)}(\alpha) \right| > c \max_{0 \leq j \leq 4} |\lambda_j|^{-d5^d + 1 - \varepsilon}.$$

In particular, for any $\alpha \in \overline{\mathbb{Q}}^*$ of degree d and $\varepsilon > 0$, there exists $c = c(\varepsilon, \alpha) > 0$ such that for any $(p, q) \in \mathbb{Z} \times \mathbb{N}$ with $q \neq 0$,

$$(1.1) \quad \left| \mathcal{A}_{2,2}(\alpha) - \frac{p}{q} \right| > \frac{c}{q^{d5^d + \varepsilon}}.$$

It seems reasonable to make the following conjecture, which probably belongs to folklore. This conjecture is Roth's Theorem if $f(z_0)$ is algebraic.

CONJECTURE 1. — *Let f be an E -function and $z_0 \in \overline{\mathbb{Q}}$. For any $\varepsilon > 0$, there exists $c > 0$ such that for any $(p, q) \in \mathbb{Z} \times \mathbb{N}$ with $q \neq 0$, either $qf(z_0) - p = 0$ or*

$$\left| f(z_0) - \frac{p}{q} \right| > \frac{c}{q^{2+\varepsilon}}.$$

Zudilin has proved this conjecture in [26] in a stronger form but under additional assumptions (which even imply $f(z_0) \notin \mathbb{Q}$), namely: f is an E -function with rational coefficients, $z_0 \in \mathbb{Q}^*$ is not a singularity of a differential system satisfied by $1, f, f_2, \dots, f_N$ over $\mathbb{Q}(z)$ for some $N \geq 2$ and f, f_2, \dots, f_N are algebraically independent over $\mathbb{Q}(z)$. It would be interesting to know if $\mathcal{A}_{2,2}, \mathcal{A}'_{2,2}, \mathcal{A}''_{2,2}, \mathcal{A}'''_{2,2}$ are algebraically independent over $\mathbb{Q}(z)$, in which case the exponent 5 could be improved to 2 in (1.1) when $\alpha \in \mathbb{Q}^*$.

With the notation of Theorem 1, for any $D \geq 1$ one may consider the $\binom{N+D-1}{N-1}$ functions $f_1^{i_1} \dots f_N^{i_N}$, with non-negative integers i_1, \dots, i_N such that $i_1 + \dots + i_N = D$. They make up a vector solution of a first order differential system, to which Theorem 1 applies. Taking $f_1 = 1$ and $f_2 = f$, this provides the following transcendence measure, valid for any value of an E -function at an algebraic point.

COROLLARY 3. — *Let f be an E -function, with coefficients in a number field \mathbb{K} of degree d , solution of an inhomogeneous linear differential equation of order $N - 1$. Let $z_0 \in \mathbb{K}$. Then for any $D \geq 1$, any $\varepsilon > 0$, and any $a_0, \dots, a_D \in \mathbb{Z}$, either $\sum_{j=0}^D a_j f(z_0)^j = 0$ or*

$$\left| \sum_{j=0}^D a_j f(z_0)^j \right| \geq c \left(\max(|a_0|, \dots, |a_D|) \right)^{-\kappa - \varepsilon},$$

where c depends only on f, z_0, D and ε , and $\kappa = d \binom{N+D-1}{N-1}^d - 1$.

With our notations, the transcendence measure of Lang-Galochkin provides the exponent $\frac{2^N(N-1)^{N-1}}{(N-1)!} d^N D^{N-1}$: for fixed N , it is smaller than κ when d and D are large but it is sometimes larger for small values of d or D as mentioned after Corollary 2, and moreover it holds under an algebraic independence assumption on the E -functions in the differential system. No such assumption is necessary in Corollary 3, which adds value to this result. Both results can be applied when f is the exponential function (with $N = 2$): we obtain a transcendence measure of e^{z_0} for any $z_0 \in \overline{\mathbb{Q}}^*$ of degree d with $\kappa = d(D+1)^d - 1$ whereas Lang-Galochkin's result yields $4d^2D$. Note that when $d = 1$ and $N = 2$ as for the exponential, we obtain $\kappa = D$ which is best possible. However this is not a new result for the exponential at rational points (see [6, Ch. 10], and [20, p. 135, Satz 3] when $z_0 = 1$).

Corollary 3 shows that no value of an E -function at an algebraic point can be a U -number, in Mahler's classification (see for instance [13, Ch. 3]). It seems reasonable to conjecture that it can neither be a T -number, so that values of E -functions at algebraic points would be either algebraic numbers or S -numbers.

The second goal of this paper is to understand the structure of the ring \mathbf{E} of all values at algebraic points of E -functions; this algebraic point can always be assumed to be 1 because if $f(z)$ is an E -function, so is $f(\alpha z)$ for any $\alpha \in \overline{\mathbb{Q}}$. Elements of \mathbf{E} are related to exponential periods (see [17, §4.3]).

For any subfield \mathbb{K} of $\overline{\mathbb{Q}}$, we shall also consider the subring $\mathbf{E}_{\mathbb{K}}$ of \mathbf{E} which consists of the evaluations $f(1)$ where f is an E -function with coefficients in \mathbb{K} (the number 1 could be replaced by any non-zero element of \mathbb{K} without changing $\mathbf{E}_{\mathbb{K}}$). Note that \mathbf{E} is the union of all $\mathbf{E}_{\mathbb{K}}$, where \mathbb{K} is a number field, since for any E -function f the holonomy property implies the existence of a number field that contains all coefficients of f .

We have defined and studied [15] analogous rings $\mathbf{G}_{\mathbb{K}}$ with G -functions instead of E -functions; it turns out that $\mathbf{G}_{\mathbb{K}}$ is nearly independent from \mathbb{K} (precisely, $\mathbf{G}_{\mathbb{K}} = \mathbf{G}_{\mathbb{Q}}$ if $\mathbb{K} \subset \mathbb{R}$, and $\mathbf{G}_{\mathbb{K}} = \mathbf{G}_{\mathbb{Q}(i)}$ otherwise). The situation is completely different for E -functions. A first hint in this direction was given in [16, Th. 4] (stated as Lemma 1 in §4 below). The way $\mathbf{E}_{\mathbb{K}}$ depends on \mathbb{K} is completely described by the following result.

THEOREM 2. — *Let \mathbb{K} be a subfield of $\overline{\mathbb{Q}}$. Elements of $\mathbf{E}_{\mathbb{K}}$ are linearly independent over $\overline{\mathbb{Q}}$ if, and only if, they are linearly independent over \mathbb{K} .*

We refer to [2, Th. 1.7] for a similar result concerning Mahler functions. Theorem 2 means that the \mathbb{K} -algebras $\mathbf{E}_{\mathbb{K}}$ and $\overline{\mathbb{Q}}$ are linearly disjoint, and the natural map $\mathbf{E}_{\mathbb{K}} \otimes_{\mathbb{K}} \overline{\mathbb{Q}} \rightarrow \mathbf{E}$ (sending $\xi \otimes z$ to ξz) is a \mathbb{K} -algebra isomorphism. The definition and properties of linearly disjoint algebras can be found in [12, Ch. V, §2, No. 5]; using Theorem 2 with $\mathbb{K} = \mathbb{Q}$ they imply the following.

COROLLARY 4. — *Let \mathbb{K} be a number field, and $(\omega_1, \dots, \omega_d)$ be a basis of the \mathbb{Q} -vector space \mathbb{K} . Then*

$$\mathbf{E}_{\mathbb{K}} = \omega_1 \mathbf{E}_{\mathbb{Q}} \oplus \dots \oplus \omega_d \mathbf{E}_{\mathbb{Q}}.$$

In other words, for any $\xi \in \mathbf{E}_{\mathbb{K}}$ there exists a unique d -tuple $(\xi_1, \dots, \xi_d) \in \mathbf{E}_{\mathbb{Q}}^d$ such that $\xi = \omega_1 \xi_1 + \dots + \omega_d \xi_d$.

The original definition of E -functions, given by Siegel [25], is slightly less restrictive: instead of geometric bounds, he allowed growths bounded by $n!^{\varepsilon}$ (for any given $\varepsilon > 0$, provided n is large enough with respect to ε). Shidlovskii's Theorem A holds for E -functions in Siegel's sense, and Beukers' Theorem B was later proved by André [5] in this general setting, by a different method. All other results in Beukers' paper [10] have been adapted by Lepetit [19]. Therefore all the results of the present paper also hold for E -functions in Siegel's sense.

The structure of this paper is as follows. In §2 we prove Proposition 1, which is crucial in the proof of both Theorems 1 and 2. We also derive from it a version of Beukers' desingularization process over a number field. This enables us to prove Theorem 1 in §3, and also to obtain in §4 a decomposition of an E -function over a number field, involving an E -function that takes only transcendental values at non-zero algebraic points. At last we apply the previous results in §5 to study the structure of $\mathbf{E}_{\mathbb{K}}$ and prove Theorem 2. The final section is devoted to an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$

on the set of values of E -functions; it is not used in the paper but it illustrates how our method presents similarities with Liouville's theorem.

Acknowledgements. — We warmly thank the referees for their very constructive comments.

2. MAIN TOOLS

2.1. CONJUGATES OF E -FUNCTIONS. — Let $f(z) = \sum_{n=0}^{\infty} a_n z^n / n!$ be an E -function with coefficients $a_n \in \overline{\mathbb{Q}}$. For any $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ we let $f^\sigma(z) = \sum_{n=0}^{\infty} \sigma(a_n) z^n / n!$. The definition shows that f^σ is also an E -function, and if g is an E -function then for any σ, τ we have $(f + g)^\sigma = f^\sigma + g^\sigma$, $(fg)^\sigma = f^\sigma g^\sigma$ and $(f^\sigma)^\tau = f^{\tau \circ \sigma}$. Moreover if f has coefficients in a number field \mathbb{K} , then f^σ has coefficients in the number field $\sigma(\mathbb{K})$.

REMARK. — Denoting by σ the complex conjugation, for any E -function f we can consider $\frac{1}{2}(f + f^\sigma)$ and $\frac{1}{2i}(f - f^\sigma)$. These E -functions have real coefficients, which are respectively the real and imaginary parts of those of f . In particular, the real and imaginary parts of any element of \mathbf{E} belong to \mathbf{E} .

The following result is central in the present paper; we refer to [11, Prop. 3.5] for a similar result. We agree that minimal polynomials of algebraic elements have leading coefficient 1.

PROPOSITION 1. — *Let f be an E -function with coefficients in a number field \mathbb{K} , and $z_0 \in \overline{\mathbb{Q}}^*$. Then the following assertions are equivalent:*

- (i) f vanishes at z_0 .
- (ii) There exists $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ such that f^σ vanishes at $\sigma(z_0)$.
- (iii) For any $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, f^σ vanishes at $\sigma(z_0)$.
- (iv) There exists an E -function g with coefficients in \mathbb{K} such that

$$f(z) = D(z)g(z) \text{ where } D \text{ is the minimal polynomial of } z_0 \text{ over } \mathbb{K}.$$

In particular, if z_0 is rational and f vanishes at z_0 , then all conjugates f^σ of f also vanish at z_0 . Also, if an E -function f with rational coefficients vanishes at some $z_0 \in \overline{\mathbb{Q}}^*$, then it vanishes at all Galois conjugates of z_0 .

We remark that with $z_0 = 1$, the implication (i) \Rightarrow (iii) is used already in the proof of [10, Prop. 4.1], which is the main result Proposition 1 is based on.

Proof of Proposition 1

(iv) \Rightarrow (iii) Let $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Then $f^\sigma(z) = D^\sigma(z)g^\sigma(z)$, and $D^\sigma(\sigma(z_0)) = \sigma(D(z_0)) = 0$. Therefore $f^\sigma(\sigma(z_0)) = 0$.

(iii) \Rightarrow (ii) is trivial.

(ii) \Rightarrow (i) Enlarging \mathbb{K} if necessary, we may assume the extension \mathbb{K}/\mathbb{Q} to be Galois and to contain z_0 . Then f^σ has coefficients in \mathbb{K} , and $\sigma(z_0) \in \mathbb{K}^*$. Using [10, Prop. 4.1] there exists an E -function g such that $f^\sigma(z) = (z - \sigma(z_0))g(z)$. Then g has coefficients in \mathbb{K} ; applying σ^{-1} yields $f(z) = (z - z_0)g^{\sigma^{-1}}(z)$ so that $f(z_0) = 0$.

(i) \Rightarrow (iv) By using [10, Prop. 4.1], one shows that there exists an E -function h such that $f(z) = (z - z_0)h(z)$. Let $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{K})$, that is: σ is a field automorphism of $\overline{\mathbb{Q}}$ such that $\sigma(x) = x$ for any $x \in \mathbb{K}$. Then $f(z) = f^\sigma(z) = (z - \sigma(z_0))h^\sigma(z)$, so that f vanishes at $\sigma(z_0)$. Let $z_1 := z_0, z_2, \dots, z_\ell$ denote the (pairwise distinct) Galois conjugates of z_0 over \mathbb{K} , i.e. the elements of the form $\sigma(z_0)$ with $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{K})$; we have proved that f vanishes at z_1, \dots, z_ℓ . Applying [10, Prop. 4.1] yields, by induction on $j \in \{1, \dots, \ell\}$, the existence of an E -function g_j such that $f(z) = g_j(z) \prod_{i=1}^j (z - z_i)$. Since $D(z) = \prod_{i=1}^\ell (z - z_i)$, we have $f(z) = D(z)g_\ell(z)$. Now, $D(z) \in \mathbb{K}[z] \setminus \{0\}$, so that all coefficients of g_ℓ belong to \mathbb{K} . This concludes the proof of Proposition 1. \square

2.2. BEUKERS' DESINGULARIZATION PROCESS. — In the proof of Theorem 1 we shall use the following version of Beukers' desingularization theorem [10, Th. 1.5].

PROPOSITION 2. — *Let \mathbb{K} be a number field, and f_1, \dots, f_N be E -functions with coefficients in \mathbb{K} , linearly independent over $\mathbb{C}(z)$. Assume that the vector $\underline{f} = {}^t(f_1, \dots, f_N)$ satisfies a first-order differential system $\underline{f}' = A\underline{f}$ with $A \in M_N(\mathbb{K}(z))$.*

Then there exist E -functions e_1, \dots, e_N with coefficients in \mathbb{K} , linearly independent over $\mathbb{C}(z)$, a matrix $B \in M_N(\mathbb{K}[z, 1/z])$ and a matrix $M \in M_N(\mathbb{K}[z])$, such that with $\underline{e} = {}^t(e_1, \dots, e_N)$:

$$\underline{e}' = B\underline{e} \quad \text{and} \quad \underline{f} = M\underline{e}.$$

The new point is that e_1, \dots, e_N and the coefficients of B and M have coefficients in the number field \mathbb{K} (whereas in [10, Th. 1.5] these are simply algebraic numbers).

The proof of Proposition 2 follows [10, p. 378], using also the additional details given in [11]. Actually Proposition 2 is already proved implicitly (for $\mathbb{K} = \mathbb{Q}$) by the implementation described in [11].

In what follows we only mention the parts of the proof where special attention has to be paid. Let α be a singularity of the differential system $Y' = AY$, and $Q \in \mathbb{K}[X]$ denote the minimal polynomial of α over \mathbb{K} . Let $k \geq 1$ be the maximal order of α as a pole of a coefficient of A , and (i_0, j_0) be such that A_{i_0, j_0} has a pole of order exactly k at α . Then $Q^k A \underline{f} = Q^k \underline{f}'$ vanishes at α ; the i_0 -th coordinate of this vector provides a linear relation

$$\sum_{j=1}^N (Q^k A_{i_0, j})(\alpha) f_j(\alpha) = 0.$$

Note that for any j , the rational function $Q(z)^k A_{i_0, j}(z) \in \mathbb{K}(z)$ is holomorphic at α , and for $j = j_0$ it does not vanish at that point. Multiplying by the least common denominator of the rational functions $Q(z)^k A_{i_0, j}(z)$ we obtain coprime polynomials $P_1, \dots, P_N \in \mathbb{K}[z]$ such that

$$P_{j_0}(\alpha) \neq 0 \quad \text{and} \quad \sum_{j=1}^N P_j(\alpha) f_j(\alpha) = 0.$$

If $N = 1$ we let $P_{1,1} = 1$; otherwise there exist polynomials $P_{i,j} \in \mathbb{K}[z]$, for $2 \leq i \leq N$ and $1 \leq j \leq N$, such that letting $P_{1,j} = P_j$, the matrix $S = (P_{i,j})_{1 \leq i, j \leq N} \in M_N(\mathbb{K}[z])$

has determinant 1. Then Sf is a vector of E -functions, with coefficients in \mathbb{K} , of which the first coordinate $\sum_{j=1}^N P_j(z)f_j(z)$ vanishes at α . Using Proposition 1, we deduce that $\sum_{j=1}^N P_j(z)f_j(z)$ vanishes at $\sigma(\alpha)$ for any $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{K})$. This concludes the proof as in [10].

3. PROOF OF THEOREM 1

To prove Theorem 1 we assume $z_0 \neq 0$ since otherwise, $f_1(z_0), \dots, f_N(z_0)$ are algebraic numbers and the conclusion follows from Schmidt's subspace theorem (see for instance [14, Ch. 1, §8.2, Th. 1.37]). Considering the E -functions $f_j(z_0 z)$ instead of $f_j(z)$, we may even assume $z_0 = 1$. We also suppose that $\Lambda \neq 0$.

The proof of Theorem 1 falls into 2 steps.

Step 1. — Let us prove Theorem 1 in the special case where $\mathbb{K} = \mathbb{Q}$ (i.e., $d = 1$). In other words, we assume that $z_0 = 1 \in \mathbb{Q}$, $\lambda_1, \dots, \lambda_N \in \mathbb{Z}$, and f_1, \dots, f_N have coefficients in \mathbb{Q} .

Recall that we do not assume that $f_1(1), \dots, f_N(1)$ are linearly independent over \mathbb{Q} . Let N' be the maximal number of linearly independent numbers among them. We may assume (up to a permutation of the indices) that $f_1(1), \dots, f_{N'}(1)$ are linearly independent over \mathbb{Q} , and $f_{N'+1}(1), \dots, f_N(1)$ belong to the \mathbb{Q} -vector space they span. There exist rational numbers $\varrho_{i,j}$ such that $f_j(1) = \sum_{i=1}^{N'} \varrho_{i,j} f_i(1)$ for any $1 \leq j \leq N$, so that

$$(3.1) \quad \Lambda := \sum_{j=1}^N \lambda_j f_j(1) = \sum_{i=1}^{N'} \mu_i f_i(1) \quad \text{with} \quad \mu_i := \sum_{j=1}^N \lambda_j \varrho_{i,j} \in \mathbb{Q}.$$

Observe that the E -functions $f_1, \dots, f_{N'}$ are linearly independent over $\mathbb{C}(z)$. Indeed, otherwise they would be linearly dependent over $\mathbb{Q}(z)$ (since they have coefficients in \mathbb{Q}), and a relation $\sum_{j=1}^{N'} S_j(z)f_j(z) = 0$ would exist with $S_1, \dots, S_{N'} \in \mathbb{Q}(z)$ not all zero. Upon multiplying by $(z-1)^k$ for a suitable $k \in \mathbb{Z}$, we may assume that none of the S_j has a pole at 1, and that at least one of them does not vanish at 1. This provides a non-trivial linear relation $\sum_{j=1}^{N'} S_j(1)f_j(1) = 0$, which contradicts the definition of N' .

Therefore $f_1, \dots, f_{N'}$ are linearly independent over $\mathbb{C}(z)$. Denote by N'' the dimension of the vector space generated over $\mathbb{C}(z)$ by f_1, \dots, f_N ; we have $N' \leq N'' \leq N$. Notice that it could happen that $N'' > N'$, for instance if $f_{N'+1}(1) = 0$. Up to a permutation of the indices, we may assume that $f_1, \dots, f_{N''}$ are linearly independent over $\mathbb{C}(z)$, and that $f_{N''+1}, \dots, f_N$ belong to the vector space they span over $\mathbb{C}(z)$.

Since $f_1, \dots, f_{N''}$ are linearly independent over $\mathbb{C}(z)$, and satisfy a linear differential system of order 1 by definition of N'' , Proposition 2 (applied with $\mathbb{K} = \mathbb{Q}$) provides E -functions $e_1, \dots, e_{N''}$ with rational coefficients and matrices $B \in M_{N''}(\mathbb{Q}[z, 1/z])$ and $M = (P_{i,j}) \in M_{N''}(\mathbb{Q}[z])$ such that $\underline{e}' = B\underline{e}$ and $f_i(z) = \sum_{j=1}^{N''} P_{i,j}(z)e_j(z)$.

Since $N' \leq N''$, Equation (3.1) yields

$$(3.2) \quad \Lambda = \sum_{j=1}^{N''} \nu_j e_j(1) \quad \text{with} \quad \nu_j := \sum_{i=1}^{N'} \mu_i P_{i,j}(1) \in \mathbb{Q}.$$

Now, Shidlovskii's lower bound stated as Theorem A in the introduction applies to the E -functions $e_1, \dots, e_{N''}$ with rational coefficients, which are linearly independent over $\mathbb{C}(z)$ and solution of a linear differential system of order 1 of which 1 is not a singularity. Denoting by δ a common denominator of the rational numbers $P_{i,j}(1)$ and $\varrho_{i,j}$ (appearing in Equations (3.1) and (3.2)), we obtain that $\delta^2 \Lambda$ is a \mathbb{Z} -linear combination of $e_1(1), \dots, e_{N''}(1)$ with coefficients bounded (in absolute value) by cH , where $c > 0$ and δ depend only on f_1, \dots, f_N . For any $\varepsilon > 0$, Theorem A yields $|\delta^2 \Lambda| > c_0 H^{-N''+1-\varepsilon} \geq c_0 H^{-N+1-\varepsilon}$ for some $c_0 > 0$ which depends only on f_1, \dots, f_N and ε . This concludes the proof of Theorem 1 in the case where $\mathbb{K} = \mathbb{Q}$.

Step 2. — Let us prove Theorem 1 for any number field \mathbb{K} . For simplicity of the exposition, we assume \mathbb{K} to be a Galois extension of \mathbb{Q} ; see the end of Step 2 for the general case. We denote by G the Galois group of \mathbb{K}/\mathbb{Q} , and consider the complex number

$$(3.3) \quad \varpi := \prod_{\sigma \in G} \left(\sum_{j=1}^N \sigma(\lambda_j) f_j^\sigma(1) \right).$$

To begin with, let us prove that $\varpi \neq 0$. Indeed, consider the E -function $g(z) = \sum_{j=1}^N \lambda_j f_j(z)$; it has coefficients in \mathbb{K} (because f_1, \dots, f_N do), and $g(1) = \Lambda \neq 0$. For any $\sigma \in G$, Proposition 1 yields $g^\sigma(1) \neq 0$. Now, $g^\sigma(1) = \sum_{j=1}^N \sigma(\lambda_j) f_j^\sigma(1)$, so that $\varpi = \prod_{\sigma \in G} g^\sigma(1) \neq 0$.

Denote by \mathcal{J} the set of all tuples of integers $\underline{i} = (i_\sigma)_{\sigma \in G}$ such that $1 \leq i_\sigma \leq N$ for any $\sigma \in G$. Expanding the product in the definition (3.3) of ϖ yields

$$(3.4) \quad \varpi = \sum_{\underline{i} \in \mathcal{J}} \prod_{\sigma \in G} \sigma(\lambda_{i_\sigma}) f_{i_\sigma}^\sigma(1) = \sum_{\underline{i} \in \mathcal{J}} \left(\prod_{\sigma \in G} \sigma(\lambda_{i_\sigma}) \right) g_{\underline{i}}(1),$$

where

$$(3.5) \quad g_{\underline{i}}(z) := \prod_{\sigma \in G} f_{i_\sigma}^\sigma(z)$$

is an E -function with coefficients in \mathbb{K} .

The normal basis theorem (see for instance [21, Th.5.18]) provides an element $\alpha \in \mathbb{K}$ such that the $\sigma(\alpha)$, for $\sigma \in G$, make up a basis of the \mathbb{Q} -vector space \mathbb{K} . Upon multiplying α by a suitable positive integer, we may assume that $\alpha \in \mathcal{O}_{\mathbb{K}}$ (so that $\sigma(\alpha) \in \mathcal{O}_{\mathbb{K}}$ for any $\sigma \in G$). Expanding all coefficients of $g_{\underline{i}}(z)$ in this basis yields (using [24, Ch. 3, Lem. 12]) E -functions $g_{\underline{i},\sigma}(z)$ with coefficients in \mathbb{Q} , for $\sigma \in G$, such that

$$(3.6) \quad g_{\underline{i}}(z) = \sum_{\sigma \in G} \sigma(\alpha) g_{\underline{i},\sigma}(z) \quad \text{for any } \underline{i} \in \mathcal{J}.$$

In the sequel, it is important to observe that these E -functions $g_{\underline{i},\sigma}(z)$ are uniquely determined by $g_{\underline{i}}(z)$, since for each n their coefficients of z^n are given by the expansion in the basis $(\sigma(\alpha))_{\sigma \in G}$ of the corresponding coefficient of $g_{\underline{i}}(z)$.

For $\underline{i} \in \mathcal{I}$ and $\sigma \in G$, we denote by $\sigma(\underline{i})$ the tuple $\underline{j} \in \mathcal{I}$ defined by $j_\tau = i_{\sigma \circ \tau}$ for any $\tau \in G$. Let us prove that

$$(3.7) \quad g_{\underline{i},\sigma}(z) = g_{\sigma(\underline{i}),\text{Id}}(z) \quad \text{for any } \underline{i} \in \mathcal{I} \text{ and any } \sigma \in G.$$

Indeed we have:

$$\begin{aligned} \sum_{\tau \in G} \tau(\alpha) g_{\sigma(\underline{i}),\tau}(z) &= g_{\sigma(\underline{i})}(z) \text{ using Equation (3.6)} \\ &= \prod_{\tau \in G} f_{i_{\sigma \circ \tau}}^\tau(z) \text{ by definition of } g_{\sigma(\underline{i})}(z) \\ &= \prod_{\tau' \in G} f_{i_{\tau'}}^{\sigma^{-1} \circ \tau'}(z) \quad \text{by letting } \tau' = \sigma \circ \tau \\ &= \left(\prod_{\tau' \in G} f_{i_{\tau'}}^{\tau'} \right)^{\sigma^{-1}}(z) \\ &= g_{\underline{i}}^{\sigma^{-1}}(z) \\ &= \sum_{\tau \in G} \sigma^{-1}(\tau(\alpha)) g_{\underline{i},\tau}(z) \quad \text{since } g_{\underline{i},\tau} \text{ has coefficients in } \mathbb{Q}. \end{aligned}$$

Comparing the coefficient of α on both sides yields Equation (3.7) since an expansion for $g_{\sigma(\underline{i})}(z)$ like the one of Equation (3.6) is unique.

Let us prove now that the vector of E -functions $g_{\underline{i},\text{Id}}(z)$, for $\underline{i} \in \mathcal{I}$, is solution of a first-order linear differential system. By assumption we have $f'_i(z) = \sum_{j=1}^N A_{i,j}(z) f_j(z)$ with $A_{i,j} \in \mathbb{K}(z)$, so that

$$(f_i^\sigma)'(z) = (f_i')^\sigma(z) = \sum_{j=1}^N \sigma(A_{i,j})(z) f_j^\sigma(z)$$

and

$$\begin{aligned} g_{\underline{i}}'(z) &= \sum_{\sigma \in G} (f_{i_\sigma}^\sigma)'(z) \prod_{\tau \neq \sigma} f_{i_\tau}^\tau(z) \\ &= \sum_{\sigma \in G} \sum_{j=1}^N \sigma(A_{i_\sigma,j})(z) f_j^\sigma(z) \prod_{\tau \neq \sigma} f_{i_\tau}^\tau(z) \\ &= \sum_{\underline{i}' \in \mathcal{I}} B_{\underline{i},\underline{i}'}(z) g_{\underline{i}'}(z) \quad \text{for some } B_{\underline{i},\underline{i}'}(z) \in \mathbb{K}(z) \\ &= \sum_{\underline{i}' \in \mathcal{I}} B_{\underline{i},\underline{i}'}(z) \sum_{\sigma \in G} \sigma(\alpha) g_{\sigma(\underline{i}'),\text{Id}}(z) \quad \text{using Equations (3.6) and (3.7)} \\ &= \sum_{\underline{i}'' \in \mathcal{I}} C_{\underline{i},\underline{i}''}(z) g_{\underline{i}'',\text{Id}}(z) \quad \text{for some } C_{\underline{i},\underline{i}''}(z) \in \mathbb{K}(z). \end{aligned}$$

Each $C_{\underline{i}, \underline{i}''}(z) \in \mathbb{K}(z)$ can be written as $N_{\underline{i}, \underline{i}''}(z)/D_{\underline{i}, \underline{i}''}(z)$ with $D_{\underline{i}, \underline{i}''}(z) \in \mathbb{Q}[z] \setminus \{0\}$ and $N_{\underline{i}, \underline{i}''}(z) \in \mathbb{K}[z]$. Writing $N_{\underline{i}, \underline{i}''}(z)$ as a $\mathbb{Q}[z]$ -linear combination of the $\sigma(\alpha)$, $\sigma \in G$, yields an expression

$$(3.8) \quad g'_{\underline{i}}(z) = \sum_{\sigma \in G} \sigma(\alpha) \sum_{\underline{i}'' \in \mathcal{J}} R_{\underline{i}, \underline{i}'', \sigma}(z) g_{\underline{i}'', \text{Id}}(z)$$

with $R_{\underline{i}, \underline{i}'', \sigma}(z) \in \mathbb{Q}(z)$. On the other hand, Equation (3.6) yields

$$(3.9) \quad g'_{\underline{i}}(z) = \sum_{\sigma \in G} \sigma(\alpha) g'_{\underline{i}, \sigma}(z).$$

Comparing the components on α of Equations (3.8) and (3.9), unicity of such an expression yields

$$g'_{\underline{i}, \text{Id}}(z) = \sum_{\underline{i}'' \in \mathcal{J}} R_{\underline{i}, \underline{i}'', \text{Id}}(z) g_{\underline{i}'', \text{Id}}(z).$$

This concludes the proof that the vector of E -functions $g_{\underline{i}, \text{Id}}(z)$, $\underline{i} \in \mathcal{J}$, satisfies a first-order linear differential system with coefficients in $\mathbb{Q}(z)$.

Now, we come back to ϖ : combining Equations (3.4), (3.6) and (3.7) yields

$$(3.10) \quad \varpi = \sum_{\underline{i} \in \mathcal{J}} \left(\prod_{\sigma \in G} \sigma(\lambda_{i_\sigma}) \right) \sum_{\tau \in G} \tau(\alpha) g_{\tau(\underline{i}), \text{Id}}(1) = \sum_{\underline{i}' \in \mathcal{J}} \kappa_{\underline{i}'} g_{\underline{i}', \text{Id}}(1)$$

upon letting

$$\kappa_{\underline{i}'} = \sum_{\tau \in G} \tau(\alpha) \prod_{\sigma \in G} \sigma(\lambda_{i'_{\tau^{-1} \circ \sigma}}) \in \mathbb{K}.$$

Let us prove that $\kappa_{\underline{i}} \in \mathbb{Z}$ for any $\underline{i} \in \mathcal{J}$. Indeed for any $\gamma \in G$, we have:

$$\gamma(\kappa_{\underline{i}}) = \sum_{\tau \in G} \gamma(\tau(\alpha)) \prod_{\sigma \in G} \gamma(\sigma(\lambda_{i_{\tau^{-1} \circ \sigma}})) = \sum_{\tau' \in G} \tau'(\alpha) \prod_{\sigma' \in G} \sigma'(\lambda_{i_{\tau'^{-1} \circ \sigma'}}) = \kappa_{\underline{i}}$$

by letting $\tau' = \gamma \circ \tau$ and $\sigma' = \gamma \circ \sigma$, since $\tau'^{-1} \circ \sigma' = \tau^{-1} \circ \sigma$. Using the fact that all $\tau(\alpha)$ and all λ_j belong to $\mathcal{O}_{\mathbb{K}}$, we deduce that $\kappa_{\underline{i}} \in \mathcal{O}_{\mathbb{K}} \cap \mathbb{Q} = \mathbb{Z}$.

To sum up, Equation (3.10) shows that ϖ is a \mathbb{Z} -linear combination of the values at 1 of a family of cardinality $\text{Card}(\mathcal{J}) = N^d$ of E -functions with coefficients in \mathbb{Q} , solution of a first order differential system. Therefore Step 1 applies with $H' := H^d \sum_{\tau \in G} |\tau(\alpha)|$, since $|\kappa_{\underline{i}}| \leq H'$ for any \underline{i} . We obtain $|\varpi| > cH^{-dN^d+d-\varepsilon}$ for any $\varepsilon > 0$, where $c > 0$ depends on ε . Now, Equation (3.3) yields $|\varpi| \leq c'H^{d-1}|\Lambda|$ by bounding trivially the factors corresponding to all $\sigma \neq \text{Id}$; here c' is a positive constant that depends only on f_1, \dots, f_N and \mathbb{K} . Combining these estimates yields $|\Lambda| > c''H^{-dN^d+1-\varepsilon}$ for some constant c'' ; this concludes Step 2 in the case where \mathbb{K} is a Galois extension of \mathbb{Q} .

If \mathbb{K}/\mathbb{Q} (of degree d) is not assumed to be Galois, we consider a finite Galois extension \mathbb{L} of \mathbb{Q} such that $\mathbb{K} \subset \mathbb{L}$. We now explain the changes that must be made to the above construction. We let $G_0 = \text{Gal}(\mathbb{L}/\mathbb{Q})$ and $H = \text{Gal}(\mathbb{L}/\mathbb{K})$. In the definition of ϖ , namely Equation (3.3), the product is now taken over the d cosets $\sigma \in G_0/H$; indeed $\sigma(\lambda_j)$ and f_j^σ are the same for all σ in a given coset, because λ_j and the coefficients of f_j belong to \mathbb{K} . In the products of Equations (3.4) and (3.5), σ ranges through G_0/H , and $\mathcal{J} = \{1, \dots, N\}^{G_0/H}$. However the normal basis theorem is applied to the Galois

extension \mathbb{L}/\mathbb{Q} , so that $\alpha \in \mathcal{O}_{\mathbb{L}}$ and σ ranges through G_0 in Equations (3.6) to (3.9). In Equation (3.10), the product is over $\sigma \in G_0/H$ and the sum over $\tau \in G_0$. We have $\kappa_{\underline{i}} \in \mathbb{L}$, and deduce that $\kappa_{\underline{i}} \in \mathbb{Q}$ since $\gamma(\kappa_{\underline{i}}) = \kappa_{\underline{i}}$ for any $\gamma \in G_0$. We conclude the proof in the same way since $\text{Card}(\mathcal{J}) = N^d$.

Alternative proof of Step 2 if $\lambda_j \in \mathbb{Z}$ for any j . — For the reader's convenience we give now a slightly different proof in this special case. It is based on the same idea of considering ϖ , but its expansion and the way Step 1 is applied are not the same. As in Step 2, we assume \mathbb{K}/\mathbb{Q} to be Galois (the general case is dealt with as explained at the end of Step 2), let $G = \text{Gal}(\mathbb{K}/\mathbb{Q})$ and consider

$$(3.11) \quad \varpi = \prod_{\sigma \in G} \left(\sum_{j=1}^N \lambda_j f_j^\sigma(1) \right)$$

since we have $\sigma(\lambda_j) = \lambda_j$ now; we still have $\varpi \neq 0$. To expand the product in the definition of ϖ , we denote by \mathcal{N} the set of all tuples $\underline{n} = (n_1, \dots, n_N)$ of non-negative integers such that $n_1 + \dots + n_N = d$. For any $\underline{n} \in \mathcal{N}$, we denote by $I(\underline{n})$ the set of all tuples $\underline{i} = (i_\sigma)_{\sigma \in G}$ consisting of integers $i_\sigma \in \{1, \dots, N\}$ such that for any $j \in \{1, \dots, N\}$ we have:

$$\text{Card}\{\sigma \in G, i_\sigma = j\} = n_j.$$

Then Equation (3.11) yields

$$(3.12) \quad \varpi = \sum_{\underline{n} \in \mathcal{N}} \lambda_1^{n_1} \cdots \lambda_N^{n_N} \varphi_{\underline{n}}(1) \quad \text{upon letting} \quad \varphi_{\underline{n}}(z) := \sum_{\underline{i} \in I(\underline{n})} \prod_{\sigma \in G} f_{i_\sigma}^\sigma(z).$$

Let us prove that $\varphi_{\underline{n}}(z)$, which is an E -function with coefficients in \mathbb{K} , actually has coefficients in \mathbb{Q} for any $\underline{n} \in \mathcal{N}$. For any $\tau \in G$ we have:

$$\begin{aligned} \varphi_{\underline{n}}^\tau &= \sum_{\underline{i} \in I(\underline{n})} \prod_{\sigma \in G} \left(f_{i_\sigma}^\sigma \right)^\tau \\ &= \sum_{\underline{i} \in I(\underline{n})} \prod_{\sigma \in G} f_{i_\sigma}^{\tau \circ \sigma} \\ &= \sum_{\underline{i} \in I(\underline{n})} \prod_{\sigma' \in G} f_{i_{\tau^{-1} \circ \sigma'}}^{\sigma'} \quad \text{by letting } \sigma' = \tau \circ \sigma \\ &= \sum_{\underline{i}' \in I(\underline{n})} \prod_{\sigma' \in G} f_{i'_{\sigma'}}^{\sigma'}, \end{aligned}$$

where the last equality comes from letting $i'_\sigma = i_{\tau^{-1} \circ \sigma}$ for any $\sigma \in G$; indeed this defines a bijective map $I(\underline{n}) \rightarrow I(\underline{n})$. Therefore $\varphi_{\underline{n}}^\tau = \varphi_{\underline{n}}$ for any $\tau \in G$, and the E -function $\varphi_{\underline{n}}(z)$ has coefficients in \mathbb{Q} .

We denote by \mathcal{E} the vector space spanned over $\overline{\mathbb{Q}}(z)$ by the functions $\prod_{\sigma \in G} f_{i_\sigma}^\sigma$ for all tuples $\underline{i} = (i_\sigma)_{\sigma \in G}$ consisting of integers $i_\sigma \in \{1, \dots, N\}$. There are N^d such tuples, so $\dim(\mathcal{E}) \leq N^d$. Moreover we have $g' \in \mathcal{E}$ for any $g \in \mathcal{E}$.

Let δ denote the dimension of the vector space spanned over $\mathbb{Q}(z)$ by the functions $\varphi_{\underline{n}}$ for $\underline{n} \in \mathcal{N}$. We can choose δ functions h_1, \dots, h_δ among the $\varphi_{\underline{n}}$ which are linearly independent, and span the same $\mathbb{Q}(z)$ -vector space. Choosing among the

successive derivatives of h_1, \dots, h_δ it is possible to find an integer $\delta' \geq \delta$ and functions h_i , for $\delta + 1 \leq i \leq \delta'$, such that $h_1, \dots, h_{\delta'}$ are linearly independent over $\mathbb{Q}(z)$ and satisfy a linear differential system of order 1. Since they have rational coefficients, they are also linearly independent over $\overline{\mathbb{Q}}(z)$; now they all belong to \mathcal{E} , so we have $\delta' \leq \dim(\mathcal{E}) \leq N^d$.

Proposition 2 with $\mathbb{K} = \mathbb{Q}$ yields a vector of E -functions $e_1, \dots, e_{\delta'}$ with rational coefficients, solution of a first-order differential system with no finite non-zero singularity, such that each h_i is a linear combination of $e_1, \dots, e_{\delta'}$ with coefficients in $\mathbb{Q}[z]$. There exist $R_{\underline{n},i}, S_{\underline{n},i} \in \mathbb{Q}(z)$ for $\underline{n} \in \mathcal{N}$ and $1 \leq i \leq \delta'$ such that, for any \underline{n} ,

$$\varphi_{\underline{n}}(z) = \sum_{i=1}^{\delta'} R_{\underline{n},i}(z) h_i(z) = \sum_{i=1}^{\delta'} S_{\underline{n},i}(z) e_i(z).$$

If no $S_{\underline{n},i}$ has a pole at $z = 1$, we can take $z = 1$ in this equation. To deal with the general case, we expand the right hand side as a polynomial in $1/(z - 1)$, up to an additive term which is holomorphic and vanishes at $z = 1$. Since $\varphi_{\underline{n}}(z)$ is holomorphic at 1, all polar contributions cancel out and the value at $z = 1$ is given by the constant term of the above-mentioned polynomial. This provides an expression of the form

$$\varphi_{\underline{n}}(1) = \sum_{i=1}^{\delta'} \sum_{j=0}^J a_{\underline{n},i,j} e_i^{(j)}(1)$$

with $a_{\underline{n},i,j} \in \mathbb{Q}$. Since ${}^t(e_1, \dots, e_{\delta'})$ is solution of a first-order differential system with coefficients in $\mathbb{Q}[z, 1/z]$, hence with no finite non-zero singularity, we obtain finally

$$(3.13) \quad \varphi_{\underline{n}}(1) = \sum_{i=1}^{\delta'} b_{\underline{n},i} e_i(1)$$

with $b_{\underline{n},i} \in \mathbb{Q}$ (where simply $b_{\underline{n},i} := S_{\underline{n},i}(1)$ in the “no pole at $z = 1$ ” case considered above). Using Equation (3.13) into Equation (3.12) yields

$$\varpi = \sum_{i=1}^{\delta'} \mu_i e_i(1) \quad \text{with} \quad \mu_i = \sum_{\underline{n} \in \mathcal{N}} \lambda_1^{n_1} \cdots \lambda_N^{n_N} b_{\underline{n},i} \in \mathbb{Q}.$$

This enables us to apply the special case of Theorem 1 where $\mathbb{K} = \mathbb{Q}$, proved in Step 1, with N replaced with $\delta' \leq N^d$. Indeed we denote by $\alpha \in \mathbb{Z}$ a common positive denominator of the rational numbers $b_{\underline{n},i}$; then we have $\alpha\mu_1, \dots, \alpha\mu_{\delta'} \in \mathbb{Z}$. Since $\varpi \neq 0$ we obtain $|\alpha\varpi| > cH'^{-N^d+1-\varepsilon}$ with

$$H' = \max_{1 \leq i \leq \delta'} |\alpha\mu_i| \leq \beta \max_{\underline{n} \in \mathcal{N}} |\lambda_1^{n_1} \cdots \lambda_N^{n_N}| \leq \beta H^d,$$

where $\beta > 0$ depends only on f_1, \dots, f_N and \mathbb{K} . Then we conclude the proof as in Step 2.

4. DECOMPOSITION OF E -FUNCTIONS OVER A NUMBER FIELD

In the same spirit as Proposition 1, it is possible to prove the following result. The weaker version with \mathbb{K} replaced by $\overline{\mathbb{Q}}$ was first proved in the unpublished note [23], and the special case $\mathbb{K} = \mathbb{Q}$ in [11].

PROPOSITION 3. — *Let f be an E -function with coefficients in a number field \mathbb{K} . Then there exist polynomials $P, Q \in \mathbb{K}[z]$, and an E -function g with coefficients in \mathbb{K} , such that*

$$f(z) = P(z) + Q(z)g(z) \text{ and } g(z_0) \text{ is transcendental for all } z_0 \in \overline{\mathbb{Q}}^*.$$

In this setting, the non-zero algebraic numbers z at which a transcendental f takes an algebraic value are exactly the roots of Q . Moreover, replacing P with its remainder in its Euclidean division by Q , we may assume $\deg P < \deg Q$ provided $Q \neq 0$ (i.e., when f is not a polynomial) and unicity then holds if Q is monic such that $Q(0) \neq 0$ (properties which can both be assumed without loss of generality); see [11, Prop. 3.3].

Proposition 3 is a generalization of the following result, which will be used in the proof. It is stated as [16, Th. 4] and its proof is due to the referee of [15].

LEMMA 1. — *Let f be an E -function with coefficients in a number field \mathbb{K} , and $\alpha \in \overline{\mathbb{Q}}$ be such that $f(\alpha)$ is algebraic. Then $f(\alpha) \in \mathbb{K}(\alpha)$.*

This lemma asserts that $\mathbf{E}_{\mathbb{K}(\alpha)} \cap \overline{\mathbb{Q}} = \mathbb{K}(\alpha)$; it is a consequence of Theorem 2.

For the convenience of the reader, let us deduce Lemma 1 from Proposition 1. Let $\beta = f(\alpha)$, and \mathbb{L} be a finite Galois extension of $\mathbb{K}(\alpha)$ such that $\beta \in \mathbb{L}$. Since $f(z) - \beta$ vanishes at α , Proposition 1 shows that for any $\sigma \in \text{Gal}(\mathbb{L}/\mathbb{K}(\alpha))$ the E -function $f(z) - \sigma(\beta) = f^\sigma(z) - \sigma(\beta)$ vanishes at $\sigma(\alpha) = \alpha$, so that $\sigma(\beta) = f(\alpha) = \beta$. This concludes the proof of Lemma 1.

Proof of Proposition 3. — To prove Proposition 3, we first remark that the result is obvious if f is algebraic, hence a polynomial: we simply take $P = f$ and $Q = 0$. Let us now assume that f is transcendental and argue by induction on the number of non-zero algebraic numbers α such that $f(\alpha) \in \overline{\mathbb{Q}}$. By Beukers' theorem this number is finite, indeed, since f is transcendental any such α must be one of the finitely many singularities of an appropriate differential equation of which f is a solution.

If this number is 0, one may choose $P = 0$ and $Q = 1$. Now, if $f(\alpha) \in \overline{\mathbb{Q}}$, Lemma 1 proves that $f(\alpha)$ belongs to $\mathbb{K}(\alpha)$: there exists $P_0 \in \mathbb{K}[X]$ such that $f(\alpha) = P_0(\alpha)$. Therefore the E -function $f - P_0$, with coefficients in \mathbb{K} , vanishes at α . Proposition 1 yields an E -function g_0 with coefficients in \mathbb{K} such that $f = P_0 + Dg_0$ where D is the minimal polynomial of α over \mathbb{K} . If $g_0(\alpha) \in \overline{\mathbb{Q}}$, the same procedure can be carried out with g_0 , leading to $P_1 \in \mathbb{K}[z]$ and an E -function g_1 with coefficients in \mathbb{K} such that $g_0 = P_1 + Dg_1$. After finitely many steps, this procedure terminates and provides g_ℓ such that $g_\ell(\alpha) \notin \overline{\mathbb{Q}}$ (see the proof of [11, Th. 3.4]). This concludes the proof of Proposition 3. \square

5. STRUCTURE OF $\mathbf{E}_{\mathbb{K}}$

In this section we prove Theorem 2 stated in the introduction. Let \mathbb{K} be a subfield of $\overline{\mathbb{Q}}$. Recall that $\mathbf{E}_{\mathbb{K}}$ is the ring of all values $f(1)$ where f is an E -function with coefficients in \mathbb{K} ; in particular $\mathbf{E}_{\overline{\mathbb{Q}}} = \mathbf{E}$.

Let f_1, \dots, f_N be E -functions with coefficients in \mathbb{K} . If $f_1(1), \dots, f_N(1)$ are linearly independent over $\overline{\mathbb{Q}}$, then obviously they are linearly independent over \mathbb{K} . Conversely, let us assume that they are linearly independent over \mathbb{K} . Let $\lambda_1, \dots, \lambda_N$ be algebraic numbers, not all zero, such that $\lambda_1 f_1(1) + \dots + \lambda_N f_N(1) = 0$. Up to a permutation of the indices we may assume that $\lambda_1 \neq 0$; then dividing by λ_1 we assume that $\lambda_1 = 1$. Let us consider a finite Galois extension \mathbb{L} of \mathbb{K} that contains $\lambda_2, \dots, \lambda_N$. Then $g(z) = \sum_{i=1}^N \lambda_i f_i(z)$ is an E -function with coefficients in \mathbb{L} , and it vanishes at $z = 1$. For any $\sigma \in \text{Gal}(\mathbb{L}/\mathbb{K})$, Proposition 1 yields $g^\sigma(1) = 0$, that is $\sum_{i=1}^N \sigma(\lambda_i) f_i(1) = 0$ since all f_i have coefficients in \mathbb{K} . Summing these relations, as σ varies, yields

$$\sum_{i=1}^N \text{Tr}_{\mathbb{L}/\mathbb{K}}(\lambda_i) f_i(1) = 0$$

with $\text{Tr}_{\mathbb{L}/\mathbb{K}}(\lambda_i) = \sum_{\sigma \in \text{Gal}(\mathbb{L}/\mathbb{K})} \sigma(\lambda_i) \in \mathbb{K}$ and $\text{Tr}_{\mathbb{L}/\mathbb{K}}(\lambda_1) = \text{Tr}_{\mathbb{L}/\mathbb{K}}(1) = [\mathbb{L} : \mathbb{K}] \neq 0$. This is a non-trivial linear relation, with coefficients in \mathbb{K} , between $f_1(1), \dots, f_N(1)$. This contradiction concludes the proof that elements of $\mathbf{E}_{\mathbb{K}}$ are linearly independent over $\overline{\mathbb{Q}}$ if, and only if, they are linearly independent over \mathbb{K} .

6. A GALOIS ACTION ON VALUES OF E -FUNCTIONS

In this section, we define an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on the set \mathbf{E} of values of E -functions. This action is not used in the paper but it sheds a different light on the proof of Theorem 1: it presents similarities with Liouville's proof that irrational algebraic numbers are not too well approximated by rationals (i.e., are not Liouville numbers).

Given $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ and $\xi \in \mathbf{E}$, there exists an E -function f such that $\xi = f(1)$; then we let $\sigma(\xi) := f^\sigma(1)$. The crucial point is to prove that $\sigma(\xi)$ depends only on σ and ξ , not on the choice of f . Indeed if g is another E -function such that $\xi = g(1)$, then $f - g$ vanishes at the point 1. Proposition 1 shows that $f^\sigma - g^\sigma$ vanishes at 1 too, so that $g^\sigma(1) = f^\sigma(1)$: this concludes the proof.

Theorem 2 shows that as a Galois representation, \mathbf{E} is isomorphic to $\mathbf{E}_{\mathbb{Q}} \otimes \overline{\mathbb{Q}}$, where $\mathbf{E}_{\mathbb{Q}}$ is a \mathbb{Q} -vector space with trivial Galois action. Therefore this does not provide a way to understand the absolute Galois group of \mathbb{Q} better. However, it sheds a new light on the proof of Theorem 1 (see §3): Step 2 is very similar to Liouville's proof that irrational algebraic numbers are not too well approximated by rationals (i.e., are not Liouville numbers). Indeed let us recall briefly Liouville's proof, stated in terms of Galois action. Let ξ be an algebraic number of degree $d \geq 2$, and assume (for simplicity) that the extension $\mathbb{Q}(\xi)/\mathbb{Q}$ is Galois. To bound from below $|q\xi - p|$ for $(p, q) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$, consider

$$\varpi := \prod_{\sigma \in \text{Gal}(\mathbb{Q}(\xi)/\mathbb{Q})} (q\sigma(\xi) - p).$$

Then $\varpi \neq 0$ (since $\sigma(\xi)$ is irrational for any σ), and $\varpi \in \mathbb{Q}$ (since it is the norm of $q\xi - p$ with respect to the extension $\mathbb{Q}(\xi)/\mathbb{Q}$). Letting $\delta \in \mathbb{Z}$ denote a positive integer such that $\delta\xi$ is an algebraic integer, we have $\delta^d\varpi \in \mathbb{Z} \setminus \{0\}$ since $\mathcal{O}_{\mathbb{Q}(\xi)} \cap \mathbb{Q} = \mathbb{Z}$. Therefore $|\delta^d\varpi| \geq 1$ so that

$$\delta^{-d} \leq |\varpi| \leq |q\xi - p|(\lceil \xi \rceil + 1)^{d-1} H^{d-1}$$

by bounding $|q\sigma(\xi) - p|$ trivially for $\sigma \neq \text{Id}$, where $H = \max(|p|, |q|)$. Dividing by q yields $|\xi - p/q| \geq cH^{-d}$ where $c > 0$ depends only on ξ .

Step 2 of the proof of Theorem 1 presents similarities, except that elements of $\mathbb{K} \subset \overline{\mathbb{Q}}$ are replaced with values of E -functions in $\mathbf{E}_{\mathbb{K}}$; the lower bound used by Liouville (namely, $\delta^d\varpi \in \mathbb{Z} \setminus \{0\}$ implies $|\delta^d\varpi| \geq 1$) is replaced accordingly by Shidlovskii's lower bound recalled in Theorem A.

REFERENCES

- [1] B. ADAMCZEWSKI & J. CASSAIGNE – “Diophantine properties of real numbers generated by finite automata”, *Compositio Math.* **142** (2006), no. 6, p. 1351–1372.
- [2] B. ADAMCZEWSKI & C. FAVERJON – “Méthode de Mahler: relations linéaires, transcendance et applications aux nombres automatiques”, *Proc. London Math. Soc. (3)* **115** (2017), no. 1, p. 55–90.
- [3] Y. ANDRÉ – *G-functions and geometry*, Aspects of Math., vol. E13, Friedr. Vieweg & Sohn, Braunschweig, 1989.
- [4] ———, “Séries Gevrey de type arithmétique. I. Théorèmes de pureté et de dualité”, *Ann. of Math. (2)* **151** (2000), no. 2, p. 705–740.
- [5] ———, “Solution algebras of differential equations and quasi-homogeneous varieties: a new differential Galois correspondence”, *Ann. Sci. École Norm. Sup. (4)* **47** (2014), no. 2, p. 449–467.
- [6] A. BAKER – *Transcendental number theory*, second ed., Cambridge Math. Library, Cambridge University Press, Cambridge, 1990.
- [7] J. P. BELL, Y. BUGEAUD & M. COONS – “Diophantine approximation of Mahler numbers”, *Proc. London Math. Soc. (3)* **110** (2015), no. 5, p. 1157–1206.
- [8] D. BERTRAND & F. BEUKERS – “Équations différentielles linéaires et majorations de multiplicités”, *Ann. Sci. École Norm. Sup. (4)* **18** (1985), no. 1, p. 181–192.
- [9] D. BERTRAND, V. CHIRSKII & J. YEBBOU – “Effective estimates for global relations on Euler-type series”, *Ann. Fac. Sci. Toulouse Math. (6)* **13** (2004), no. 2, p. 241–260.
- [10] F. BEUKERS – “A refined version of the Siegel-Shidlovskii theorem”, *Ann. of Math. (2)* **163** (2006), no. 1, p. 369–379.
- [11] A. BOSTAN, T. RIVOAL & B. SALVY – “Minimization of differential equations and algebraic values of E -functions”, 2022, [arXiv:2209.01827](https://arxiv.org/abs/2209.01827).
- [12] N. BOURBAKI – *Elements of mathematics. Algebra II. Chapters 4–7*, Springer, Berlin, 2003.
- [13] Y. BUGEAUD – *Approximation by algebraic numbers*, Cambridge Tracts in Math., vol. 160, Cambridge University Press, Cambridge, 2004.
- [14] N. I. FELDMAN & Y. V. NESTERENKO – “Transcendental numbers”, in *Number theory, IV*, Encyclopaedia Math. Sci., vol. 44, Springer, Berlin, 1998, p. 1–345.
- [15] S. FISCHLER & T. RIVOAL – “On the values of G -functions”, *Comment. Math. Helv.* **89** (2014), no. 2, p. 313–341.
- [16] ———, “Arithmetic theory of E -operators”, *J. Éc. polytech. Math.* **3** (2016), p. 31–65.
- [17] ———, “Microsolutions of differential operators and values of arithmetic Gevrey series”, *Amer. J. Math.* **140** (2018), no. 2, p. 317–348.
- [18] L.-C. KAPPE – “Zur Approximation von e^α ”, *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* **9** (1966), p. 3–14.
- [19] G. LEPETIT – “The André-Chudnovsky-Katz theorem in the broad sense”, *North-West. Eur. J. Math.* **7** (2021), p. 83–149.

- [20] K. MAHLER – “Zur Approximation der Exponentialfunktion und des Logarithmus. I”, *J. reine angew. Math.* **166** (1931), p. 118–136.
- [21] J. MILNE – “Fields and Galois theory”, 2022, version 5.10, available at www.jmilne.org/math, 144 pages.
- [22] YU. V. NESTERENKO & A. B. SHIDLOVSKIĬ – “On the linear independence of values of E -functions”, *Mat. Sb.* **187** (1996), no. 8, p. 93–108.
- [23] T. RIVOAL – “Valeurs algébriques de E -fonctions aux points algébriques”, HAL:hal-03676576.
- [24] A. B. SHIDLOVSKIĬ – *Transcendental numbers*, De Gruyter Studies in Math., vol. 12, Walter de Gruyter & Co., Berlin, 1989.
- [25] C. L. SIEGEL – “Über einige Anwendungen diophantischer Approximationen.”, *Abh. Preuß. Akad. Wiss., Phys.-Math. Kl.* **1929** (1929), no. 1.
- [26] V. V. ZUDILIN – “On rational approximations of values of a certain class of entire functions”, *Mat. Sb.* **186** (1995), no. 4, p. 555–590.

Manuscript received 25th April 2023

accepted 24th November 2023

STÉPHANE FISCHLER, Université Paris-Saclay, CNRS, Laboratoire de mathématiques d’Orsay,
91405 Orsay, France

E-mail : stephane.fischler@universite-paris-saclay.fr

Url : <https://www.imo.universite-paris-saclay.fr/~stephane.fischler/>

TANGUY RIVOAL, Université Grenoble Alpes, CNRS, Institut Fourier, CS 40700,
38058 Grenoble cedex 9, France

E-mail : tanguy.rivoal@univ-grenoble-alpes.fr

Url : <https://rivoal.perso.math.cnrs.fr/>