



ANNALES

DE

L'INSTITUT FOURIER

Nalini ANANTHARAMAN & Stéphane NONNENMACHER

Half-delocalization of eigenfunctions for the Laplacian on an Anosov manifold

Tome 57, n° 7 (2007), p. 2465-2523.

http://aif.cedram.org/item?id=AIF_2007__57_7_2465_0

© Association des Annales de l'institut Fourier, 2007, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

HALF-DELOCALIZATION OF EIGENFUNCTIONS FOR THE LAPLACIAN ON AN ANOSOV MANIFOLD

by Nalini ANANTHARAMAN & Stéphane
NONNENMACHER

ABSTRACT. — We study the high-energy eigenfunctions of the Laplacian on a compact Riemannian manifold with Anosov geodesic flow. The localization of a semiclassical measure associated with a sequence of eigenfunctions is characterized by the Kolmogorov-Sinai entropy of this measure. We show that this entropy is necessarily bounded from below by a constant which, in the case of constant negative curvature, equals half the maximal entropy. In this sense, high-energy eigenfunctions are at least half-delocalized.

RÉSUMÉ. — Nous étudions la limite de haute énergie pour les fonctions propres du laplacien, sur une variété riemannienne compacte dont le flot géodésique est d'Anosov. La localisation d'une mesure semiclassique associée à une suite de fonctions propres peut être mesurée par son entropie de Kolmogorov-Sinai. Nous obtenons pour cette entropie une borne inférieure qui, dans le cas des variétés à courbure négative constante, vaut la moitié de l'entropie maximale. En ce sens, on peut dire que les fonctions propres de haute énergie sont au moins à demi délocalisées.

The theory of quantum chaos tries to understand how the chaotic behaviour of a classical Hamiltonian system is reflected in its quantum version. For instance, let M be a compact Riemannian C^∞ manifold, such that the geodesic flow has the Anosov property — the ideal chaotic behaviour. The corresponding quantum dynamics is the unitary flow generated by the Laplace-Beltrami operator on $L^2(M)$. One expects that the chaotic properties of the geodesic flow influence the spectral theory of the Laplacian. The Random Matrix conjecture [6] asserts that the high-lying eigenvalues should, after proper renormalization, statistically resemble those of a large random matrix, at least for a generic Anosov metric. The Quantum Unique Ergodicity conjecture [27] (see also [5, 30]) deals with the corresponding eigenfunctions ψ : it claims that the probability density $|\psi(x)|^2 dx$ should approach (in a weak sense) the Riemannian volume, when the eigenvalue

Keywords: Quantum chaos, semiclassical measure, ergodic theory, entropy, Anosov flows.
Math. classification: 81Q50, 35Q40, 35P20, 37D40, 58J40, 28D20.

corresponding to ψ tends to infinity. In fact a stronger property should hold for the *Wigner transform* W_ψ , a distribution on the cotangent bundle T^*M which describes the distribution of the wave function ψ on the phase space T^*M . We will adopt a semiclassical point of view, that is consider the eigenstates of eigenvalue unity of the semiclassical Laplacian $-\hbar^2\Delta$, in the semiclassical limit $\hbar \rightarrow 0$. Weak limits of the distributions W_ψ are called *semiclassical measures*: they are invariant measures of the geodesic flow on the unit energy layer \mathcal{E} . The Quantum Unique Ergodicity conjecture asserts that on an Anosov manifold there exists a unique semiclassical measure, namely the Liouville measure on \mathcal{E} ; in other words, in the semiclassical régime all eigenfunctions become uniformly distributed over \mathcal{E} .

For manifolds with an ergodic geodesic flow (with respect to the Liouville measure), it has been shown by Schnirelman, Zelditch and Colin de Verdière that *almost all* eigenfunctions become uniformly distributed over \mathcal{E} , in the semiclassical limit: this property is dubbed as Quantum Ergodicity [29, 32, 8]. The possibility of *exceptional sequences* of eigenstates with different semiclassical limits remains open in general. The Quantum Unique Ergodicity conjecture states that such sequences do not exist for an Anosov manifold [27].

So far the most precise results on this question were obtained for Anosov manifolds M with *arithmetic* properties: see Rudnick-Sarnak [27], Wolpert [31]. Recently, Lindenstrauss [24] proved the asymptotic equidistribution of all “arithmetic” eigenstates (these are believed to exhaust the full family of eigenstates). The proof, unfortunately, cannot be extended to general Anosov manifolds.

To motivate the conjecture, one may instead invoke the following dynamical explanation. By the Heisenberg uncertainty principle, an eigenfunction cannot be strictly localized on a submanifold in phase space. Its microlocal support must contain a symplectic cube of volume \hbar^d , where d is the dimension of M . Since ψ is invariant under the quantum dynamics, which is semiclassically approximated by the geodesic flow, the *fast mixing* property of the latter will spread this cube throughout the energy layer, showing that the support of the eigenfunction must also spread throughout \mathcal{E} .

This argument is however too simplistic. First, Colin de Verdière and Parisse showed that, on a surface of revolution of negative curvature, eigenfunctions can concentrate on a single periodic orbit in the semiclassical limit, despite the exponential instability of that orbit [9]. Their construction shows that one cannot use purely local features, such as instability, to rule out localization of eigenfunctions on closed geodesics. Second, the

argument above is based on the classical dynamics, and does not take into account the *interferences* of the wavefunction with itself, after a long time. Faure, Nonnenmacher and De Bièvre exhibited in [15] a simple example of a symplectic Anosov dynamical system, namely the action of a linear hyperbolic automorphism on the 2-torus (also called “Arnold’s cat map”), the quantization of which does not satisfy the Quantum Unique Ergodicity conjecture. Precisely, they construct a family of eigenstates for which the semiclassical measure consists in two ergodic components: half of it is the Liouville measure, while the other half is a Dirac peak on a single unstable periodic orbit. It was also shown that — in the case of the “cat map” — this half-localization on a periodic orbit is *maximal* [14]. Another type of semiclassical measures was recently exhibited by Kelmer for quantized automorphisms on higher-dimensional tori and some of their perturbations [19, 20]: it consists in the Lebesgue measure on some invariant co-isotropic subspace of the torus. In those cases, the existence of exceptional eigenstates is due to some nongeneric algebraic properties of the classical and quantized systems.

In a previous paper [3], we discovered how to use an information-theoretic variant of the uncertainty principle [22, 25], called the *Entropic Uncertainty Principle*, to constrain the localization properties of eigenfunctions in the case of another toy model, the Walsh-quantized baker’s map. For any dynamical system, the complexity of an invariant measure can be described through its *Kolmogorov–Sinai entropy*. In the case of the Walsh-baker’s map, we showed that the entropy of semiclassical measures must be at least half the entropy of the Lebesgue measure. Thus, our result can be interpreted as a “half-delocalization” of eigenstates. The Walsh-baker model being very special, it was not clear whether the strategy could be generalized to more realistic systems, like geodesic flows or more general symplectic systems quantized *à la* Weyl.

In this paper we show that it is the case: the strategy used in [3] is rather general, and its implementation to the case of Anosov geodesic flows only requires more technical suffering.

1. Main result

Let M be a compact Riemannian manifold. We will denote by $|\cdot|_x$ the norm on T_x^*M given by the metric. The geodesic flow $(g^t)_{t \in \mathbb{R}}$ is the Hamiltonian flow on T^*M generated by the Hamiltonian

$$H(x, \xi) = \frac{|\xi|_x^2}{2}.$$

In the semiclassical setting, the corresponding quantum operator is $-\frac{\hbar^2 \Delta}{2}$, which generates the unitary flow $(U^t) = (\exp(it\hbar \frac{\Delta}{2}))$ acting on $L^2(M)$.

We denote by $(\psi_k)_{k \in \mathbb{N}}$ an orthonormal basis of $L^2(M)$ made of eigenfunctions of the Laplacian, and by $(\frac{1}{\hbar_k^2})_{k \in \mathbb{N}}$ the corresponding eigenvalues:

$$-\hbar_k^2 \Delta \psi_k = \psi_k, \text{ with } \hbar_{k+1} \leq \hbar_k.$$

We are interested in the high-energy eigenfunctions of $-\Delta$, in other words the semiclassical limit $\hbar_k \rightarrow 0$.

The Wigner distribution associated to an eigenfunction ψ_k is defined by

$$W_k(a) = \langle \text{Op}_{\hbar_k}(a) \psi_k, \psi_k \rangle_{L^2(M)}, \quad a \in C_c^\infty(T^*M).$$

Here Op_{\hbar_k} is a quantization procedure, set at the scale \hbar_k , which associates a bounded operator on $L^2(M)$ to any smooth phase space function a with nice behaviour at infinity (see for instance [10]). If a is a function on the manifold M , we have $W_k(a) = \int_M a(x) |\psi_k(x)|^2 dx$: the distribution W_k is a *microlocal lift* of the probability measure $|\psi_k(x)|^2 dx$ into a phase space distribution. Although the definition of W_k depends on a certain number of choices, like the choice of local coordinates, or of the quantization procedure (Weyl, anti-Wick, “right” or “left” quantization...), its asymptotic behaviour when $\hbar_k \rightarrow 0$ does not. Accordingly, we call *semiclassical measures* the limit points of the sequence $(W_k)_{k \in \mathbb{N}}$, in the distribution topology.

Using standard semiclassical arguments, one easily shows the following [8]:

PROPOSITION 1.1. — *Any semiclassical measure is a probability measure carried on the energy layer $\mathcal{E} = H^{-1}(\frac{1}{2})$ (which coincides with the unit cotangent bundle $\mathcal{E} = S^*M$). This measure is invariant under the geodesic flow.*

If the geodesic flow has the Anosov property — for instance if M has negative sectional curvature — then there exist many invariant probability measures on \mathcal{E} , in addition to the Liouville measure. The geodesic flow has countably many periodic orbits, each of them carrying an invariant probability measure. There are still many others, like the equilibrium states obtained by variational principles [18]. The Kolmogorov–Sinai entropy, also called metric entropy, of a (g^t) -invariant probability measure μ is a non-negative number $h_{KS}(\mu)$ that describes, in some sense, the complexity of a μ -typical orbit of the flow. For instance, a measure carried on a closed geodesic has zero entropy. An upper bound on the entropy is given by the Ruelle inequality: since the geodesic flow has the Anosov property, the energy layer \mathcal{E} is foliated into unstable manifolds of the flow, and for any

invariant probability measure μ one has

$$(1.1) \quad h_{KS}(\mu) \leq \left| \int_{\mathcal{E}} \log J^u(\rho) d\mu(\rho) \right|.$$

In this inequality, $J^u(\rho)$ is the *unstable Jacobian* of the flow at the point $\rho \in \mathcal{E}$, defined as the Jacobian of the map g^{-1} restricted to the unstable manifold at the point $g^1\rho$. Although $\log J^u(\rho)$ depends on the metric structure on \mathcal{E} , its *average* over any invariant measure does not, and this average is always negative. If M has dimension d and has constant sectional curvature -1 , one has $\log J^u \equiv -(d - 1)$, so this inequality just reads $h_{KS}(\mu) \leq d - 1$. The equality holds in (1.1) if and only if μ is the Liouville measure on \mathcal{E} [23]. Our central result is the following

THEOREM 1.2. — *Let M be a compact smooth Riemannian manifold with Anosov geodesic flow. Let μ be a semiclassical measure associated to the eigenfunctions of the Laplacian on M . Then the metric entropy of μ with respect to the geodesic flow satisfies*

$$(1.2) \quad h_{KS}(\mu) \geq \frac{3}{2} \left| \int_{\mathcal{E}} \log J^u(\rho) d\mu(\rho) \right| - (d - 1)\lambda_{\max},$$

where $d = \dim M$ and $\lambda_{\max} = \lim_{t \rightarrow \pm\infty} \frac{1}{t} \log \sup_{\rho \in \mathcal{E}} |dg_{\rho}^t|$ is the maximal expansion rate of the geodesic flow on \mathcal{E} .

In particular, if M has constant sectional curvature -1 (and thus $\lambda_{\max} = 1$), this means that

$$(1.3) \quad h_{KS}(\mu) \geq \frac{d - 1}{2}.$$

The first author proved in [2] that the entropy of such a semiclassical measure is bounded from below by a positive (hardly explicit) constant. The bound (1.3) in the above theorem is much sharper in the case of constant curvature. On the other hand, if the curvature varies a lot (still being negative everywhere), the right hand side of (1.2) may actually be negative, in which case the above bound is trivial. In fact, if the sectional curvatures vary in the interval $[-K_2^2, -K_1^2]$ then λ_{\max} can be as large as K_2 , whereas $\left| \int_{\mathcal{E}} \log J^u(\rho) d\mu(\rho) \right|$ may vary between $(d-1)K_1$ and $(d-1)K_2$, depending on the measure μ . This “problem” is not very surprising, since the above bound is generally not optimal. Indeed, in a subsequent work in collaboration with Herbert Koch [1], we have managed to slightly improve the above bound to

$$(1.4) \quad h_{KS}(\mu) \geq \left| \int_{\mathcal{E}} \log J^u(\rho) d\mu(\rho) \right| - \frac{(d - 1)\lambda_{\max}}{2}.$$

The numbers $3/2$ and $(d - 1)$ appearing in (1.2), which follow from the estimate in Theorem 2.7, are thus not fundamental, but result from some choices we have made. Indeed, the improved bound in (1.4) results from a modified, more symmetrical version of that estimate. This lower bound can still be negative in case of a strongly varying curvature. We believe that, by further improving the method presented below, the following bound could be obtained:

$$(1.5) \quad h_{KS}(\mu) \geq \frac{1}{2} \left| \int_{\mathcal{E}} \log J^u(\rho) d\mu(\rho) \right|.$$

This conjectured lower bound is now strictly positive for any invariant measure, and improves the bounds (1.2, 1.4) proved so far.

Remark 1.3. — Proposition 1.1 and Theorem 1.2 still apply if μ is not associated to a subsequence of eigenstates, but rather a sequence $(u_{\hbar})_{\hbar \rightarrow 0}$ of *quasimodes* of the Laplacian, of the following order:

$$\|(-\hbar^2 \Delta - 1)u_{\hbar}\| = o(\hbar |\log \hbar|^{-1}) \|u_{\hbar}\|, \quad \hbar \rightarrow 0.$$

This extension of the theorem requires little modifications, which we leave to the reader. It is also possible to prove lower bounds on the entropy in the case of quasimodes of the type

$$\|(-\hbar^2 \Delta - 1)u_{\hbar}\| \leq c\hbar |\log \hbar|^{-1} \|u_{\hbar}\|, \quad \hbar \rightarrow 0,$$

as long as $c > 0$ is sufficiently small. However, this extension is not as straightforward as in [2], so we defer it to a future work.

Remark 1.4. — In this article we only treat the case of Anosov geodesic flows. The same method could hopefully be extended to the case of manifolds with nonpositive sectional curvature, or even assuming only that there are no conjugate points (and maybe some growth condition on the volume of spheres). Such manifolds include the surfaces considered by H. Donnelly [11], which contain a flat cylinder supporting “bouncing ball quasimodes”. One future objective is to prove (1.5) in this more general context (with an adequate definition of $\log J^u$). Notice that any semiclassical measure associated with bouncing ball quasimodes has vanishing entropy, and is supported on a set of geodesics which are not unstable, so that $\log J^u$ vanishes on this support. The bound (1.5) therefore still makes sense in that case, but is trivial. On the other hand, proving (1.5) for such manifolds would have the nontrivial consequence to forbid semiclassical measures from being supported on an unstable closed orbit, like for Anosov manifolds.

In a more straightforward way, the present results can easily be adapted to the case of a general Hamiltonian flow, assumed to be Anosov on some compact energy layer. The quantum operator can then be any self-adjoint \hbar -quantization of the Hamilton function. The tools needed to prove estimates of the type (2.18) in a more general setting have been used in [26].

Although this paper is overall in the same spirit as [2], certain aspects of the proof are quite different. We recall that the proof given in [2] required to study the quantum dynamics far beyond the Ehrenfest time — *i.e.* the time needed by the classical flow to transform wavelengths ~ 1 into wavelengths $\sim \hbar$. In this paper we will study the dynamics until *twice* the Ehrenfest time, but not beyond. In variable curvature, the fact that the Ehrenfest time depends on the initial position seems to be the reason why the bounds (1.2, 1.4) are not optimal.

Quantum Unique Ergodicity would mean that

$$h_{KS}(\mu) = \left| \int_{\mathcal{E}} \log J^u(\rho) d\mu(\rho) \right|.$$

We believe however that (1.5) is the optimal result that can be obtained without using more precise information, like for instance upper bounds on the multiplicities of eigenvalues. Indeed, in the above mentioned examples of Anosov systems where Quantum Unique Ergodicity fails, the bound (1.5) is actually *sharp* [15, 19, 3]. In those examples, the spectrum has high degeneracies in the semiclassical limit, which allows for a lot of freedom to select the eigenstates. Such high degeneracies are not expected to happen in the case of the Laplacian on a negatively curved manifold. Yet, for the moment we have no clear understanding of the relationship between spectral degeneracies and failure of Quantum Unique Ergodicity.

Acknowledgements. — Both authors were partially supported by the Agence Nationale de la Recherche, under the grant ANR-05-JCJC-0107-01. They are grateful to Yves Colin de Verdière and Steve Zelditch for their comments and encouragement. They thank Jens Marklof for organizing their stay at Bristol University in June 2006, during which part of this article was written. S. Nonnenmacher also thanks Maciej Zworski and Didier Robert for interesting discussions, and Herbert Koch for his enlightening remarks on the Riesz-Thorin theorem.

2. Outline of the proof

2.1. Weighted entropic uncertainty principle

Our main tool is an adaptation of the entropic uncertainty principle conjectured by Kraus in [22] and proven by Maassen and Uffink [25]. This principle states that if a unitary matrix has “small” entries, then any of its eigenvectors must have a “large” Shannon entropy. For our purposes, we need an elaborate version of this uncertainty principle, which we shall prove in Section 6.

Let $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ be a complex Hilbert space, and denote $\|\psi\| = \sqrt{\langle \psi, \psi \rangle}$ the associated norm. Let $\pi = (\pi_k)_{k=1, \dots, \mathcal{N}}$ be a quantum partition of unity, that is, a family of operators on \mathcal{H} such that

$$(2.1) \quad \sum_{k=1}^{\mathcal{N}} \pi_k \pi_k^* = \text{Id}.$$

In other words, for all $\psi \in \mathcal{H}$ we have

$$\|\psi\|^2 = \sum_{k=1}^{\mathcal{N}} \|\psi_k\|^2, \text{ where we denote } \psi_k = \pi_k^* \psi \text{ for all } k = 1, \dots, \mathcal{N}.$$

If $\|\psi\| = 1$, we define the entropy of ψ with respect to the partition π as

$$h_\pi(\psi) = - \sum_{k=1}^{\mathcal{N}} \|\psi_k\|^2 \log \|\psi_k\|^2.$$

We extend this definition by introducing the notion of pressure, associated to a family $(\alpha_k)_{k=1, \dots, \mathcal{N}}$ of positive real numbers: it is defined by

$$p_{\pi, \alpha}(\psi) = - \sum_{k=1}^{\mathcal{N}} \|\psi_k\|^2 \log \|\psi_k\|^2 - \sum_{k=1}^{\mathcal{N}} \|\psi_k\|^2 \log \alpha_k^2.$$

In Theorem 2.1 below, we use two families of weights $(\alpha_k)_{k=1, \dots, \mathcal{N}}$, $(\beta_j)_{j=1, \dots, \mathcal{N}}$, and consider the corresponding pressures $p_{\pi, \alpha}$, $p_{\pi, \beta}$.

Besides the appearance of the weights α, β , we also modify the statement in [25] by introducing an auxiliary operator O — for reasons that should become clear later.

THEOREM 2.1. — *Let O be a bounded operator and \mathcal{U} an isometry on \mathcal{H} . Define $A = \max_k \alpha_k$, $B = \max_j \beta_j$ and*

$$c_O^{(\alpha, \beta)}(\mathcal{U}) \stackrel{\text{def}}{=} \sup_{j, k} \alpha_k \beta_j \|\pi_j^* \mathcal{U} \pi_k O\|_{\mathcal{L}(\mathcal{H})}.$$

Then, for any $\vartheta \geq 0$, for any normalized $\psi \in \mathcal{H}$ satisfying

$$\forall k = 1, \dots, \mathcal{N}, \quad \|(\text{Id} - O)\pi_k^* \psi\| \leq \vartheta,$$

the pressures $p_{\pi, \beta}(\mathcal{U}\psi)$, $p_{\pi, \alpha}(\psi)$ satisfy

$$p_{\pi, \beta}(\mathcal{U}\psi) + p_{\pi, \alpha}(\psi) \geq -2 \log (c_O^{(\alpha, \beta)}(\mathcal{U}) + \mathcal{N}AB\vartheta).$$

Remark 2.2. — The result of [25] corresponds to the case where \mathcal{H} is an \mathcal{N} -dimensional Hilbert space, $O = \text{Id}$, $\vartheta = 0$, $\alpha_k = \beta_j = 1$, and the operators π_k are orthogonal projectors on an orthonormal basis of \mathcal{H} . In this case, the theorem reads

$$h_\pi(\mathcal{U}\psi) + h_\pi(\psi) \geq -2 \log c(\mathcal{U}),$$

where $c(\mathcal{U})$ is the supremum of all matrix elements of \mathcal{U} in the orthonormal basis defined by π .

2.2. Applying the entropic uncertainty principle to the Laplacian eigenstates

In the whole article, we consider a certain subsequence of eigenstates $(\psi_{k_j})_{j \in \mathbb{N}}$ of the Laplacian, such that the corresponding sequence of Wigner functions (W_{k_j}) converges to a certain semiclassical measure μ (see the discussion preceding Proposition 1.1). The subsequence (ψ_{k_j}) will simply be denoted by $(\psi_{\hbar})_{\hbar \rightarrow 0}$, using the slightly abusive notation $\psi_{\hbar} = \psi_{\hbar k_j}$ for the eigenstate ψ_{k_j} . Each state ψ_{\hbar} satisfies

$$(2.2) \quad (-\hbar^2 \Delta - 1)\psi_{\hbar} = 0,$$

and we assume that

$$(2.3) \quad \text{the Wigner measures } W_{\psi_{\hbar}} \xrightarrow{\hbar \rightarrow 0} \mu \text{ in the weak-* topology.}$$

In this section we define the data to input in Theorem 2.1, in order to obtain informations on the eigenstates ψ_{\hbar} and the measure μ . Only the Hilbert space is fixed, $\mathcal{H} \stackrel{\text{def}}{=} L^2(M)$. All other data depend on the semiclassical parameter \hbar : the quantum partition π , the operator O , the positive real number ϑ , the weights (α_j) , (β_k) and the unitary operator \mathcal{U} .

2.2.1. Smooth partition of unity

As usual when computing the Kolmogorov–Sinai entropy, we start by decomposing the manifold M into small cells of diameter $\varepsilon > 0$. More precisely, let $(\Omega_k)_{k=1,\dots,K}$ be an open cover of M such that all Ω_k have diameters $\leq \varepsilon$, and let $(P_k)_{k=1,\dots,K}$ be a family of smooth real functions on M , with $\text{supp } P_k \Subset \Omega_k$, such that

$$(2.4) \quad \forall x \in M, \quad \sum_{k=1}^K P_k^2(x) = 1.$$

Most of the time, the notation P_k will actually denote the operator of multiplication by $P_k(x)$ on the Hilbert space $L^2(M)$: the above equation shows that they form a quantum partition of unity (2.1), which we will call $\mathcal{P}^{(0)}$.

2.2.2. Refinement of the partition under the Schrödinger flow

We denote the quantum propagator by $U^t = \exp(it\hbar \Delta / 2)$. With no loss of generality, we will assume that the injectivity radius of M is greater than 2, and work with the propagator at time unity, $U = U^1$. This propagator quantizes the flow at time one, g^1 . The \hbar -dependence of U will be implicit in our notations.

As one does to compute the Kolmogorov–Sinai entropy of an invariant measure, we define a new quantum partition of unity by evolving and refining the initial partition $\mathcal{P}^{(0)}$ under the quantum evolution. For each time $n \in \mathbb{N}$ and any sequence of symbols $\epsilon = (\epsilon_0 \cdots \epsilon_n)$, $\epsilon_i \in [1, K]$ (we say that the sequence ϵ is of length $|\epsilon| = n$), we define the operators

$$(2.5) \quad \begin{aligned} P_\epsilon &= P_{\epsilon_n} U P_{\epsilon_{n-1}} \cdots U P_{\epsilon_0} \\ \tilde{P}_\epsilon &= U^{-n} P_\epsilon = P_{\epsilon_n}(n) P_{\epsilon_{n-1}}(n-1) \cdots P_{\epsilon_0}. \end{aligned}$$

Throughout the paper we will use the notation $A(t) = U^{-t} A U^t$ for the quantum evolution of an operator A . From (2.4) and the unitarity of U , the family of operators $\{P_\epsilon\}_{|\epsilon|=n}$ obviously satisfies the resolution of identity $\sum_{|\epsilon|=n} P_\epsilon P_\epsilon^* = \text{Id}_{L^2}$, and therefore forms a quantum partition which we call $\mathcal{P}^{(n)}$. The operators \tilde{P}_ϵ also have this property, they will be used in the proof of the subadditivity, see sections 2.2.7 and 4.

2.2.3. Energy localization

In the semiclassical setting, the eigenstate ψ_{\hbar} of (2.2) is associated with the energy layer $\mathcal{E} = \mathcal{E}(1/2) = \{\rho \in T^*M, H(\rho) = 1/2\}$. Starting from the cotangent bundle T^*M , we restrict ourselves to a compact phase space by introducing an energy cutoff (actually, several cutoffs) near \mathcal{E} . To optimize our estimates, we will need this cutoff to depend on \hbar in a sharp way. For some fixed $\delta \in (0, 1)$, we consider a smooth function $\chi_{\delta} \in C^{\infty}(\mathbb{R}; [0, 1])$, with $\chi_{\delta}(t) = 1$ for $|t| \leq e^{-\delta/2}$ and $\chi_{\delta}(t) = 0$ for $|t| \geq 1$. Then, we rescale that function to obtain a family of \hbar -dependent cutoffs near \mathcal{E} :

$$(2.6) \quad \forall \hbar \in (0, 1), \forall n \in \mathbb{N}, \forall \rho \in T^*M,$$

$$\chi^{(n)}(\rho; \hbar) \stackrel{\text{def}}{=} \chi_{\delta}(e^{-n\delta} \hbar^{-1+\delta}(H(\rho) - 1/2)).$$

The cutoff $\chi^{(0)}$ is localized in an energy interval of length $2\hbar^{1-\delta}$. Choosing $0 < C_{\delta} < \delta^{-1} - 1$, we will only consider indices $n \leq C_{\delta} |\log \hbar|$, such that the “widest” cutoff will be supported in an interval of *microscopic* length $2\hbar^{1-(1+C_{\delta})\delta} \ll 1$. In our applications, we will always take δ small enough, so that we can take C_{δ} of the form

$$(2.7) \quad 4/\lambda_{\max} < C_{\delta} < \delta^{-1} - 1.$$

These cutoffs can be quantized into pseudodifferential operators $\text{Op}(\chi^{(n)}) = \text{Op}_{\mathcal{E}, \hbar}(\chi^{(n)})$ described in Section 5.1 (the quantization uses a nonstandard pseudodifferential calculus drawn from [28]). It is shown there (see Proposition 5.4) that the eigenstate ψ_{\hbar} satisfies

$$(2.8) \quad \|(\text{Op}(\chi^{(0)}) - 1)\psi_{\hbar}\| = \mathcal{O}(\hbar^{\infty})\|\psi_{\hbar}\|.$$

Here and below, the norm $\|\cdot\|$ will either denote the Hilbert norm on $\mathcal{H} = L^2(M)$, or the corresponding operator norm.

Remark 2.3. — Although the use of these sharp cutoffs is quite tedious (due to the nonstandard pseudodifferential calculus they require), their use seems necessary to obtain the lower bound (1.2) (and similarly for (1.4)). Using cutoffs localizing in an energy strip of width $\hbar^{1/2-\delta}$ would allow us to use more standard symbol classes (of the type (5.4)), but it would lead to the lower bound $3/2|\int_{\mathcal{E}} \log J^u d\mu| - d\lambda_{\max}$ for the entropy, which is worse than (1.2) (see the remark following Theorem 2.7). In constant curvature, in particular, we would get the lower bound $\frac{d-3}{2}$ which is trivial for $d \leq 3$.

Remark 2.4. — We will constantly use the fact that sharp energy localization is almost preserved by the operators P_{ϵ} . Indeed, using results of section 5.4, namely the first statement of Corollary 5.6 and the norm estimate

(5.12), we obtain that for \hbar small enough and any $m, m' \leq C_\delta |\log \hbar|/2$,

$$(2.9) \quad \forall |\epsilon| = m, \|\text{Op}(\chi^{(m'+m)})P_\epsilon^* \text{Op}(\chi^{(m')}) - P_\epsilon^* \text{Op}(\chi^{(m')})\| = \mathcal{O}(\hbar^\infty).$$

Here the implied constants are uniform with respect to m, m' — and of course the same estimates hold if we replace P_ϵ^* by P_ϵ . Similarly, from §5 one can easily show that

$$\forall |\epsilon| = m, \|P_\epsilon \text{Op}(\chi^{(m')}) - P_\epsilon^f \text{Op}(\chi^{(m')})\| = \mathcal{O}(\hbar^\infty),$$

where $P_{\epsilon_j}^f \stackrel{\text{def}}{=} \text{Op}_\hbar(P_{\epsilon_j} f)$, f is a smooth, compactly supported function in T^*M which takes the value 1 in a neighbourhood of \mathcal{E} — and $P_\epsilon^f = P_{\epsilon_m}^f U P_{\epsilon_{m-1}}^f \cdots U P_{\epsilon_0}^f$.

In the whole paper, we will fix a small $\delta' > 0$, and call “Ehrenfest time” the \hbar -dependent integer

$$(2.10) \quad n_E(\hbar) \stackrel{\text{def}}{=} \lfloor \frac{(1 - \delta') |\log \hbar|}{\lambda_{\max}} \rfloor.$$

Unless indicated otherwise, the integer n will always be taken equal to n_E . For us, the significance of the Ehrenfest time is that it is the largest time interval on which the (non-commutative) dynamical system formed by (U^t) acting on pseudodifferential operators can be treated as being, approximately, commutative (see (4.2)).

Using the estimates (2.9) with $m = n, m' = 0$ together with (2.8), one easily checks the following

PROPOSITION 2.5. — *Fix $\delta > 0, C_\delta > 0$ satisfying (2.7). For any fixed $L > 0$, there exists $\hbar_{\delta,L} > 0$ such that, for any $\hbar \leq \hbar_{\delta,L}$, any $n \leq C_\delta |\log \hbar|$, the Laplacian eigenstate ψ_\hbar satisfies*

$$(2.11) \quad \forall \epsilon, |\epsilon| = n, \|(\text{Op}(\chi^{(n)}) - \text{Id})P_\epsilon^* \psi_\hbar\| \leq \hbar^L \|\psi_\hbar\|.$$

Notice that this estimate includes the sequences ϵ of length $n = n_E(\hbar)$.

2.2.4. Applying the entropic uncertainty principle

We now precise some of the data we will use in the entropic uncertainty principle, Theorem 2.1:

- the quantum partition π is given by the family of operators $\{P_\epsilon, |\epsilon| = n = n_E\}$. In the semiclassical limit, this partition has cardinality $\mathcal{N} = K^n \asymp \hbar^{-K_0}$ for some fixed $K_0 > 0$.
- the operator O is $O = \text{Op}(\chi^{(n)})$, and by Proposition 2.5, we can take $\vartheta = \hbar^L$, where L will be chosen very large (see §2.2.6).

- the isometry will be $\mathcal{U} = U^n = U^{n_E}$.
- the weights $\alpha_\epsilon, \beta_\epsilon$ will be selected in §2.2.6. They will be semiclassically tempered, meaning that there exists $K_1 > 0$ such that, for \hbar small enough, all $\alpha_\epsilon, \beta_\epsilon$ are contained in the interval $[1, \hbar^{-K_1}]$.

As in Theorem 2.1, the entropy and pressures associated with a normalized state $\phi \in \mathcal{H}$ are given by

$$(2.12) \quad h_n(\phi) = h_{\mathcal{P}(n)}(\phi) = - \sum_{|\epsilon|=n} \|P_\epsilon^* \phi\|^2 \log (\|P_\epsilon^* \phi\|^2),$$

$$(2.13) \quad p_{n,\alpha}(\phi) = h_n(\phi) - 2 \sum_{|\epsilon|=n} \|P_\epsilon^* \phi\|^2 \log \alpha_\epsilon.$$

We may apply Theorem 2.1 to any sequence of states satisfying (2.11), in particular the eigenstates ψ_\hbar .

COROLLARY 2.6. — Consider the assumptions of Prop. 2.5. Define

$$(2.14) \quad c_{\text{Op}(\chi^{(n)})}^{\alpha,\beta}(U^n) \stackrel{\text{def}}{=} \max_{|\epsilon|=|\epsilon'|=n} (\alpha_\epsilon \beta_{\epsilon'} \|P_{\epsilon'}^* U^n P_\epsilon \text{Op}(\chi^{(n)})\|).$$

Then for any $\hbar \leq \hbar_{\delta,L}$, and for any normalized state ϕ satisfying the property (2.11) with $n = n_E(\hbar)$, we have

$$p_{n,\beta}(U^n \phi) + p_{n,\alpha}(\phi) \geq -2 \log (c_{\text{Op}(\chi^{(n)})}^{\alpha,\beta}(U^n) + \hbar^{L-K_0-2K_1}).$$

Most of Section 3 will be devoted to obtaining a good upper bound for the norms $\|P_{\epsilon'}^* U^n P_\epsilon \text{Op}(\chi^{(n)})\|$ involved in the above quantity. The bound is given in Theorem 2.7 below. Our choice for the weights $\alpha_\epsilon, \beta_\epsilon$ will then be guided by these upper bounds.

2.2.5. Unstable Jacobian for the geodesic flow

We need to recall a few definitions pertaining to Anosov flows. For any $\lambda > 0$, the geodesic flow g^t is Anosov on the energy layer $\mathcal{E}(\lambda) = H^{-1}(\lambda) \subset T^*M$. This implies that for each $\rho \in \mathcal{E}(\lambda)$, the tangent space $T_\rho \mathcal{E}(\lambda)$ splits into

$$T_\rho \mathcal{E}(\lambda) = E^u(\rho) \oplus E^s(\rho) \oplus \mathbb{R}X_H(\rho)$$

where E^u is the unstable subspace and E^s the stable subspace. The unstable Jacobian $J^u(\rho)$ at the point ρ is defined as the Jacobian of the map g^{-1} , restricted to the unstable subspace at the point $g^1 \rho$: $J^u(\rho) = \det (dg_{|_{E^u(g^1 \rho)}}^{-1})$ (the unstable spaces at ρ and $g^1 \rho$ are equipped with the induced Riemannian metric). This Jacobian can be “coarse-grained” as

follows in a neighbourhood $\mathcal{E}^\varepsilon \stackrel{\text{def}}{=} \mathcal{E}([1/2 - \varepsilon, 1/2 + \varepsilon])$ of \mathcal{E} . For any pair $(\epsilon_0, \epsilon_1) \in [1, K]^2$, we define

$$(2.15) \quad J_1^u(\epsilon_0, \epsilon_1) \stackrel{\text{def}}{=} \sup \{ J^u(\rho) : \rho \in T^*\Omega_{\epsilon_0} \cap \mathcal{E}^\varepsilon, g^1 \rho \in T^*\Omega_{\epsilon_1} \}$$

if the set on the right hand side is not empty, and $J_1^u(\epsilon_0, \epsilon_1) = e^{-\Lambda}$ otherwise, where $\Lambda > 0$ is a fixed large number. For any sequence of symbols ϵ of length n , we define the coarse-grained Jacobian

$$(2.16) \quad J_n^u(\epsilon) \stackrel{\text{def}}{=} J_1^u(\epsilon_0, \epsilon_1) \cdots J_1^u(\epsilon_{n-1}, \epsilon_n).$$

Although J^u and $J_1^u(\epsilon_0, \epsilon_1)$ are not necessarily everywhere smaller than unity, there exists $C, \lambda_+, \lambda_- > 0$ such that, for any $n > 0$, all the coarse-grained Jacobians of length n satisfy

$$(2.17) \quad C^{-1} e^{-n(d-1)\lambda_+} \leq J_n^u(\epsilon) \leq C e^{-n(d-1)\lambda_-}.$$

One can take $\lambda_+ = \lambda_{\max}(1 + \varepsilon)$. We can now give our central estimate, proven in Section 3.

THEOREM 2.7. — *Given $\delta' \in (0, 1)$, $\delta > 0$, small enough to satisfy (2.7), define the Ehrenfest time $n_E(\hbar)$ by (2.10), and the family of cut-off functions $\chi^{(n)}$ as in (2.6).*

Given a partition $\mathcal{P}^{(0)}$, there exists $\hbar_{\mathcal{P}^{(0)}, \delta, \delta'}$ such that, for any $\hbar \leq \hbar_{\mathcal{P}^{(0)}, \delta, \delta'}$, for any positive integer $n \leq n_E(\hbar)$, and any pair of sequences ϵ, ϵ' of length n ,

$$(2.18) \quad \|P_{\epsilon'}^* U^n P_\epsilon \text{Op}(\chi^{(n)})\| \leq C \hbar^{-(d-1+c\delta)} J_n^u(\epsilon)^{1/2} J_n^u(\epsilon').$$

Here $d = \dim M$, and the constants c, C only depend on the Riemannian manifold (M, g) .

We notice that the numbers appearing in the lower bound (1.2) already appear in the above right hand side: the power of \hbar leads to the factor $-(d - 1)$, while adding the powers of the two Jacobians gives the factor $3/2$. The use of the sharp cutoffs $\chi^{(n)}$ is crucial to get (2.18): with cutoffs of width $\geq \hbar^{1/2}$, the first factor on the right hand side would have been $C \hbar^{-(d+c\delta)}$.

2.2.6. Choice of the weights

There remains to choose the weights $(\alpha_\epsilon, \beta_\epsilon)$ to use in Theorem 2.1. Our choice is guided by the following idea: in the quantity (2.14), the weights should balance the variations (with respect to ϵ, ϵ') in the norms, such as to

make all terms in (2.14) of the same order. Using the upper bounds (2.18), we end up with the following choice for all ϵ of length n :

$$(2.19) \quad \alpha_\epsilon \stackrel{\text{def}}{=} J_n^u(\epsilon)^{-1/2} \text{ and } \beta_\epsilon \stackrel{\text{def}}{=} J_n^u(\epsilon)^{-1}.$$

All these quantities are defined using the Ehrenfest time $n = n_E(\hbar)$. From (2.17), there exists $K_1 > 0$ such that, for \hbar small enough, all the weights are bounded by

$$(2.20) \quad 1 \leq |\alpha_\epsilon| \leq \hbar^{-K_1}, \quad 1 \leq |\beta_\epsilon| \leq \hbar^{-K_1},$$

as announced in §2.2.4. The estimate (2.18) can then be rewritten as

$$c_{\text{Op}(X^{(n)})}^{\alpha, \beta}(U^n) \leq C \hbar^{-(d-1+c\delta)}.$$

We now apply Corollary 2.6 to the particular case of the eigenstates ψ_\hbar . We choose L large enough such that $\hbar^{L-K_0-2K_1}$ is negligible in comparison with $\hbar^{-(d-1+c\delta)}$, and consider the parameter $\hbar_0 = \min(\hbar_{\delta, L}, \hbar_{\mathcal{P}^{(0)}, \delta, \delta'})$, where $\hbar_{\delta, L}, \hbar_{\mathcal{P}^{(0)}, \delta, \delta'}$ appear respectively in Proposition 2.5 and Theorem 2.7.

PROPOSITION 2.8. — *Let $(\psi_\hbar)_{\hbar \rightarrow 0}$ be our sequence of eigenstates (2.2). Then, for $\hbar < \hbar_0$, the pressures of ψ_\hbar at the Ehrenfest time $n = n_E(\hbar)$ (see (2.10)) relative to the weights (2.19) satisfy*

$$(2.21) \quad \begin{aligned} p_{n, \alpha}(\psi_\hbar) + p_{n, \beta}(\psi_\hbar) &\geq 2(d - 1 + c\delta) \log \hbar - \tilde{C} \\ &\geq -2 \frac{(d - 1 + c\delta) \lambda_{\max}}{(1 - \delta')} n - \tilde{C}. \end{aligned}$$

Here \tilde{C} only depends on the constant C appearing in (2.18).

2.2.7. Subadditivity until the Ehrenfest time

Before taking the limit $\hbar \rightarrow 0$, we prove that a similar lower bound holds if we replace $n \asymp |\log \hbar|$ by some fixed n_o , and $\mathcal{P}^{(n)}$ by the corresponding partition $\mathcal{P}^{(n_o)}$. This is due to the following *subadditivity property*, which is the semiclassical analogue of the classical subadditivity of pressures for invariant measures.

PROPOSITION 2.9 (Subadditivity). — *Let $\delta' > 0$ and define the Ehrenfest time $n_E(\hbar)$ as in (2.10). There exists a real number $R > 0$ independent of δ' and a function $R(\bullet, \bullet)$ on $\mathbb{N} \times (0, 1]$ such that*

$$\forall n_o \in \mathbb{N}, \limsup_{\hbar \rightarrow 0} |R(n_o, \hbar)| \leq R,$$

and with the following properties. For any $\hbar \in (0, 1]$, any $n_o, m \in \mathbb{N}$ with $n_o + m \leq n_E(\hbar)$, for ψ_\hbar any normalized eigenstate satisfying (2.2), the pressures associated with the weights α_ϵ of (2.19) satisfy

$$p_{n_o+m, \alpha}(\psi_\hbar) \leq p_{n_o, \alpha}(\psi_\hbar) + p_{m-1, \alpha}(\psi_\hbar) + R(n_o, \hbar).$$

The same inequality holds for the pressures $p_{n_o+m, \beta}(\psi_\hbar)$ associated with the weights β_ϵ .

The proof is given in §4. The time $n_o + m$ needs to be smaller than the Ehrenfest time because, in order to show the subadditivity, the various operators $P_{\epsilon_i}(i)$ composing \tilde{P}_ϵ have to approximately commute with each other. Indeed, for $m \geq n_E(\hbar)$ the commutator $[P_{\epsilon_m}(m), P_{\epsilon_0}]$ may have a norm of order unity.

Equipped with this subadditivity, we may finish the proof of Theorem 1.2. Let $n_o \in \mathbb{N}$ be fixed and $n = n_E(\hbar)$. Using the Euclidean division $n = q(n_o + 1) + r$, with $r \leq n_o$, Proposition 2.9 implies that for \hbar small enough,

$$\frac{p_{n, \alpha}(\psi_\hbar)}{n} \leq \frac{p_{n_o, \alpha}(\psi_\hbar)}{n_o} + \frac{p_{r, \alpha}(\psi_\hbar)}{n} + \frac{R(n_o, \hbar)}{n_o}.$$

Using (2.21) and the fact that $p_{r, \alpha}(\psi_\hbar) + p_{r, \beta}(\psi_\hbar)$ stays uniformly bounded (by a quantity depending on n_o) when $\hbar \rightarrow 0$, we find

$$(2.22) \quad \frac{p_{n_o, \alpha}(\psi_\hbar)}{n_o} + \frac{p_{n_o, \beta}(\psi_\hbar)}{n_o} \geq -2 \frac{(d-1+c\delta)\lambda_{\max}}{(1-\delta')} - 2 \frac{R(n_o, \hbar)}{n_o} + \mathcal{O}_{n_o}(1/n).$$

We are now dealing with the partition $\mathcal{P}^{(n_o)}$, n_o being independent of \hbar .

2.2.8. End of the proof

Because ψ_\hbar are eigenstates of U , the norms appearing in the definition of $h_{n_o}(\psi_\hbar)$ can be alternatively written as

$$(2.23) \quad \|P_\epsilon^* \psi_\hbar\| = \|\tilde{P}_\epsilon^* \psi_\hbar\| = \|P_{\epsilon_0} P_{\epsilon_1}(1) \cdots P_{\epsilon_{n_o}}(n_o) \psi_\hbar\|.$$

We may take the limit $\hbar \rightarrow 0$ (so that $n \rightarrow \infty$) in (2.22). The assumption (2.3) implies that, for any sequence ϵ of length n_o , $\|\tilde{P}_\epsilon^* \psi_\hbar\|^2$ converges to $\mu(\{\epsilon\})$, where $\{\epsilon\}$ is the function $P_{\epsilon_0}^2(P_{\epsilon_1}^2 \circ g^1) \cdots (P_{\epsilon_{n_o}}^2 \circ g^{n_o})$ on T^*M . Thus $h_{n_o}(\psi_\hbar)$ semiclassically converges to the classical entropy

$$h_{n_o}(\mu) = h_{n_o}(\mu, (P_k^2)) = - \sum_{|\epsilon|=n_o} \mu(\{\epsilon\}) \log \mu(\{\epsilon\}).$$

As a result, the left hand side of (2.22) converges to

$$(2.24) \quad \frac{2}{n_o} h_{n_o}(\mu) + \frac{3}{n_o} \sum_{|\epsilon|=n_o} \mu(\{\epsilon\}) \log J_{n_o}^u(\epsilon).$$

Since the semiclassical measure μ is g^t -invariant and $J_{n_o}^u$ has the multiplicative structure (2.16), the second term in (2.24) can be simplified:

$$\sum_{|\epsilon|=n_o} \mu(\{\epsilon\}) \log J_{n_o}^u(\epsilon) = n_o \sum_{\epsilon_0, \epsilon_1} \mu(\{\epsilon_0 \epsilon_1\}) \log J_1^u(\epsilon_0, \epsilon_1).$$

We have thus obtained the lower bound

$$(2.25) \quad \frac{h_{n_o}(\mu)}{n_o} \geq -\frac{3}{2} \sum_{\epsilon_0, \epsilon_1} \mu(\{\epsilon_0 \epsilon_1\}) \log J_1^u(\epsilon_0, \epsilon_1) - \frac{(d-1+c\delta)\lambda_{\max}}{(1-\delta')} - 2\frac{R}{n_o}.$$

δ and δ' could be taken arbitrarily small, and at this stage they can be let vanish.

The Kolmogorov–Sinai entropy of μ is by definition the limit of the first term $\frac{h_{n_o}(\mu)}{n_o}$ when n_o goes to infinity, with the notable difference that the smooth functions P_k should be replaced by characteristic functions associated with some partition of M , $M = \bigsqcup_k O_k$. Thus, let us consider such a partition of diameter $\leq \varepsilon/2$, such that μ does not charge the boundaries of the O_k . This last requirement can be easily enforced, if necessary by slightly shifting the O_k . One may for instance construct a “hypercubic partition” defined locally by a finite family of hypersurfaces $(S_i)_{i=1, \dots, I}$. If some of the S_i charge μ , one can (using local coordinates) translate them by arbitrarily small amounts \vec{v}_i such that $\mu(S_i + \vec{v}_i) = 0$. The boundary of the partition defined by those translated hypersurfaces does not charge μ (see e.g. [2, Appendix A2] for a similar construction).

By convolution we can smooth the characteristic functions $(\mathbb{1}_{O_k})$ into a smooth partition of unity (P_k) satisfying the conditions of section 2.2.1 (in particular, each P_k is supported on a set Ω_k of diameter $\leq \varepsilon$). The lower bound (2.25) holds with respect to the smooth partition (P_k^2) , and does not depend on the derivatives of the P_k : as a result, the same bound carries over to the characteristic functions $(\mathbb{1}_{O_k})$.

We can finally let n_o tend to $+\infty$, then let the diameter $\varepsilon/2$ of the partition tend to 0. From the definition (2.15) of the coarse-grained Jacobian, the first term in the right hand side of (2.25) converges to the integral $-\frac{3}{2} \int_{\mathcal{E}} \log J^u(\rho) d\mu(\rho)$ as $\varepsilon \rightarrow 0$. Since the integral of $\log J^u$ is negative, this proves (1.2). □

The next sections are devoted to proving, successively, Theorem 2.7, Proposition 2.9 and Theorem 2.1.

3. The main estimate: proof of Theorem 2.7

3.1. Strategy of the proof

We want to bound from above the norm of the operator $P_{\epsilon'}^* U^n P_{\epsilon} \text{Op}(\chi^{(n)})$. This norm can be obtained as follows:

$$\|P_{\epsilon'}^* U^n P_{\epsilon} \text{Op}(\chi^{(n)})\| = \sup \left\{ |\langle P_{\epsilon'} \Phi, U^n P_{\epsilon} \text{Op}(\chi^{(n)}) \Psi \rangle| : \Psi, \Phi \in \mathcal{H}, \|\Psi\| = \|\Phi\| = 1 \right\}.$$

Using Remark 2.4, we may insert $\text{Op}(\chi^{(4n)})$ on the right of $P_{\epsilon'}$, up to an error $\mathcal{O}_{L^2}(\hbar^\infty)$. In this section we will prove the following

PROPOSITION 3.1. — *For \hbar small enough, for any time $n \leq n_E(\hbar)$, for any sequences ϵ, ϵ' of length n and any normalized states $\Psi, \Phi \in L^2(M)$, one has*

$$(3.1) \quad |\langle P_{\epsilon'} \text{Op}(\chi^{(4n)}) \Phi, U^n P_{\epsilon} \text{Op}(\chi^{(n)}) \Psi \rangle| \leq C \hbar^{-(d-1)-c\delta} J_n^u(\epsilon)^{1/2} J_n^u(\epsilon').$$

Here we have taken δ small enough such that $C_\delta > 4/\lambda_{\max}$, and $n_E(\hbar)$ is the Ehrenfest time (2.10). The constants C and $c = 2 + 5/\lambda_{\max}$ only depend on the Riemannian manifold M .

For such times n , the right hand side in the above bound is larger than $C \hbar^{\frac{1}{2}(d-1)}$, in comparison to which the errors $\mathcal{O}(\hbar^\infty)$ are negligible. Theorem 2.7 therefore follows from the above proposition.

The idea in Proposition 3.1 is rather simple, although the technical implementation becomes cumbersome. We first show that any state of the form $\text{Op}(\chi^{(*)})\Psi$, as those appearing on both sides of the scalar product (3.1), can be decomposed as a superposition of essentially $\hbar^{-\frac{(d-1)}{2}}$ “elementary” Lagrangian states, supported on Lagrangian manifolds transverse to the stable leaves of the flow: see §3.2. In fact, our elementary Lagrangian states, defined in (3.2), are truncated δ -functions, microlocally supported on Lagrangians of the form $\cup_t g^t S_z^* M$, where $S_z^* M$ is the unit sphere at the point z . Any function of the form $\text{Op}(\chi^{(*)})\Psi$ is a superposition, in the z variable, of such states. The action of the Schrödinger flow U^t on a Lagrangian state is analyzed by the WKB method, described in §3.3.1. For each time t , U^t is a Fourier integral operator associated with g^t , the geodesic flow at time t . Then, the action of the operator $P_\epsilon = P_{\epsilon_n} U P_{\epsilon_{n-1}} U \cdots U P_{\epsilon_0}$ on a Lagrangian state is intuitively simple to understand: each application of U amounts to applying g^1 to the underlying Lagrangian manifold, which it stretches in the unstable direction (the rate of elongation being described

by the unstable Jacobian), while each multiplication by P_ϵ cuts out a small piece of the stretched Lagrangian. This iteration of stretching and cutting accounts for the exponential decay, see §3.4.2.

3.2. Decomposition of $\text{Op}(\chi)\Psi$ into elementary Lagrangian states

In Proposition 3.1, we apply the cutoff $\text{Op}(\chi^{(n)})$ on Ψ , respectively $\text{Op}(\chi^{(4n)})$ on Φ . To avoid too cumbersome notations, we treat both cases at the same time, denoting both cutoffs by $\chi = \chi^{(*)}$, and their associated quantization by $\text{Op}(\chi)$. The original notations will be restored only when needed. The energy cutoff χ is supported on a microscopic energy interval, where it varies between 0 and 1. In spite of those fast variations in the direction transverse to \mathcal{E} , it can be quantized such as to satisfy some sort of pseudodifferential calculus. As explained in Section 5.3, the quantization $\text{Op} \stackrel{\text{def}}{=} \text{Op}_{\mathcal{E}, \hbar}$ (see (5.10)) uses a *finite* family of Fourier Integral Operators (U_{κ_j}) associated with local canonical maps (κ_j). Each κ_j sends an open bounded set $\mathcal{V}_j \subset T^*M$ intersecting \mathcal{E} to $\mathcal{W}_j \subset \mathbb{R}^{2d}$, endowed with coordinated $(y, \eta) = (y_1, \dots, y_d, \eta_1, \dots, \eta_d)$, such that $H \circ \kappa_j^{-1} = \eta_1 + 1/2$. In other words, each κ_j defines a set of local flow-box coordinates (y, η) , such that y_1 is the time variable and $\eta_1 + 1/2$ the energy, while $(y', \eta') \in \mathbb{R}^{2(d-1)}$ are symplectic coordinates in a Poincaré section transverse to the flow.

3.2.1. Integral representation of U_{κ_j}

Since κ_j is defined only on \mathcal{V}_j , one may assume that $U_{\kappa_j}u = 0$ for functions $u \in L^2(M \setminus \pi\mathcal{V}_j)$ (here and below π will represent either the projection from T^*M to M along fibers, or from $\mathbb{R}^{2d}_{y, \eta}$ to \mathbb{R}^d_y). If \mathcal{V}_j is small enough, the action of U_{κ_j} on a function $\Psi \in L^2(M)$ can be represented as follows:

$$[U_{\kappa_j}\Psi](y) = (2\pi\hbar)^{-\frac{D+d}{2}} \int_{\pi\mathcal{V}_j} e^{\frac{i}{\hbar}S(z,y,\theta)} a_{\hbar}(z, y, \theta) \Psi(z) dz d\theta,$$

where

- θ takes values in an open set $\Theta_j \subset \mathbb{R}^D$ for some integer $D \geq 0$,
- the Lagrangian manifold generated by S is the graph of κ_j ,
- $a_{\hbar}(z, y, \theta)$ has an asymptotic expansion $a_{\hbar} \sim \sum_{l \geq 0} \hbar^l a_l$, and it is supported on $\pi\mathcal{V}_j \times \pi\mathcal{W}_j \times \Theta_j$.

When applying the definition (5.10) to the cutoff χ , we notice that the product $\chi(1 - \phi) \equiv 0$, so that $\text{Op}(\chi)$ is given by the sum of operators

$\text{Op}(\chi)_j = U_{\kappa_j}^* \text{Op}_h^w(\chi_j) U_{\kappa_j}$, each of them effectively acting from $L^2(\pi\mathcal{V}_j)$ to itself. We denote by $\delta_j(x; z)$ the kernel of the operator $\text{Op}(\chi)_j$: it is given by the integral

$$(3.2) \quad \delta_j(x; z) = (2\pi\hbar)^{-(D+2d)} \int e^{-\frac{i}{\hbar}S(x,y,\theta)} e^{\frac{i}{\hbar}\langle y-\tilde{y}, \eta \rangle} e^{\frac{i}{\hbar}S(z,\tilde{y},\tilde{\theta})} \\ \times \bar{a}_\hbar(x, y, \theta) a_\hbar(z, \tilde{y}, \tilde{\theta}) \varphi_j(y, \eta) \chi(\eta_1) \, dy \, d\theta \, d\tilde{y} \, d\tilde{\theta} \, d\eta.$$

For any wavefunction $\Psi \in L^2(M)$, we have therefore

$$(3.3) \quad [\text{Op}(\chi)\Psi](x) = \sum_j \int_{\pi\mathcal{V}_j} \Psi(z) \delta_j(x; z) \, dz.$$

We temporarily restore the dependence of $\delta_j(x; z)$ on the cutoffs, calling $\delta_j^{(n)}(x; z)$ the kernel of the operator $\text{Op}(\chi^{(n)})_j$. In order to prove Proposition 3.1, we will for each set (j, j', z, z') , obtain approximate expressions for the wavefunctions $U^t P_\epsilon \delta_j^{(n)}(z)$, respectively $P_{\epsilon'} \delta_{j'}^{(4n)}(z')$, and use these expressions to bound from above their overlaps:

LEMMA 3.2. — *Under the assumptions and notations of Proposition 3.1, the upper bound*

$$\left| \langle U^{-n/2} P_{\epsilon'} \delta_{j'}^{(4n)}(z'), U^{n/2} P_\epsilon \delta_j^{(n)}(z) \rangle \right| \leq C \hbar^{-(d-1)-c\delta} J_n^u(\epsilon)^{1/2} J_n^u(\epsilon').$$

holds uniformly for any j, j' , any points $z \in \pi\mathcal{V}_j$, $z' \in \pi\mathcal{V}_{j'}$ and any n -sequences ϵ, ϵ' .

Using (3.3) and the Cauchy-Schwarz inequality $\|\Psi\|_{L^1} \leq \sqrt{\text{Vol}(M)} \|\Psi\|_{L^2}$, this Lemma yields Proposition 3.1.

In the following sections we study the action of the operator P_ϵ on the state $\delta(z) = \delta_j^{(*)}(z)$ of the form (3.2). By induction on n , we propose an Ansatz for that state, valid for times $n = |\epsilon|$ of the order of $|\log \hbar|$. Apart from the sharp energy cutoff, this Ansatz is similar to the one described in [2].

3.3. WKB Ansatz for the first step

The first step of the evolution consists in applying the operator $U P_{\epsilon_0}$ to $\delta(z)$. For this aim, we will use the decomposition (3.2) into WKB states of the form $a(x) e^{iS(x)/\hbar}$, and evolve such states individually through the above operator. We briefly review how the propagator $U^t = e^{it\hbar\Delta/2}$ evolves such states.

Remark 3.3. — The \hbar -dependent presentation we use here is called the WKB method, from the work of Wentzell, Kramers and Brillouin on the Schrödinger equation. An alternative approach would have been to use the wave equation, and the Hadamard parametrix for the kernel of its propagator : see [16] and Bérard’s paper [4] for a modern presentation. An advantage of the wave equation lies in its finite speed of propagation. On the other hand, the fact that the Hamiltonian $\frac{|\xi|_x^2}{2}$ (vs. $|\xi|_x$) is smooth and strictly convex makes the parametrix of the Schrödinger equation somewhat simpler to write, and avoids certain technical difficulties.

In absence of conjugate points, if we consider a *fixed* energy cutoff χ compactly supported inside $\{|H(x, \xi) - 1/2| \leq \delta\}$, the parametrix of the Schrödinger equation reads :

$$(3.4) \quad [U^t \text{Op}_{\hbar}(\chi)](x, z) \sim \frac{e^{i\frac{d^2(x,z)}{2\hbar t}}}{(2i\pi\hbar t)^{d/2}} \sum_{k \geq 0} \hbar^k a_k(t, x, z),$$

this expansion being valid if x is such that there is at most one trajectory starting in the support of χ , and joining z to x in time t . In case there are several such trajectories (belonging to different homotopy classes), one needs to sum over their individual contributions. The coefficients a_k can be explicitly expressed in terms of χ , the geodesic flow and its derivatives, in particular a_0 is related to the Jacobian of the exponential map.

Paragraphs 3.3.1 and 3.3.2 recall the construction of this parametrix. Because our cutoff χ is not fixed, but its support shrinks rapidly as $\hbar \rightarrow 0$, we cannot apply the stationary phase method with respect to the energy parameter in the expression (3.13), which thus has to replace the simpler (3.4). We take the opportunity to introduce a certain number of notations and recall how the remainder terms behave in L^2 -norm.

3.3.1. Evolution of a WKB state

Consider an initial state $u(0)$ of the form $u(0, x) = a_{\hbar}(0, x)e^{\frac{i}{\hbar}S(0,x)}$, where $S(0, \bullet)$, $a_{\hbar}(0, \bullet)$ are smooth functions defined on a subset $\Omega \subset M$, and a_{\hbar} expands as $a_{\hbar} \sim \sum_k \hbar^k a_k$. This represents a WKB (or Lagrangian) state, supported on the Lagrangian manifold $\mathcal{L}(0) = \{(x, d_x S(0, x)), x \in \Omega\}$.

Then, for any integer N , the state $\tilde{u}(t) \stackrel{\text{def}}{=} U^t u(0)$ can be approximated, to order \hbar^N , by a WKB state $u(t)$ of the following form:

$$(3.5) \quad u(t, x) = e^{\frac{iS(t,x)}{\hbar}} a_{\hbar}(t, x) = e^{\frac{iS(t,x)}{\hbar}} \sum_{k=0}^{N-1} \hbar^k a_k(t, x).$$

Since we want $u(t)$ to solve $\frac{\partial u}{\partial t} = i\hbar \frac{\Delta_x u}{2}$ up to a remainder of order \hbar^N , the functions S and a_k must satisfy the following partial differential equations:

$$(3.6) \quad \begin{cases} \frac{\partial S}{\partial t} + H(x, d_x S) = 0 & \text{(Hamilton-Jacobi equation)} \\ \frac{\partial a_0}{\partial t} = -\langle d_x a_0, d_x S(t, x) \rangle - a_0 \frac{\Delta_x S(t, x)}{2} & \text{(0-th transport equation),} \\ \frac{\partial a_k}{\partial t} = \frac{i\Delta_x a_{k-1}}{2} - \langle d_x a_k, d_x S \rangle - a_k \frac{\Delta_x S}{2} & \text{(k-th transport equation).} \end{cases}$$

Assume that, on a certain time interval — say $s \in [0, 1]$ — the above equations have a well defined smooth solution $S(s, x)$, meaning that the transported Lagrangian manifold $\mathcal{L}(s) = g^s \mathcal{L}(0)$ is of the form $\mathcal{L}(s) = \{(x, d_x S(s, x))\}$, where $S(s)$ is a smooth function on the open set $\pi \mathcal{L}(s)$.

DEFINITION 3.4. — We shall say that a Lagrangian manifold \mathcal{L} is “projectible” if the projection $\pi : \mathcal{L} \rightarrow M$ is a diffeomorphism onto its image. If $\pi(\mathcal{L})$ is simply connected, this implies that \mathcal{L} is the graph of dS for some function S : we say that \mathcal{L} is generated by S .

We thus assume that $\mathcal{L}(s)$ is projectible for $s \in [0, 1]$, and is generated by $S(s)$. Under these conditions, we denote as follows the induced flow on M :

$$(3.7) \quad g_{S(s)}^t : x \in \pi \mathcal{L}(s) \mapsto \pi g^t(x, d_x S(s, x)) \in \pi \mathcal{L}(s+t).$$

This flow satisfies the property $g_{S(s+\tau)}^t \circ g_{S(s)}^\tau = g_{S(s)}^{t+\tau}$. We then introduce the following (unitary) operator $T_{S(s)}^t$, which transports functions on $\pi \mathcal{L}(s)$ into functions on $\pi \mathcal{L}(s+t)$:

$$(3.8) \quad T_{S(s)}^t(a)(x) = a \circ g_{S(s+t)}^{-t}(x) (J_{S(s+t)}^{-t}(x))^{1/2}.$$

Here $J_{S(s)}^t(x)$ is the Jacobian of the map $g_{S(s)}^t$ at the point x (measured with respect to the Riemannian volume on M). It is given by

$$(3.9) \quad J_{S(s)}^t(x) = \exp \left\{ \int_0^t \Delta S(s+\tau, g_{S(s)}^\tau(x)) d\tau \right\}.$$

The 0-th transport equation in (3.6) is explicitly solved by

$$(3.10) \quad a_0(t) = T_{S(0)}^t a_0, \quad t \in [0, 1],$$

and the higher-order terms $k \geq 1$ are given by

$$(3.11) \quad a_k(t) = T_{S(0)}^t a_k + \int_0^t T_{S(s)}^{t-s} \left(\frac{i\Delta_x a_{k-1}}{2}(s) \right) ds.$$

The function $u(t, x)$ defined by (3.5) satisfies the approximate equation

$$\frac{\partial u}{\partial t} = i\hbar \frac{\Delta_x u}{2} - i\hbar^N e^{\frac{i}{\hbar} S(t, x)} \frac{\Delta_x a_{N-1}}{2}(t, x).$$

From Duhamel’s principle and the unitarity of U^t , the difference between $u(t)$ and the exact solution $\tilde{u}(t)$ is bounded, for $t \in [0, 1]$, by

$$\begin{aligned}
 (3.12) \quad \|u(t) - \tilde{u}(t)\|_{L^2} &\leq \frac{\hbar^N}{2} \int_0^t \|\Delta a_{N-1}(s)\|_{L^2} ds \\
 &\leq Ct\hbar^N \left(\sum_{k=0}^{N-1} \|a_k(0)\|_{C^{2(N-k)}} \right).
 \end{aligned}$$

The constant C is controlled by the volumes of the sets $\pi\mathcal{L}(s)$ ($0 \leq s \leq 1$), and by a certain number of derivatives of the flow $g_{S(s+t)}^{-t}$ ($0 \leq s+t \leq 1$).

3.3.2. The Ansatz for time $n = 1$

We now apply the above analysis to study the evolution of the state $\delta(z)$ given by the integral (3.2). Until section 3.5.2, we will consider a single point z . Selecting in (3.2) a pair (y, θ) in the support of a_{\hbar} , we consider the state

$$u(0, x) = e^{-\frac{i}{\hbar}S(x,y,\theta)} \bar{a}_{\hbar}^{\epsilon_0}(x, y, \theta), \text{ where } a_{\hbar}^{\epsilon_0}(x, y, \theta) \stackrel{\text{def}}{=} P_{\epsilon_0}(x) a_{\hbar}(x, y, \theta).$$

Notice that this state is compactly supported in Ω_{ϵ_0} . We will choose a (large) integer $N > 0$ (see the condition at the very end of §3.6), truncate the \hbar -expansion of $\bar{a}_{\hbar}^{\epsilon_0}$ to the order $\tilde{N} = N + D + 2d$, and apply to that state the WKB evolution described in the previous section, up to order \tilde{N} and for times $0 \leq t \leq 1$. We then obtain an approximate state $\bar{a}_{\hbar}^{\epsilon_0}(t, x; y, \theta) e^{-\frac{i}{\hbar}S(t,x;y,\theta)}$. By the superposition principle, we get the following representation for the state $U^t P_{\epsilon_0} \delta(z)$:

$$(3.13) \quad [U^t P_{\epsilon_0} \delta(z)](x) = (2\pi\hbar)^{-\frac{d+1}{2}} \int v(t, x; z, \eta_1) \chi(\eta_1) d\eta_1 + \mathcal{O}_{L^2}(\hbar^N),$$

where for each energy parameter η_1 we took

$$\begin{aligned}
 (3.14) \quad v(t, x; z, \eta_1) &= (2\pi\hbar)^{-D-\frac{3d-1}{2}} \int e^{-\frac{i}{\hbar}S(t,x;y,\theta)} e^{\frac{i}{\hbar}\langle y-\tilde{y}, \eta \rangle} e^{\frac{i}{\hbar}S(z,\tilde{y},\tilde{\theta})} \\
 &\quad \times \bar{a}_{\hbar}^{\epsilon_0}(t, x; y, \theta) a_{\hbar}(z; \tilde{y}, \tilde{\theta}) \varphi_j(y, \eta) dy d\theta d\tilde{y} d\tilde{\theta} d\eta'
 \end{aligned}$$

(here $\eta' = (\eta_2, \dots, \eta_d)$). The reason why we integrate over all variables but η_1 lies in the sharp cutoff χ : due to this cutoff one cannot apply a stationary phase analysis in the variable η_1 .

At time $t = 0$, the state $v(0, \bullet, z, \eta_1)$ is a WKB state, supported on the Lagrangian manifold

$$\mathcal{L}_{\eta_1}^0(0) = \{ \rho \in \mathcal{E}(1/2 + \eta_1), \pi(\rho) \subset \Omega_{\epsilon_0}, \exists \tau \in [-1, 1], g^{-\tau} \rho \in T_z^* M \}.$$

This Lagrangian is obtained by propagating the sphere

$$S_{z,\eta_1}^* M = \left\{ \rho = (z, \xi), |\xi|_z = \sqrt{1 + 2\eta_1} \right\}$$

on the interval $\tau \in [-1, 1]$, and keeping only the points situated above Ω_{ϵ_0} . The projection of $\mathcal{L}_{\eta_1}^0(0)$ on M is not one-to-one: the point z has infinitely many preimages, while other points $x \in \Omega_{\epsilon_0}$ have in general two preimages (x, ξ_x) and $(x, -\xi_x)$.




Figure 3.1. Sketch of the Lagrangian manifold $\mathcal{L}_{\eta_1}^0(0)$ situated above Ω_{ϵ_0} and centered at z (center ellipse, dark grey), its image $\mathcal{L}_{\eta_1}^0(1)$ through the flow (external annulus, light grey) and the intersection $\mathcal{L}_{\eta_1}^1(0)$ of the latter with $T^*\Omega_{\epsilon_1}$. The thick arrows show the possible momenta at points $x \in M$ (black dots).

Let us assume that the diameter of the partition ε is less than $1/6$. For $0 < t \leq 1$, $v(t; z, \eta_1)$ is a WKB state supported on $\mathcal{L}_{\eta_1}^0(t) = g^t \mathcal{L}_{\eta_1}^0(0)$. If the time is small, $\mathcal{L}_{\eta_1}^0(t)$ still intersects $T_z^* M$. On the other hand, all points in $\mathcal{E}(1/2 + \eta_1)$ move at a speed $\sqrt{1 + 2\eta_1} \in [1 - 2\varepsilon, 1 + \varepsilon]$, so for times $t \in [3\varepsilon, 1]$ any point $x \in \pi \mathcal{L}_{\eta_1}^0(t)$ is at distance greater than ε from Ω_{ϵ_0} . Since the injectivity radius of M is ≥ 2 , such a point x is connected to z by a single short geodesic arc. Furthermore, since x is outside Ω_{ϵ_0} , there is

no ambiguity about the sign of the momentum at x : in conclusion, there is a unique $\rho \in \mathcal{L}_{\eta_1}^0(t)$ sitting above x (Fig. 3.1).

For times $3\varepsilon \leq t \leq 1$, the Lagrangian $\mathcal{L}_{\eta_1}^0(t)$ is therefore projectible, and it is generated by a function $S^0(t, \bullet; z, \eta_1)$. Equivalently, for any x in the support of $v(t, \bullet; z, \eta_1)$, the integral (3.14) is stationary at a unique set of parameters $\bullet_c = (y_c, \theta_c, \tilde{y}_c, \tilde{\theta}_c, \eta'_c)$, and leads to an expansion (up to order \hbar^N):

$$(3.15) \quad v(t; z, \eta_1) = v^0(t; z, \eta_1) + \mathcal{O}(\hbar^N),$$

$$\text{where } v^0(t, x; z, \eta_1) = e^{\frac{i}{\hbar} S^0(t, x; z, \eta_1)} b_{\hbar}^0(t, x; z, \eta_1).$$

The above discussion shows that $\mathcal{L}_{\eta_1}^0 \stackrel{\text{def}}{=} \cup_{3\varepsilon \leq t \leq 1} \mathcal{L}_{\eta_1}^0(t)$ is a projectible Lagrangian manifold which can be generated by a single function $S^0(\bullet; z, \eta_1)$ defined on $\pi \mathcal{L}_{\eta_1}^0$. The phase functions $S^0(t, \bullet; z, \eta_1)$ obtained through the stationary phase analysis depend very simply on time:

$$S^0(t, x; z, \eta_1) = S^0(x; z, \eta_1) - (1/2 + \eta_1)t.$$

The symbol b_{\hbar}^0 is given by a truncated expansion $b_{\hbar}^0 = \sum_{k=0}^{N-1} \hbar^k b_k^0$. The principal symbol reads

$$b_0^0(t, x; z, \eta_1) = \bar{a}_0^{\varepsilon_0}(t, x; y_c, \theta_c) a_0(z; \tilde{y}_c, \tilde{\theta}_c),$$

while higher order terms b_k^0 are given by linear combination of derivatives of $\bar{a}_{\hbar}^{\varepsilon_0}(t, x; \bullet) a_{\hbar}(z; \bullet)$ at the critical point $\bullet = \bullet_c$. Since $\bar{a}_{\hbar}^{\varepsilon_0}(0, \bullet; y_c, \theta_c)$ is supported inside Ω_{ε_0} , the transport equation (3.11) shows that $b_{\hbar}^0(t, \bullet; z, \eta_1)$ is supported inside $\pi \mathcal{L}_{\eta_1}^0(t)$.

If we take in particular $t = 1$, the state

$$(3.16) \quad v^0(1; z) = (2\pi\hbar)^{-\frac{d+1}{2}} \int v^0(1; z, \eta_1) \chi(\eta_1) d\eta_1$$

provides an approximate expression for $UP_{\varepsilon_0} \delta(z)$, up to a remainder $\mathcal{O}_{L^2}(|\text{supp } \chi| \hbar^{N - \frac{d+1}{2}})$.

3.4. Iteration of the WKB Ansätze

In this section we will obtain an approximate Ansatz for $P_{\varepsilon_n} \cdots UP_{\varepsilon_1} UP_{\varepsilon_0} \delta(z)$. Above we have already performed the first step, obtaining an approximation $v^0(1; z)$ of $UP_{\varepsilon_0} \delta(z)$, which was decomposed into fixed-energy WKB states $v^0(1; z, \eta_1)$. The next steps will be performed by evolving each component $v^0(1; z, \eta_1)$ individually, and integrating over η_1 only at the

end. Until Lemma 3.5 we will fix the variables (z, η_1) , and omit them in our notations when no confusion may arise.

Applying the multiplication operator P_{ϵ_1} to the state $v^0(1) = v^0(1; z, \eta_1)$, we obtain another WKB state which we denote as follows:

$$v^1(0, x) = b_h^1(0, x)e^{\frac{i}{h}S^1(0,x)}, \text{ with } \begin{cases} S^1(0, x) = S^0(1, x; z, \eta_1), \\ b_h^1(0, x) = P_{\epsilon_1}(x)b_h^0(1, x; z, \eta_1). \end{cases}$$

This state is associated with the manifold

$$\mathcal{L}^1(0) = \mathcal{L}_{\eta_1}^0(1) \cap T^*\Omega_{\epsilon_1}.$$

If this intersection is empty, then $v^1(0) = 0$, which means that $P_{\epsilon_1}Uv(0; z, \eta_1) = \mathcal{O}(\hbar^N)$. In the opposite case, we can evolve $v^1(0)$ following the procedure described in §3.3.1. For $t \in [0, 1]$, and up to an error $\mathcal{O}_{L^2}(\hbar^N)$, the evolved state $U^t v^1(0)$ is given by the WKB Ansatz

$$v^1(t, x) = b_h^1(t, x)e^{\frac{i}{h}S^1(t,x)}, \quad b_h^1(t) = \sum_{k=0}^{N-1} \hbar^k b_k^1(t).$$

The state $v^1(t)$ is associated with the Lagrangian $\mathcal{L}^1(t) = g^t \mathcal{L}^1(0)$, and the function $b_h^1(t)$ is supported inside $\pi \mathcal{L}^1(t)$. The Lagrangian $\mathcal{L}^1 \stackrel{\text{def}}{=} \cup_{0 \leq t \leq 1} \mathcal{L}^1(t)$ is generated by the function $S^1(0, x)$, and for any $t \in [0, 1]$ we have $S^1(t, x) = S^1(0, x) - (1/2 + \eta_1) t$.

3.4.1. Evolved Lagrangians

We can iterate this procedure, obtaining a sequence of approximations

$$(3.17) \quad v^j(t) = U^t P_{\epsilon_j} v^{j-1}(1) + \mathcal{O}(\hbar^N), \text{ where } v^j(t, x) = b_h^j(t, x)e^{\frac{i}{h}S^j(t,x)}.$$

To show that this procedure is consistent, we must check that the Lagrangian manifold $\mathcal{L}^j(t)$ supporting $v^j(t)$ is always projectible : that it does not develop caustics through the evolution ($t \in [0, 1]$), and that the projection $\pi : \mathcal{L}^j(t) \rightarrow M$ is injective. This will ensure that $\mathcal{L}^j(t)$ is generated by a function $S^j(t)$. We now show that these properties hold, due to the assumptions on the classical flow.

The manifolds $\mathcal{L}^j(t)$ are obtained by the following procedure. Knowing $\mathcal{L}^{j-1}(1)$, which is generated by the phase function $S^{j-1}(1)$, we take for $\mathcal{L}^j(0)$ the intersection

$$\mathcal{L}^j(0) = \mathcal{L}^{j-1}(1) \cap T^*\Omega_{\epsilon_j}.$$

If this set is empty, we then stop the construction. Otherwise, this Lagrangian is evolved into $\mathcal{L}^j(t) = g^t \mathcal{L}^j(0)$ for $t \in [0, 1]$. Notice that the

Lagrangian $\mathcal{L}^j(t)$ corresponds to the evolution at time $j + t$ of a piece of $\mathcal{L}^0(0)$; the latter is contained in the union $\cup_{|\tau| \leq 1} g^\tau S_{z, \eta_1}^* M$, where $S_{z, \eta_1}^* M$ is the sphere of energy $1/2 + \eta_1$ above z . If the geodesic flow is Anosov, it has no conjugate points [21]. This implies that $g^t \mathcal{L}^0(0)$ will not develop caustics: in other words, the phase functions $S^j(t)$ will never become singular.

On the other hand, when $j \rightarrow \infty$ the Lagrangian $g^{j+t} \mathcal{L}^0(0)$ is no longer projectible, it covers any $x \in M$ many times, so that several local generating functions are needed to describe the different sheets (see §3.5.3). However, the small piece $\mathcal{L}^j(t) \subset g^{j+t} \mathcal{L}^0(0)$ is generated by only one of them. Indeed, because the injectivity radius is ≥ 2 , any point $x \in \Omega_{\epsilon_j}$ can be connected to another point $x' \in M$ by at most one geodesic of length $\sqrt{1 + 2\eta_1} \leq 1 + \epsilon$. Also, if ϵ is small enough, it is impossible for two geodesics, issued from the same point z and going successively through the same Ω_{ϵ_j} at integer times, to intersect again at a later time : otherwise they would be homotopic, and thus give rise to two geodesics with same endpoints in the universal cover. This ensures that, for any $j \geq 1$, the manifold $\mathcal{L}^j = \cup_{t \in [0, 1]} \mathcal{L}^j(t)$ is projectible, and thus generated by a function $S^j(0)$ defined on $\pi \mathcal{L}^j$, or equivalently by $S^j(t) = S^j(0) - (1/2 + \eta_1) t$.

Finally, since the flow on $\mathcal{E}(1/2 + \eta_1)$ is Anosov, the sphere bundle $\{S_{z, \eta_1}^* M, z \in M\}$ is uniformly transverse to the strong stable foliation [21]. As a result, under the flow a piece of sphere becomes exponentially close to an unstable leaf when $t \rightarrow +\infty$. The Lagrangians \mathcal{L}^j thus become exponentially close to the weak unstable foliation as $j \rightarrow \infty$. This transversality argument is crucial in our choice to decompose the state Ψ into components $\delta_j(z)$.

3.4.2. Exponential decay of the symbols

We now analyze the behaviour of the symbols $b_{\hbar}^j(t, x)$ appearing in (3.17), when $j \rightarrow \infty$. These symbols are constructed iteratively: starting from the function $b_{\hbar}^{j-1}(1) = \sum_{k=0}^{N-1} b_k^{j-1}(1)$ supported inside $\pi \mathcal{L}^{j-1}(1)$, we define

$$(3.18) \quad b_{\hbar}^j(0, x) = P_{\epsilon_j}(x) b_{\hbar}^{j-1}(1, x), \quad x \in \pi \mathcal{L}^j(0).$$

The WKB procedure of §3.3.1 shows that for any $t \in [0, 1]$,

$$(3.19) \quad U^t v^j(0) = v^j(t) + R_N^j(t),$$

where the transported symbol $b_h^{j-1}(t) = \sum_{k=0}^{N-1} \hbar^k b_k^{j-1}(t)$ is supported inside $\pi\mathcal{L}^j(t)$. The remainder satisfies

$$(3.20) \quad \|R_N^j(t)\|_{L^2} \leq Ct\hbar^N \left(\sum_{k=0}^{N-1} \|b_k^j(0)\|_{C^{2(N-k)}} \right).$$

To control this remainder when $j \rightarrow \infty$, we need to bound from above the derivatives of b_h^j . Lemma 3.5 below shows that all terms $b_k^j(t)$ and their derivatives decay exponentially when $j \rightarrow \infty$, due to the Jacobian appearing in (3.8).

To understand the reasons of the decay, we first consider the principal symbols $b_0^j(1, x)$. They satisfy the following recurrence:

$$(3.21) \quad \begin{aligned} b_0^j(1, x) &= T_{S^j}^1(P_{\epsilon_j} \times b_0^{j-1}(1))(x) \\ &= P_{\epsilon_j}(g_{S^j}^{-1}(x))b_0^{j-1}(1, g_{S^j}^{-1}(x)) \sqrt{J_{S^j}^{-1}(x)}. \end{aligned}$$

Iterating this expression, and using the fact that $0 \leq P_{\epsilon_j} \leq 1$, we get at time n and for any $x \in \pi\mathcal{L}^n(0)$:

$$(3.22) \quad |b_0^n(0, x)| \leq |b_0^0(1, g_{S^n}^{-n+1}(x))| \times \left(J_{S^{n-1}}^{-1}(x)J_{S^{n-2}}^{-1}(g_{S^{n-2}}^{-1}(x)) \cdots J_{S^1}^{-1}(g_{S^n}^{-n+2}(x)) \right)^{1/2}.$$

Since the Lagrangians \mathcal{L}^j converge exponentially fast to the weak unstable foliation, the associated Jacobians satisfy for some $C > 0$:

$$\forall j \geq 2, \forall \rho = (x, \xi) \in \mathcal{L}^j(0), \left| \frac{J_{S^j}^{-1}(x)}{J_{S^u}^{-1}(\rho)(x)} - 1 \right| \leq Ce^{-j/C}.$$

Here $S^u(\rho)$ generates the local weak unstable manifold at the point ρ (which is a Lagrangian submanifold in $\mathcal{E}(1/2 + \eta_1)$). The product of Jacobians in (3.22) therefore satisfies, uniformly with respect to n and $\rho \in \mathcal{L}^n(0)$:

$$\begin{aligned} \prod_{j=1}^{n-1} J_{S^{n-j}}^{-1}(g_{S^n}^{-j+1}(x)) &= e^{\mathcal{O}(1)} \prod_{j=1}^{n-1} J_{S^u(g^{-j+1}\rho)}^{-1}(g_{S^n}^{-j+1}(x)) \\ &= e^{\mathcal{O}(1)} J_{S^u(\rho)}^{1-n}(x), \quad n \rightarrow \infty. \end{aligned}$$

The Jacobian $J_{S^u(\rho)}^{-1}$ measures the contraction of g^{-1} along $E^u(\rho)$: so does the Jacobian $J^u(\rho)$ defined in §2.2.5, but with respect to different coordinates. When iterating the contraction n times, the ratio of these Jacobians remains bounded:

$$J_{S^u(\rho)}^{1-n}(x) = e^{\mathcal{O}(1)} \prod_{j=1}^{n-1} J^u(g^{-j+1}\rho), \quad n \rightarrow \infty.$$

We finally express the upper bound in terms of the “coarse-grained” Jacobians (2.15, 2.16). Since $\rho \in \mathcal{L}^n(0) \subset T^*\Omega_{\epsilon_n}$ and $g^{-j}\rho \in T^*\Omega_{\epsilon_{n-j}}$ for all $j = 1, \dots, n - 1$, we obtain the following estimate on the principal symbol $b_0^n(0)$:

$$(3.23) \quad \forall n \geq 1, \quad \|b_0^n(0)\|_{L^\infty} \leq C \|b_0^0(1; z, \eta_1)\|_{L^\infty} J_{n-1}^u(\epsilon_1 \cdots \epsilon_n)^{1/2}.$$

The constant C only depends on the Riemannian manifold M . By construction the symbol $b_0^0(1; z, \eta_1)$ is bounded uniformly with respect to the variables (z, η_1) (assuming $|\eta_1| < \varepsilon$).

The following lemma shows that the above bound extends to the full symbol $b_h^n(0, x)$ and its derivatives (which are supported on $\pi\mathcal{L}^n(0)$).

LEMMA 3.5. — *Take any index $0 \leq k \leq N$ and $m \leq 2(N - k)$. Then there exists a constant $C(k, m)$ such that*

$$\forall n \geq 1, \quad \forall x \in \pi\mathcal{L}^n(0), \quad |d^m b_k^n(0, x)| \leq C(k, m) n^{m+3k} J_n^u(\epsilon_0 \cdots \epsilon_n)^{1/2}.$$

This bound is uniform with respect to the parameters (z, η_1) . For $(k, m) \neq (0, 0)$, the constant $C(k, m)$ depends on the partition $\mathcal{P}^{(0)}$, while $C(0, 0)$ does not.

Before giving the proof of this lemma, we draw some consequences. Taking into account the fact that the remainders $R_N^j(1)$ are dominated by the derivatives of the b_k^j (see (3.20)), the above statement translates into

$$\forall j \geq 1, \quad \|R_N^j(1)\|_{L^2} \leq C(N) j^{3N} J_j^u(\epsilon_0 \cdots \epsilon_j)^{1/2} \hbar^N.$$

A crucial fact for us is that the above bound also holds for the propagated remainder $P_{\epsilon_n} U \cdots U P_{\epsilon_{j+1}} R_N^j(1)$, due to the fact that the operators $P_{\epsilon_j} U$ are contracting. As a result, the total error at time n is bounded from above by the sum of the errors $\|R_N^j(1)\|_{L^2}$. We obtain the following estimate for any $n > 0$:

$$(3.24) \quad \|P_{\epsilon_n} U P_{\epsilon_{n-1}} \cdots P_{\epsilon_1} U v(0; z, \eta_1) - v^n(0; z, \eta_1)\|_{L^2} \leq C(N) \hbar^N \sum_{j=0}^n j^{3N} J_j^u(\epsilon_0 \cdots \epsilon_j)^{1/2}.$$

From the fact that the Jacobians J_j^u decay exponentially with j , the last term is bounded by $C(N) \hbar^N$. This bound is uniform with respect to the data (z, η_1) .

By the superposition principle, we obtain the following

COROLLARY 3.6. — For small enough $\hbar > 0$, any point $z \in \pi\mathcal{V}_j$, and any sequence ϵ of arbitrary length $n \geq 0$, we have

$$P_\epsilon \delta_j(z) = (2\pi\hbar)^{-\frac{d+1}{2}} \int v^n(0; z, \eta_1) \chi(\eta_1) d\eta_1 + \mathcal{O}_{L^2}(|\text{supp } \chi| \hbar^{N-\frac{d+1}{2}}).$$

Here we may take $\chi = \chi^{(n')}$ with an arbitrary $0 \leq n' \leq C_\delta |\log \hbar|$ (see (2.6) and the following discussion).

Proof of Lemma 3.5

The transport equation (3.10, 3.11) linking b^j to b^{j-1} ,

$$(3.25) \quad \begin{aligned} b_k^j(t) &= T_{S^j}^t b_k^j(0) + (1 - \delta_{k,0}) \int_0^t T_{S^j}^{t-s} \left(\frac{i \Delta b_{k-1}^j(s)}{2} \right) ds, \\ & \qquad \qquad \qquad k = 0, \dots, N-1, \\ b_k^j(0) &= P_{\epsilon_j} \times b_k^{j-1}(1), \end{aligned}$$

can be m times differentiated. We can write the recurrence equations for the m -differential forms $d^m b_k^j(t)$ as follows:

$$(3.26) \quad \begin{aligned} d^m b_k^j(t, x) &= \sum_{\ell \leq m} T_{S^j}^t d^\ell b_k^{j-1}(1, x) \cdot \theta_{m\ell}^j(t, x) \\ & \quad + \sum_{\ell \leq m} \int_0^t T_{S^j}^{t-s} d^{\ell+2} b_{k-1}^j(s, x) \cdot \alpha_{m\ell}^j(t, s, x) ds. \end{aligned}$$

Above we have extended the transport operator T_S^t defined in (3.8) to multi-differential forms on M . Namely,

$$(T_{S^j}^t d^\ell b)(x) \stackrel{\text{def}}{=} \sqrt{J_{S^j}^{-t}(x)} d^\ell b(g_{S^j}^{-t}(x))$$

is an ℓ -form on $(T_{g_S^{-t}(x)} M)^\ell$. The linear form $\theta_{m\ell}^j(t, x)$ sends $(T_x M)^m$ to $(T_{g_{S^j}^{-t}(x)} M)^\ell$ (resp. $\alpha_{m\ell}^j(t, s, x)$ sends $(T_x M)^m$ to $(T_{g_{S^j}^{s-t}(x)} M)^{\ell+2}$). These forms can be expressed in terms of derivatives of the maps $g_{S^j}^{-t}, g_{S^j}^{s-t}$ at the point x , and $\theta_{m\ell}^j$ also depends on $m - \ell$ derivatives of the function P_{ϵ_j} . These forms are uniformly bounded with respect to j, x and $t \in [0, 1]$. We only need to know the explicit expression for θ_{mm}^j :

$$(3.27) \quad \theta_{mm}^j(t, x) = P_{\epsilon_j}(g_{S^j}^{-t}(x)) \times \left(dg_{S^j}^{-t}(x) \right)^{\otimes m}.$$

Since the above expressions involve several sets of parameters, to facilitate the bookkeeping we arrange the functions $b_k^j(t, x)$ and the m -differential

forms $d^m b_k^j(t, x)$, $m \leq 2(N - k)$, inside a vector \mathbf{b}^j . We will denote the entries by $\mathbf{b}_{(k,m)}^j = d^m b_k^j$, and with $0 \leq k \leq N - 1$, $m \leq 2(N - k)$:

$$(3.28) \quad \mathbf{b}^j = \mathbf{b}^j(t, x) \stackrel{\text{def}}{=} \begin{pmatrix} b_0^j, db_0^j, \dots, d^{2N} b_0^j, \\ b_1^j, db_1^j, \dots, d^{2(N-1)} b_1^j, \\ \dots, \\ b_{N-1}^j, db_{N-1}^j, d^2 b_{N-1}^j. \end{pmatrix}$$

The set of recurrence equations (3.26) may then be cast in a compact form, using three operator-valued matrices \mathbf{M}_*^j (here the subscript j is not a power, but refers to the Lagrangian \mathcal{L}^j on which the transformation is based):

$$(3.29) \quad (\mathbf{I} - \mathbf{M}_1^j) \mathbf{b}^j = (\mathbf{M}_{0,0}^j + \mathbf{M}_{0,1}^j) \mathbf{b}^{j-1}.$$

The first matrix acts as follows on the indices (k, m) :

$$(\mathbf{M}_1^j \mathbf{b}^j)_{(k,m)}(t) = \sum_{\ell \leq m} \int_0^t ds T_{S^j}^{t-s} \mathbf{b}_{(k-1, \ell+2)}^j(s) \cdot \alpha_{m\ell}^j(t, s).$$

Since \mathbf{M}_1^j relates b_k to b_{k-1} , it is obviously a nilpotent matrix of order N . The matrix $\mathbf{M}_{0,1}^j$:

$$(\mathbf{M}_{0,1}^j \mathbf{b}^{j-1})_{(k,m)}(t) = \sum_{\ell < m} T_{S^j}^t \mathbf{b}_{(k,\ell)}^{j-1}(1) \cdot \theta_{m\ell}^j(t),$$

which relates m -derivatives to ℓ -derivatives, $\ell < m$, is also nilpotent. Finally, the last matrix $\mathbf{M}_{0,0}^j$ acts diagonally on the indices (k, m) :

$$(3.30) \quad (\mathbf{M}_{0,0}^j \mathbf{b}^{j-1})_{(k,m)}(t) = T_{S^j}^t \mathbf{b}_{(k,m)}^{j-1}(1) \cdot \theta_{mm}^j(t).$$

From the nilpotence of \mathbf{M}_1^j , we can invert (3.29) into

$$\mathbf{b}^j = \left(\sum_{k=0}^{N-1} [\mathbf{M}_1^j]^k \right) (\mathbf{M}_{0,0}^j + \mathbf{M}_{0,1}^j) \mathbf{b}^{j-1},$$

where $[\mathbf{M}]^k$ denotes the k -th power of the matrix \mathbf{M} . The above expression can be iterated:

$$(3.31) \quad \mathbf{b}^n = \sum_{k_1, \dots, k_n=0}^{N-1} \sum_{\alpha_1, \dots, \alpha_n=0}^1 [\mathbf{M}_1^n]^{k_n} \mathbf{M}_{0,\alpha_n}^n [\mathbf{M}_1^{n-1}]^{k_{n-1}} \mathbf{M}_{0,\alpha_{n-1}}^{n-1} \dots [\mathbf{M}_1^1]^{k_1} \mathbf{M}_{0,\alpha_1}^1 \mathbf{b}^0.$$

Notice that the first step $\mathbf{M}_{0,\alpha_1}^1 \mathbf{b}^0$ only uses the vector \mathbf{b}^0 at time $t = 1$, where it is well-defined.

From the nilpotence properties of \mathbf{M}_1^j and $\mathbf{M}_{0,1}^j$, the terms contributing to $\mathbf{b}_{(k,m)}^n$ must satisfy $\sum k_j \leq k$ and $\sum \alpha_j \leq m + 2(\sum k_j)$. In particular, $\sum k_j \leq N$, $\sum \alpha_j \leq 2N$, so for n large, all terms in (3.31) are made of few (long) strings of successive matrices $\mathbf{M}_{0,0}^j$, separated by a few matrices $\mathbf{M}_{0,1}^j$ or \mathbf{M}_1^j (the total number of matrices $\mathbf{M}_{0,1}^j$ or \mathbf{M}_1^j in each term is at most $3N$). As a result, the total number of terms in the expression (3.31) of $\mathbf{b}_{(k,m)}^n$ grows at most like $\mathcal{O}(n^{m+3k})$ when $n \rightarrow \infty$.

Using the fact that $\theta_{m\ell}^j$ and $\alpha_{m\ell}^j$ are uniformly bounded, the actions of the nilpotent matrices \mathbf{M}_1^j , $\mathbf{M}_{0,1}^j$ induce the following bounds on the sup-norm of $\mathbf{b}_{k,m}^j(t)$:

$$(3.32) \quad \begin{aligned} \sup_{0 \leq t \leq 1} \|\mathbf{M}_1^j \mathbf{b}_{(k,m)}^j(t)\|_{L^\infty} &\leq C \max_{m' \leq m+2} \sup_{0 \leq t \leq 1} \|\mathbf{b}_{(k-1,m')}^j(t)\|_{L^\infty}, \\ \sup_{0 \leq t \leq 1} \|(\mathbf{M}_{0,1}^j \mathbf{b}^{j-1})_{(k,m)}(t)\|_{L^\infty} &\leq C(m) \max_{m' \leq m-1} \|\mathbf{b}_{(k,m')}^{j-1}(1)\|_{L^\infty}. \end{aligned}$$

The constant $C(m)$ depends on the partition $\mathcal{P}^{(0)}$: for a partition of diameter ε , it is of order ε^{-m} .

On the other hand, for any pair (k, m) , the ‘‘diagonal’’ action (3.30) on $\mathbf{b}_{(k,m)}^j$ is very similar with its action on $\mathbf{b}_{(0,0)}^j$, which is the recurrence relation (3.21). The only difference comes from the appearance of the m -forms θ_{mm}^j instead of the functions θ_{00}^j . From the explicit expression (3.27) and the fact that $0 \leq P_{\varepsilon_j} \leq 1$, one easily gets

$$|(\mathbf{M}_{0,0}^j \mathbf{b}^{j-1})_{(k,m)}(t, x)| \leq \sqrt{J_{S_j}^{-t}(x)} |dg_{S_j}^{-t}(x)|^m |\mathbf{b}_{(k,m)}^{j-1}(1, g_{S_j}^{-t}(x))|.$$

By contrast with (3.32), in the above bound there is no potentially large constant prefactor in front of the right hand side. This allows us to iterate this inequality, and obtain a bound similar with (3.22). Indeed, using the composition of the maps $g_{S_j}^{-1}$ and their derivatives, we get for any $j, j' \in \mathbb{N}$ and $t \in [0, 1]$:

$$(3.33) \quad \begin{aligned} |(\mathbf{M}_{0,0}^{j+j'} \dots \mathbf{M}_{0,0}^j \mathbf{b}^{j-1})_{(k,m)}(t, x)| \\ \leq \sqrt{J_{S^{j'+j}}^{-t-j'}(x)} |dg_{S^{j'+j}}^{-t-j'}(x)|^m |\mathbf{b}_{(k,m)}^{j-1}(1, g_{S^{j'+j}}^{-t-j'}(x))|. \end{aligned}$$

As we explained above, the flow g^t acting on \mathcal{L}^j is asymptotically expanding except in the flow direction, because $g^t \mathcal{L}^j$ converges to the weak unstable manifold. As a result, the inverse flow $g^{-j'}$ acting on $\mathcal{L}^{j+j'} \subset g^{j'} \mathcal{L}^j$, and its projection $g_{S^{j'+j}}^{-j'}$, have a tangent map $dg_{S^{j'+j}}^{-j'}$ uniformly bounded with respect to j, j' . In each ‘‘string’’ of operators $\mathbf{M}_{0,0}^*$, the factor $dg_S^{-j'}$ can be replaced by a uniform constant. For each term in (3.31), we can then

iteratively combine the bounds (3.32, 3.33), to get

$$|(\mathbf{M}^n \mathbf{M}^{n-1} \dots \mathbf{M}^1 \mathbf{b}^0)_{(k,m)}(t, x)| \leq C \sqrt{J_{S^n}^{-t-n+1}(x)} \|\mathbf{b}^0(1)\|$$

Summing over those terms, we obtain

$$(3.34) \quad |\mathbf{b}_{(k,m)}^n(t, x)| \leq \tilde{C}(k, m) n^{m+3k} \sqrt{J_{S^n}^{-t-n+1}(x)} \|\mathbf{b}^0(1)\|.$$

The Jacobian on the right hand side is the same as in the bound (3.22). We can thus follow the same reasoning and replace $J_{S^n}^{-t-n+1}$ by $J_n^u(\epsilon)$ to obtain the lemma. \square

This ends the proof of Lemma 3.5 and Corollary 3.6. We proceed with the proof of our main Lemma 3.2, and now describe the states $U^{-n/2} P_{\epsilon'} \delta_{j'}^{(4n)}(z')$ and $U^{n/2} P_{\epsilon} \delta_j^{(n)}(z)$.

3.5. Evolution under $U^{-n/2}$ and $U^{n/2}$

Applying Corollary 3.6 with $n' = 4n$, resp. $n' = n$, we have approximate expressions for the states appearing in Lemma 3.2:

$$(3.35) \quad P_{\epsilon} \delta_j^{(n)}(z) = (2\pi\hbar)^{-\frac{d+1}{2}} \int v^n(0; z, \eta_1, \epsilon) \chi^{(n)}(\eta_1) d\eta_1 + \mathcal{O}_{L^2}(e^{n\delta} \hbar^{N-\frac{d-1}{2}}),$$

$$(3.36) \quad P_{\epsilon'} \delta_{j'}^{(4n)}(z') = (2\pi\hbar)^{-\frac{d+1}{2}} \int v^n(0; z', \eta'_1, \epsilon') \chi^{(4n)}(\eta'_1) d\eta'_1 + \mathcal{O}_{L^2}(e^{4n\delta} \hbar^{N-\frac{d-1}{2}}),$$

we notice that for $n \leq n_E(\hbar)$ the remainders are of the form $\mathcal{O}(\hbar^{N-N_1})$ for some fixed N_1 .

To prove the bound of Lemma 3.2, we assume n is an even integer, and consider the individual overlaps

$$(3.37) \quad \left\langle U^{-n/2} v^n(0; z', \eta'_1, \epsilon'), U^{n/2} v^n(0; z, \eta_1, \epsilon) \right\rangle,$$

Until the end of the section, we will fix z, η_1, z', η'_1 and omit them in the notations. On the other hand, we will sometimes make explicit the dependence on the sequences ϵ', ϵ . We then need to understand the states $U^{-n/2} v^n(0; \epsilon')$ and $U^{n/2} v^n(0; \epsilon)$.

3.5.1. Evolution under $U^{-n/2}$

We use WKB approximations to describe the backwards-evolved state $U^{-t}v^n(0; \epsilon')$. Before entering into the details, let us sketch the backwards evolution of the Lagrangian $\mathcal{L}^n = \mathcal{L}^n(0; \epsilon')$ supporting $v^n(0) = v^n(0; \epsilon')$ (for a moment we omit to indicate the dependence in ϵ'). Since \mathcal{L}^n had been obtained by evolving \mathcal{L}^0 and truncating it at each step, for any $0 \leq t \leq n-1$, the Lagrangian $\mathcal{L}^n(-t) \stackrel{\text{def}}{=} g^{-t}\mathcal{L}^n$ will be contained in $\mathcal{L}^{n-\lfloor t \rfloor-1}(1 - \{t\})$, where we decomposed the time t into its integral and fractional part. This Lagrangian projects well onto the base manifold, and is generated by the function $S^n(-t) = S^{n-\lfloor t \rfloor-1}(1 - \{t\})$ (which satisfies the Hamilton-Jacobi equation for negative times). This shows that the WKB method of §3.3.1, applied to the backwards flow U^{-t} acting on $v^n(0)$, can be *formally* used for all times $0 \leq t \leq n-1$. The evolved state can be written as

$$(3.38) \quad U^{-t}v^n(0) = v^n(-t) + \hat{R}_N(-t),$$

and $v^n(-t)$ has the WKB form

$$(3.39) \quad v^n(-t) = b_h^n(-t)e^{iS^n(-t)/\hbar}, \quad b_h^n(-t) = \sum_{k=0}^{N-1} \hbar^k b_k^n(-t).$$

The symbols $b_k^n(-t)$ are obtained from $b_k^n(0)$ using the backwards transport equations (see Eqs. (3.10, 3.11)):

$$(3.40) \quad b_0^n(-t) = T_{S^n(0)}^{-t} b_0^n(0) = (J_{S^n(-t)}^t)^{1/2} b^n(0) \circ g_{S^n(-t)}^t,$$

$$(3.41) \quad b_k^n(-t) = T_{S^n(0)}^{-t} b_k^n(0) - \int_0^t T_{S^n(-t)}^{-t+s} \left(\frac{i \Delta b_{k-1}^n}{2}(-s) \right) ds.$$

These symbols are supported on $\pi\mathcal{L}^n(-t)$. We need to estimate their C^m norms uniformly in t . The inverse of the Jacobian $J_{S^n(-t)}^t$ approximately measures the volume of the Lagrangian $\mathcal{L}^n(-t)$. Since the latter remains close to the weak unstable manifold as long as $n-t \gg 1$, the backwards flow has the effect to shrink it along the unstable directions. Thus, for $n-1 \geq t \gg 1$, $\mathcal{L}^n(-t)$ consist in a thin, elongated subset of $\mathcal{L}^{n-\lfloor t \rfloor-1}$ (see figure 3.2), with a volume of order

$$(3.42) \quad \text{Vol}(\mathcal{L}^n(-t)) \leq C \left(\inf_x J_{S^n(-t)}^t(x) \right)^{-1} \\ \leq C J_{[t]}^u(\epsilon'_{n-\lfloor t \rfloor} \cdots \epsilon'_n), \quad 0 \leq t \leq n-1.$$

When differentiating $b_0^n(-t)$, the derivatives of the flow $g_{S^n(-t)}^t$ also appear. Since $\mathcal{L}^n(-t)$ is close to the weak unstable manifold, the derivatives become

large as $t \gg 1$:

$$|\partial_x^\alpha g_{S^n(-t)}^t(x)| \leq C(\alpha) e^{t\lambda_+},$$

where $\lambda_+ \stackrel{\text{def}}{=} \lambda_{\max}(1 + \delta'/2)$, $0 \leq t \leq n - 1$, $x \in \pi\mathcal{L}^n(-t)$.

Hence, for any $t \leq n - 1$ and index $0 \leq m \leq 2N$ the m -derivatives of the principal symbol can be bounded as follows:

(3.43)

$$\begin{aligned} \forall t \leq n - 1, |d^m b_0^n(-t, x)| &\leq C(J_{S^n(-t)}^t(x))^{1/2} |dg_{S^n(-t)}^t(x)|^m \|b_0^n(0)\|_{C^m} \\ &\leq C J_{[t]}^u(\epsilon'_{n-[t]} \cdots \epsilon'_n)^{-1/2} e^{tm\lambda_+} \|b_0^n(0)\|_{C^m} \\ &\leq C J_{n-[t]}^u(\epsilon'_0 \cdots \epsilon'_{n-[t]})^{1/2} e^{tm\lambda_+}. \end{aligned}$$

In the last line we used the estimates of Lemma 3.5 for $\|b^n(0)\|_{C^m}$. From now on we will abbreviate $J_{n-[t]}^u(\epsilon'_0 \cdots \epsilon'_{n-[t]})$ by $J_{n-[t]}^u(\epsilon')$. By iteration, we similarly estimate the derivatives of the higher-order symbols ($k < N$, $m \leq 2(N - k)$):

(3.44) $\forall t \leq n - 1, |d^m b_k^n(-t, x)| \leq C J_{n-[t]}^u(\epsilon')^{1/2} e^{t(m+2k)\lambda_+}.$

We see that the higher-order symbols may grow faster (with t) than the principal one. As a result, when t becomes too large, the expansion (3.39) does not make sense any more, since the remainder in (3.38) may become larger than the main term. From (3.12), this remainder is bounded by

$$\|\hat{R}_N(-t)\| \leq \frac{\hbar^N}{2} \int_0^t \|\Delta b_{N-1}^n(-s)\| ds \leq C \hbar^N e^{t2N\lambda_+} J_{n-[t]}^u(\epsilon')^{1/2}.$$

This remainder remains smaller than the previous terms if $t \leq n_E(\hbar)/2$. Since we assume $n \leq n_E(\hbar)$, the WKB expansion still makes sense if we take $t = n/2$. To ease the notations in the following sections, we call $w^{n/2} \stackrel{\text{def}}{=} v^n(-n/2)$ the WKB state approximating $U^{-n/2}v^n(0)$, its phase function $S^{n/2} = S^n(-n/2)$ and its symbol $c_{\hbar}^{n/2}(x) \stackrel{\text{def}}{=} b_{\hbar}^n(-n/2, x)$, all these data depending on ϵ' . The above discussion shows that

(3.45) $\|U^{-n/2}v^n(0; \epsilon') - w^{n/2}(\epsilon')\| = \|\hat{R}_N(-n/2)\| \leq C \hbar^{N\delta'/2} J_{n/2}^u(\epsilon')^{1/2}.$

We will select an integer N large enough ($N\delta' \gg 1$), so that the above remainder is smaller than the estimate $J_n^u(\epsilon')^{1/2}$ we have on $\|v^n(\epsilon')\|$.

3.5.2. Evolution under $U^{n/2}$

We now study the forward evolution $U^{n/2}v^n(0; \epsilon)$. From now on we omit to indicate the dependence in the parameter $t = 0$. Using the smooth

partition (2.4), we decompose $U^{n/2}$ as:

$$U^{n/2} = \sum_{\alpha_i, 1 \leq i \leq n/2} P_{\alpha_{n/2}}^2 U P_{\alpha_{n/2-1}}^2 U \cdots P_{\alpha_1}^2 U \stackrel{\text{def}}{=} \sum_{\alpha} Q_{\alpha}.$$

The operators (Q_{α}) are very similar with the (P_{α}) of Eq. (2.5): the cutoffs P_k are replaced by their squares P_k^2 . As a result, the iterative WKB method presented in the previous sections can be adapted to obtain approximate expressions for each state $Q_{\alpha} v^n(\epsilon)$, similarly as in (3.24):

$$\begin{aligned} Q_{\alpha} v^n(\epsilon) &= v^{\frac{3}{2}n}(\epsilon\alpha) + \mathcal{O}_{L^2}(\sqrt{J_n^u(\epsilon)} \hbar^N), \\ v^{\frac{3}{2}n}(x; \epsilon\alpha) &= b_{\hbar}^{\frac{3}{2}n}(x; \epsilon\alpha) e^{\frac{i}{\hbar} S^{\frac{3}{2}n}(x; \epsilon\alpha)}. \end{aligned}$$

Here $\epsilon\alpha$ is the sequence of length $3n/2$ with elements $\epsilon_0 \cdots \epsilon_n \alpha_1 \cdots \alpha_{n/2}$. That state is localized on the Lagrangian manifold $\mathcal{L}^{\frac{3}{2}n}(\epsilon\alpha)$. The symbols $b_k^{\frac{3}{2}n}(\epsilon\alpha)$ and their derivatives satisfy the bounds of Lemma 3.5. The state $U^{n/2} v^n(\epsilon)$ is therefore given by a sum of contributions

$$(3.46) \quad U^{n/2} v^n(\epsilon) = \sum_{\alpha} v^{\frac{3}{2}n}(\epsilon\alpha) + \mathcal{O}_{L^2}(\hbar^{N-N_K}).$$

Here N_K is a constant depending on the cardinal K of the partition $\mathcal{P}^{(0)}$, and we assumed $n \leq n_E(\hbar)$. The integer N will be taken large enough, such that \hbar^{N-N_K} is smaller than the remainder appearing in (3.45).

3.5.3. Grouping terms into connected Lagrangian leaves

To compute the overlap (3.37), we do not need the full sum (3.46), but only the components α such that the support of $v^{\frac{3}{2}n}(\epsilon\alpha)$ intersects the support of $w^{n/2}(\epsilon')$, which is inside $\Omega_{\epsilon'_{n/2}}$. Thus, we can restrict ourselves to the set of sequences

$$A \stackrel{\text{def}}{=} \left\{ \alpha : \pi \mathcal{L}^{\frac{3}{2}n}(\epsilon\alpha) \cap \Omega_{\epsilon'_{n/2}} \neq \emptyset \right\} \subset \{1, \dots, K\}^{n/2}.$$

For $n \gg 1$, the Lagrangian $\bigcup_{\alpha \in A} \mathcal{L}^{\frac{3}{2}n}(\epsilon\alpha)$, which is a strict subset of $g^{n/2} \mathcal{L}^n(\epsilon)$, is the disjoint union of a large number of *connected leaves*, which we denote by $\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell)$, $\ell \in [1, L]$ (see Figure 3.2). Each leaf $\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell)$ corresponds to geodesics of length $n/2$ from Ω_{ϵ_n} to $\Omega_{\epsilon'_{n/2}}$ in a definite homotopy class. As a consequence, if ρ, ρ' belong to two different leaves $\ell \neq \ell'$, there must be a time $0 < t < \frac{n}{2}$ such that the backwards images $g^{-t}\rho, g^{-t}\rho'$ are at a distance larger than $D > 0$ (D is related to the injectivity




Figure 3.2. Decomposition of $(g^{n/2}\mathcal{L}^n(\epsilon)) \cap T^*\Omega_{\epsilon'_{n/2}}$ into connected leaves (here we show two of them, in light grey). The leaf ℓ contains the components $\mathcal{L}^{\frac{3}{2}n}(\epsilon\alpha)$, $\mathcal{L}^{\frac{3}{2}n}(\epsilon\beta)$ while the leaf ℓ_0 contains $\mathcal{L}^{\frac{3}{2}n}(\epsilon\gamma)$. We also show the elongated leaf $g^{-n/2}\mathcal{L}^n(\epsilon')$ supporting the state $w^{n/2}(\epsilon')$ (black). This state might interfere with $v^{\frac{3}{2}n}(\epsilon, \ell_0)$, but not with $v^{\frac{3}{2}n}(\epsilon, \ell)$ or any other leaf above $\Omega_{\epsilon'_{n/2}}$.

radius). The total number of leaves above $\Omega_{\epsilon'_{n/2}}$ can grow at most like the full volume of $g^{n/2}\mathcal{L}(\epsilon)$, so that

$$L \leq C\epsilon^{n(d-1)\lambda_+/2} \leq C\hbar^{-(d-1)/2}.$$

Each leaf $\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell)$ is the union of a certain number of components $\mathcal{L}^{\frac{3}{2}n}(\epsilon\alpha)$, and we group the corresponding sequences α into the subset $A_\ell \subset \{1, \dots,$

$K\}^{n/2}$:

$$\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell) = \bigcup_{\alpha \in A_\ell} \mathcal{L}^{\frac{3}{2}n}(\epsilon\alpha).$$

We obviously have $A = \bigsqcup_\ell A_\ell$. All components $\mathcal{L}^{\frac{3}{2}n}(\epsilon\alpha)$ with $\alpha \in A_\ell$ are generated by the same phase function $S^{\frac{3}{2}n}(\epsilon\alpha) \stackrel{\text{def}}{=} S^{\frac{3}{2}n}(\epsilon, \ell)$, so that the state

$$(3.47) \quad v^{\frac{3}{2}n}(x; \epsilon, \ell) \stackrel{\text{def}}{=} \sum_{\alpha \in A_\ell} v^{\frac{3}{2}n}(x; \epsilon\alpha) = b_{\frac{3}{2}n}^{\frac{3}{2}n}(x; \epsilon, \ell) e^{\frac{i}{\hbar} S^{\frac{3}{2}n}(x; \epsilon, \ell)}$$

is a Lagrangian state supported on $\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell)$, with symbol

$$b_{\frac{3}{2}n}^{\frac{3}{2}n}(x; \epsilon, \ell) = \sum_{\alpha \in A_\ell} b_{\frac{3}{2}n}^{\frac{3}{2}n}(x; \epsilon\alpha).$$

By inspection one can check that, at each point $\rho \in \mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell)$, the above sum over $\alpha \in A_\ell$ has the effect to insert partitions of unity $\sum_k P_k^2 = 1$ at each preimage $g^{-j}(\rho)$, $j = 0, \dots, \frac{n}{2} - 1$. As a result, the principal symbol will satisfy the same type of upper bound as in (3.22):

$$\begin{aligned} |b_0^{\frac{3}{2}n}(x; \epsilon, \ell)| &\leq |b^n(g_S^{-n/2}(x))| J_S^{-\frac{1}{2}n}(x)^{1/2} \\ &\leq C J_S^{-\frac{3}{2}n}(x)^{1/2}, \text{ with } S = S^{\frac{3}{2}n}(\epsilon, \ell). \end{aligned}$$

The same argument holds for the higher-order terms and their derivatives. Besides, because the action of $g^{-3n/2}$ on $\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell)$ is contracting, for any $x \in \Omega_{\epsilon'}_{n/2}$ the Jacobian $J_S^{-\frac{3}{2}n}(x)$ is of the order of $J_{\frac{3}{2}n}^u(\epsilon\alpha)$, where α can be any sequence in A_ℓ (all these Jacobians are of the same order). Defining

$$J_{\frac{3}{2}n}^u(\epsilon, \ell) = \max_{\alpha \in A_\ell} J_{\frac{3}{2}n}^u(\epsilon\alpha) \geq \frac{1}{C} \min_{\alpha \in A_\ell} J_{\frac{3}{2}n}^u(\epsilon\alpha),$$

the full symbol $b_{\frac{3}{2}n}^{\frac{3}{2}n}(x; \epsilon, \ell)$ satisfies similar bounds as in Lemma 3.5:

$$(3.48) \quad |d^m b_k^{\frac{3}{2}n}(x; \epsilon, \ell)| \leq C n^{m+3k} J_{\frac{3}{2}n}^u(\epsilon, \ell)^{1/2}, \quad k \leq N - 1, \quad m \leq 2(N - k).$$

Remark 3.7. — As pointed out by the referee, the WKB Ansatz for $U^{n/2}$ on M could be deduced from a similar (but simpler) Ansatz for the flow on the universal cover \tilde{M} : the former is obtained by summing the contributions of geodesics in different homotopy classes [4, Section E]. Proceeding this way, one directly obtains, instead of (3.46), the sum over the components $v^{\frac{3}{2}n}(x; \epsilon, \ell)$ of (3.47), each $\ell \in [1, L]$ labelling a certain homotopy class.

3.6. Overlaps between the Lagrangian states

Putting together (3.45, 3.47, 3.46), the overlap (3.37) is approximated by the following sum:

$$(3.49) \quad \langle U^{-n/2}v^n(\epsilon'), U^{n/2}v^n(\epsilon) \rangle = \sum_{\ell=1}^L \langle w^{n/2}(\epsilon'), v^{\frac{3}{2}n}(\epsilon, \ell) \rangle + \mathcal{O}(\hbar^{N\delta'/2}),$$

where

$$\langle w^{n/2}(\epsilon'), v^{\frac{3}{2}n}(\epsilon, \ell) \rangle = \int e^{\frac{i}{\hbar}(S^{\frac{3}{2}n}(x; \epsilon, \ell) - S^{n/2}(x; \epsilon'))} \bar{c}_\hbar^{n/2}(x; \epsilon') b_\hbar^{\frac{3}{2}n}(x; \epsilon, \ell).$$

Each term is the overlap between the WKB state $w^{n/2}(\epsilon')$ supported on $g^{-n/2}\mathcal{L}^n(\epsilon')$, and the WKB state $v^{\frac{3}{2}n}(\epsilon, \ell)$ supported on $\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell)$, both Lagrangians sitting above $\Omega_{\epsilon'_{n/2}}$ (see Figure 3.2). The sup-norms of these two states, governed by the principal symbols $c_0^{n/2}(\epsilon')$, $b_0^{\frac{3}{2}n}(\epsilon, \ell)$, are bounded by

$$(3.50) \quad \|w^{n/2}(\epsilon')\|_{L^\infty} \leq C J_{n/2}^u(\epsilon')^{1/2}, \quad \|v^{\frac{3}{2}n}(\epsilon, \ell)\|_{L^\infty} \leq C J_{\frac{3}{2}n}^u(\epsilon, \ell)^{1/2}.$$

The integral (3.49) takes place on the support of $c_\hbar^{n/2}(x; \epsilon')$, that is (see (3.42)), on a set of volume $\mathcal{O}(J_{n/2}^u(\epsilon'_{n/2} \cdots \epsilon'_n))$. It follows that each overlap (3.6) is bounded by

$$(3.51) \quad |\langle w^{n/2}(\epsilon'), v^{\frac{3}{2}n}(\epsilon, \ell) \rangle| \leq C J_{n/2}^u(\epsilon')^{1/2} J_{\frac{3}{2}n}^u(\epsilon, \ell)^{1/2} J_{n/2}^u(\epsilon'_{n/2} \cdots \epsilon'_n).$$

We show below that the above estimate can be improved for almost all leaves ℓ , when one takes into account the phases in the integrals (3.6). Actually, for times $n \leq n_E(\hbar)$, there is at most a single term ℓ_0 in the sum (3.49) for which the above bound is sharp; for all other terms ℓ , the phase oscillates fast enough to make the integral negligible. Geometrically, this phase oscillation means that the Lagrangians $\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell)$, $g^{-n/2}\mathcal{L}^n(\epsilon') \subset \mathcal{L}^{n/2}(\epsilon')$ are “far enough” from each other (see Fig. 3.2). The “distance” between two Lagrangians above $\Omega_{\epsilon'_{n/2}}$ is actually measured by the *height*

$$Ht(\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell), \mathcal{L}^{n/2}(\epsilon')) \stackrel{\text{def}}{=} \inf_{x \in \Omega_{\epsilon'_{n/2}}} |dS^{\frac{3}{2}n}(x; \epsilon, \ell) - dS^{n/2}(x; \epsilon')|.$$

The overlap between “distant” leaves can be estimated through a nonstationary phase argument:

LEMMA 3.8. — *Assume that, for some $\delta'' < \delta'/2$, for some $\hbar > 0$ and some time $n \leq n_E(\hbar)$, the height*

$$Ht(\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell), \mathcal{L}^{n/2}(\epsilon')) \geq \hbar^{\frac{1-\delta''}{2}}.$$

Then, provided \hbar is small enough, the overlap

$$(3.52) \quad |\langle w^{n/2}(\epsilon'), v^{\frac{3}{2}n}(\epsilon, \ell) \rangle| \leq C \hbar^{N\delta''} \sqrt{J_{n/2}^u(\epsilon') J_{\frac{3}{2}n}^u(\epsilon, \ell)}.$$

The constant $C > 0$ is uniform with respect to ϵ', ϵ and the implicit parameters z, z', η_1, η'_1 .

Proof. — Let us call $s(x) = S^{\frac{3}{2}n}(x; \epsilon, \ell) - S^{n/2}(x; \epsilon')$ the phase function appearing in the integral (3.6). Notice that the assumption on the height means that $|ds(x)| \geq \hbar^{\frac{1-\delta''}{2}}$ for all x . We then expand the product $\bar{c}_{\hbar}^{n/2} b_{\hbar}^{\frac{3}{2}n}$ and keep only the first N terms:

$$\bar{c}_{\hbar}^{n/2}(x; \epsilon') b_{\hbar}^{\frac{3}{2}n}(x; \epsilon, \ell) = a_{\hbar}(x) + \text{Rem}_N(x), \quad a_{\hbar}(x) = \sum_{k=0}^{N-1} \hbar^k a_k(x).$$

From the estimates (3.44, 3.48), we control the sup-norm of the remainder:

$$\|\text{Rem}_N\|_{L^\infty} \leq C \hbar^{N\delta'/2} \sqrt{J_{n/2}^u(\epsilon') J_{\frac{3}{2}n}^u(\epsilon, \ell)}.$$

Through the Leibniz rule we control the derivatives of a_k :

$$\|a_k\|_{C^m} \leq C n^{m+3k} \sqrt{J_{n/2}^u(\epsilon') J_{\frac{3}{2}n}^u(\epsilon, \ell)} e^{\frac{\sigma}{2}(m+2k)\lambda_+}, \quad k \leq N-1, \quad m \leq 2(N-k).$$

For each $k < N$ and $m \leq 2(N-k)$, we have at our disposal the following nonstationary phase estimate [17, Section 7.7]:

$$\begin{aligned} \left| \int a_k(x) \exp\left(\frac{i}{\hbar}s(x)\right) dx \right| &\leq C \hbar^m \sum_{m' \leq m} \sup_x \left(\frac{|d^{m'} a_k(x)|}{|ds(x)|^{2m-m'}} \right) \\ &\leq C \hbar^{m\delta'' - k(1-\delta'/2)} \sqrt{J_{n/2}^u(\epsilon') J_{\frac{3}{2}n}^u(\epsilon, \ell)}. \end{aligned}$$

Here we used the assumption on $|ds(x)|$ and the fact that $\delta'' < \delta'/2$. By taking $m = N - k$ for each k and summing the estimate over k , we get:

$$\left| \int a_{\hbar}(x) \exp\left(\frac{i}{\hbar}s(x)\right) dx \right| \leq C \hbar^{N\delta''} \sqrt{J_{n/2}^u(\epsilon') J_{\frac{3}{2}n}^u(\epsilon, \ell)}.$$

Since $\delta'/2 > \delta''$, the remainder Rem_N yields a smaller contribution, which ends the proof. □

We now show that there is at most one Lagrangian leaf $\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell_o)$ which can be very close to $\mathcal{L}^{n/2}(\epsilon')$:

LEMMA 3.9. — Take as above $\delta'' < \delta'/2$, assume the diameter ε is much smaller than the injectivity radius, and for \hbar small enough take $n \leq \frac{(1-\delta'')|\log \hbar|}{\lambda_{\max}}$.

If there is some $\ell_o \in \{1, \dots, L\}$ such that the height $Ht(\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell_o), \mathcal{L}^{n/2}(\epsilon')) \leq \hbar^{\frac{1-\delta''}{2}}$, then for any $\ell \neq \ell_o$ we must have $Ht(\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell), \mathcal{L}^{n/2}(\epsilon')) > \hbar^{\frac{1-\delta''}{2}}$.

Proof. — Assume *ab absurdo* the existence of $\rho_o \in \mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell_o)$, $\rho \in \mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell)$ and $\rho'_1, \rho'_2 \in \mathcal{L}^{n/2}(\epsilon')$, such that the Riemannian distances $d(\rho_o, \rho'_1) \leq \hbar^{\frac{1-\delta''}{2}}$ and $d(\rho, \rho'_2) \leq \hbar^{\frac{1-\delta''}{2}}$. When applying the backwards flow for times $0 \leq t \leq \frac{n}{2}$, these points depart at most like

$$d(g^{-t}\rho_o, d^{-t}\rho'_1) \leq Ce^{t\lambda} \hbar^{\frac{1-\delta''}{2}} \leq C\hbar^{\delta'/4-\delta''/2},$$

$$d(g^{-t}\rho, d^{-t}\rho'_2) \leq Ce^{t\lambda} \hbar^{\frac{1-\delta''}{2}} \leq C\hbar^{\delta'/4-\delta''/2}.$$

Besides, on this time interval the points $g^{-t}\rho'_1, g^{-t}\rho'_2$ remain in the small Lagrangian piece $g^{-t}\mathcal{L}^{n/2}(\epsilon')$ of diameter $\leq \epsilon$, so that $d(g^{-t}\rho_o, g^{-t}\rho) \leq \epsilon$. Since ϵ has been chosen small, this contradicts the property that the points $g^{-t}\rho_o, g^{-t}\rho$ must depart at a distance $\geq D$ (see the discussion at the beginning of §3.5.3). □

If there exists a leaf ℓ_o such that $Ht(\mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell_o), \mathcal{L}^{n/2}(\epsilon')) \leq \hbar^{\frac{1-\delta''}{2}}$, there is a point $\rho_o \in \mathcal{L}^{\frac{3}{2}n}(\epsilon, \ell_o)$ such that $g^{-j}\rho_o$ stays at small distance from $\mathcal{L}^{n/2-j}(\epsilon')$ for all $j = 0, \dots, n/2-1$, and therefore satisfies $\pi g^{-j}\rho_o \in \Omega_{\epsilon'_{n/2-j}}$. This shows that the set A_{ℓ_o} contains the sequence $(\epsilon'_1 \dots \epsilon'_{n/2}) \stackrel{\text{def}}{=} \tilde{\epsilon}'$. The overlap corresponding to this leaf is bounded as in (3.51), and after replacing $J_{\frac{3}{2}n}^u(\epsilon, \ell_o)$ by $J_{\frac{3}{2}n}^u(\epsilon \tilde{\epsilon}')$ we obtain

$$(3.53) \quad | \langle w^{n/2}(\epsilon'), v^{\frac{3}{2}n}(\epsilon; \ell_o) \rangle | \leq C J_n^u(\epsilon') J_n^u(\epsilon)^{1/2}.$$

According to the above two Lemmas, all the remaining leaves are “far from” $\mathcal{L}^{n/2}(\epsilon')$, and their contributions to (3.49) sum up to

$$\sum_{\ell \neq \ell_o} \langle w^{n/2}(\epsilon'), v^{\frac{3}{2}n}(\epsilon; \ell) \rangle = \mathcal{O}(\hbar^{N\delta''-(d-1)/2}).$$

We take N large enough (say, $N\delta'' \gg 1$), such that this is negligible compared with (3.53). We finally get, whether such an ℓ_o exists or not:

$$| \langle U^{-n/2} v^n(z', \eta'_1, \epsilon'), U^{n/2} v^n(z, \eta_1, \epsilon) \rangle | \leq C J_n^u(\epsilon') J_n^u(\epsilon)^{1/2}.$$

To finish the proof of Lemma 3.2, there remains to integrate over the parameters η_1, η'_1 in (3.35). Since $\chi^{(n)}$ (resp. $\chi^{(4n)}$) is supported on an interval of length $\hbar^{1-\delta} e^{n\delta}$ (resp. $\hbar^{1-\delta} e^{4n\delta}$), the overlap of Lemma 3.2 finally satisfies

the following bound:

$$\begin{aligned}
 & |\langle U^{-n/2} P_{\epsilon'} \delta_{j'}^{(4n)}(z'), U^{n/2} P_{\epsilon} \delta_j^{(n)}(z) \rangle| \\
 & \leq C \hbar^{-(d+1)} e^{5\delta n} \hbar^{2-2\delta} J_n^u(\epsilon') J_n^u(\epsilon)^{1/2}.
 \end{aligned}$$

This is the estimate of Lemma 3.2, with $c = 2 + 5/\lambda_{\max}$. Proposition 3.1 and Theorem 2.7 follow. □

4. Subadditivity

The aim of this section is to prove Proposition 2.9. It is convenient here to use some notions of symbolic dynamics. Starting from our partition of unity $(P_k)_{k=1, \dots, K}$, we introduce a symbolic space $\Sigma = \{1, \dots, K\}^{\mathbb{N}}$. The shift σ acts on Σ by shifting a sequence $\epsilon = \epsilon_0 \epsilon_1 \dots$ to the left and deleting the first symbol. For $\epsilon = (\epsilon_0 \dots \epsilon_n)$, we denote $[\epsilon] \subset \Sigma$ the subset (n -cylinder) formed of sequences starting with the symbols $\epsilon_0 \dots \epsilon_n$ (throughout this section the integer n will generally differ from $n_E(\hbar)$).

To any normalized eigenfunction ψ_{\hbar} we can associate a probability measure μ_{\hbar}^{Σ} on Σ by letting, for any n -cylinder $[\epsilon]$,

$$\begin{aligned}
 \mu_{\hbar}^{\Sigma}([\epsilon]) & \stackrel{\text{def}}{=} \|P_{\epsilon_n} P_{\epsilon_{n-1}}(1) \dots P_{\epsilon_0}(n) \psi_{\hbar}\|^2 \\
 & = \|P_{\epsilon_n}(-n) P_{\epsilon_{n-1}}(-(n-1)) \dots P_{\epsilon_0} \psi_{\hbar}\|^2.
 \end{aligned}$$

If we denote $\bar{\epsilon} = (\epsilon_n \epsilon_{n-1} \dots \epsilon_0)$, this quantity is equal to $\|\tilde{P}_{\bar{\epsilon}}^* \psi_{\hbar}\|^2 = \|P_{\bar{\epsilon}}^* \psi_{\hbar}\|^2$ (see (2.23)). To ensure that this defines a probability measure on Σ , one needs to check the following compatibility condition

$$(4.1) \quad \mu_{\hbar}^{\Sigma}([\epsilon_0 \dots \epsilon_n]) = \sum_{\epsilon_{n+1}=1}^K \mu_{\hbar}^{\Sigma}([\epsilon_0 \dots \epsilon_n \epsilon_{n+1}])$$

for all n and all $\epsilon_0 \dots \epsilon_n$. This identity is obvious from (2.4).

4.1. Invariance until the Ehrenfest time

By the Egorov theorem, if μ is the weak- $*$ limit of the Wigner measures $W_{\psi_{\hbar}}$ on T^*M , then for every n and any fixed n -cylinder $[\epsilon] \subset \Sigma$ we have $\mu_{\hbar}^{\Sigma}([\epsilon]) \xrightarrow{\hbar \rightarrow 0} \mu(\{\bar{\epsilon}\})$, where $\{\bar{\epsilon}\}$ was defined in §2.2.7 as the function $P_{\epsilon_n}^2 (P_{\epsilon_{n-1}}^2 \circ g^1) \dots (P_{\epsilon_0}^2 \circ g^n)$ on T^*M . This means that the measures μ_{\hbar}^{Σ} converge to a measure μ_0^{Σ} defined by $\mu_0^{\Sigma}([\epsilon]) \stackrel{\text{def}}{=} \mu(\{\bar{\epsilon}\})$.

Since the ψ_{\hbar} are eigenfunctions, μ is localized on \mathcal{E} and is (g^t) -invariant (Prop. 1.1), so that μ_0^Σ is σ -invariant. For $\hbar > 0$ the measures μ_{\hbar}^Σ are not exactly σ -invariant; yet, we show below that μ_{\hbar}^Σ is *almost invariant* under the shift, until the Ehrenfest time.

For small $\gamma, \nu > 0$ we introduce the time $T_{\nu, \gamma, \hbar} \stackrel{\text{def}}{=} \frac{(1-\gamma)|\log \hbar|}{2(1+\nu)\lambda_{\max}}$.

PROPOSITION 4.1. — *For any given $n_o \in \mathbb{N}$, for any small enough \hbar and any $n \in \mathbb{N}$ such that $n + n_o \leq 2T_{\nu, \gamma, \hbar}$, for any cylinder $[\epsilon] = [\epsilon_0 \epsilon_1 \cdots \epsilon_{n_o}]$ of length n_o , one has*

$$\sum_{\epsilon_i, -n \leq i \leq -1} \mu_{\hbar}^\Sigma([\epsilon_{-n} \cdots \epsilon_{-1} \epsilon_0 \epsilon_1 \cdots \epsilon_{n_o}]) = \mu_{\hbar}^\Sigma([\epsilon_0 \epsilon_1 \cdots \epsilon_{n_o}]) + \mathcal{O}(\hbar^{\gamma/2}).$$

The implied constant is uniform with respect to n_o and n in the allowed interval. In other words, the measure μ_{\hbar}^Σ is almost σ -invariant:

$$\sigma_{\#}^n \mu_{\hbar}^\Sigma([\epsilon]) \stackrel{\text{def}}{=} \mu_{\hbar}^\Sigma(\sigma^{-n}[\epsilon]) = \mu_{\hbar}^\Sigma([\epsilon]) + \mathcal{O}(\hbar^{\gamma/2}).$$

Proof. — For simplicity we prove the result for $n_o = 0$; the argument can easily be adapted to any $n_o > 0$.

We use an estimate on the norm of commutators, proved in Lemma 5.2. If A is an operator on $L^2(M)$, remember that we denote $A(t) = U^{-t} A U^t$. According to Lemma 5.2, for any smooth observables a, b supported inside $\mathcal{E}^\nu = \mathcal{E}(1/2 - \nu, 1/2 + \nu)$, one has

$$(4.2) \quad \|\text{Op}_{\hbar}(a)(t), \text{Op}_{\hbar}(b)(-t)\|_{L^2(M)} = \mathcal{O}(\hbar^\gamma),$$

or equivalently

$$\|\text{Op}_{\hbar}(a)(2t), \text{Op}_{\hbar}(b)\|_{L^2(M)} = \mathcal{O}(\hbar^\gamma),$$

for any time $|t| \leq T_{\nu, \gamma, \hbar}$. This result will be applied to the observables $a = P_{\epsilon_0} f, b = P_{\epsilon_j} f$, where f is compactly supported in \mathcal{E}^ν and identically 1 near \mathcal{E} . According to Remark 2.4, inserting the cutoff f after each P_{ϵ_j} only modifies $\mu_{\hbar}^\Sigma([\epsilon])$ by an amount $\mathcal{O}(\hbar^\infty)$. In the following, we will omit to indicate these insertions and the $\mathcal{O}(\hbar^\infty)$ errors.

To prove Proposition 4.1, we first write

$$\begin{aligned} & \sum_{\epsilon_i, -n \leq i \leq -1} \mu_{\hbar}^\Sigma([\epsilon_{-n} \epsilon_{-(n-1)} \cdots \epsilon_0]) \\ &= \sum_{\epsilon_i, -n \leq i \leq -1} \|P_{\epsilon_0} P_{\epsilon_{-1}}(1) \cdots P_{\epsilon_{-n}}(n) \psi_{\hbar}\|^2 \\ &= \sum \langle P_{\epsilon_{-1}}(1) P_{\epsilon_0}^2 P_{\epsilon_{-1}}(1) \tilde{P}_{[\epsilon_{-2} \cdots \epsilon_{-n}]}^*(2) \psi_{\hbar}, \tilde{P}_{[\epsilon_{-2} \cdots \epsilon_{-n}]}^*(2) \psi_{\hbar} \rangle \end{aligned}$$

$$\begin{aligned}
 &= \sum \langle P_{\epsilon_0}^2 P_{\epsilon_{-1}} (1)^2 \tilde{P}_{[\epsilon_{-2} \dots \epsilon_{-n}]}^* (2) \psi_{\hbar}, \tilde{P}_{[\epsilon_{-2} \dots \epsilon_{-n}]}^* (2) \psi_{\hbar} \rangle \\
 &\quad + \mathcal{O}(\hbar^\gamma) \left[\sum_{\epsilon_i, -n \leq i \leq -2} \|\tilde{P}_{[\epsilon_{-2} \dots \epsilon_{-n}]}^* (2) \psi_{\hbar}\|^2 \right] \\
 &= \sum_{\epsilon_i, -n \leq i \leq -2} \langle P_{\epsilon_0}^2 \tilde{P}_{[\epsilon_{-2} \dots \epsilon_{-n}]}^* (2) \psi_{\hbar}, \tilde{P}_{[\epsilon_{-2} \dots \epsilon_{-n}]}^* (2) \psi_{\hbar} \rangle + \mathcal{O}(\hbar^\gamma).
 \end{aligned}$$

We have used the identities

$$\sum_{\epsilon_{-1}} P_{\epsilon_{-1}} (1)^2 = I \text{ and } \sum_{\epsilon_{-n}, \dots, \epsilon_{-2}} \|\tilde{P}_{[\epsilon_{-2} \dots \epsilon_{-n}]}^* \psi_{\hbar}\|^2 = 1.$$

We repeat the procedure:

$$\begin{aligned}
 &\sum_{\epsilon_i, -n \leq i \leq -2} \langle P_{\epsilon_0}^2 \tilde{P}_{[\epsilon_{-2} \dots \epsilon_{-n}]}^* (2) \psi_{\hbar}, \tilde{P}_{[\epsilon_{-2} \dots \epsilon_{-n}]}^* (2) \psi_{\hbar} \rangle \\
 &= \sum \langle P_{\epsilon_{-2}} (2) P_{\epsilon_0}^2 P_{\epsilon_{-2}} (2) \tilde{P}_{[\epsilon_{-3} \dots \epsilon_{-n}]}^* (3) \psi_{\hbar}, \tilde{P}_{[\epsilon_{-3} \dots \epsilon_{-n}]}^* (3) \psi_{\hbar} \rangle \\
 &= \sum \langle P_{\epsilon_0}^2 P_{\epsilon_{-2}} (2)^2 \tilde{P}_{[\epsilon_{-3} \dots \epsilon_{-n}]}^* (3) \psi_{\hbar}, \tilde{P}_{[\epsilon_{-3} \dots \epsilon_{-n}]}^* (3) \psi_{\hbar} \rangle \\
 &\quad + \mathcal{O}(\hbar^\gamma) \left[\sum_{\epsilon_i, -n \leq i \leq -3} \|\tilde{P}_{[\epsilon_{-3} \dots \epsilon_{-n}]}^* (3) \psi_{\hbar}\|^2 \right] \\
 &= \sum_{\epsilon_i, -n \leq i \leq -3} \langle P_{\epsilon_0}^2 \tilde{P}_{[\epsilon_{-3} \dots \epsilon_{-n}]}^* (3) \psi_{\hbar}, \tilde{P}_{[\epsilon_{-3} \dots \epsilon_{-n}]}^* (3) \psi_{\hbar} \rangle + \mathcal{O}(\hbar^\gamma).
 \end{aligned}$$

Iterating this procedure n times we obtain

$$\sum_{\epsilon_i, -n \leq i \leq -1} \mu_{\hbar}^{\Sigma}([\epsilon_{-n} \epsilon_{-(n-1)} \dots \epsilon_0]) = \langle P_{\epsilon_0}^2 \psi_{\hbar}, \psi_{\hbar} \rangle + n \mathcal{O}(\hbar^\gamma),$$

which proves the Proposition for $n_0 = 0$, since $n = \mathcal{O}(|\log \hbar|)$. The proof for any fixed $n_0 > 0$ is identical. □

4.2. Proof of Proposition 2.9

For ψ_{\hbar} an eigenstate of the Laplacian, the entropy $h_n(\psi_{\hbar})$ introduced in (2.12) can be expressed in terms of the measure μ_{\hbar}^{Σ} :

$$\begin{aligned}
 (4.3) \quad h_n(\psi_{\hbar}) &= - \sum_{|\epsilon|=n} \|\tilde{P}_{\epsilon}^* \psi_{\hbar}\|^2 \log \|\tilde{P}_{\epsilon}^* \psi_{\hbar}\|^2 \\
 &= - \sum_{|\epsilon|=n} \mu_{\hbar}^{\Sigma}([\bar{\epsilon}]) \log \mu_{\hbar}^{\Sigma}([\bar{\epsilon}]) \\
 &= - \sum_{|\epsilon|=n} \mu_{\hbar}^{\Sigma}([\epsilon]) \log \mu_{\hbar}^{\Sigma}([\epsilon]) \stackrel{\text{def}}{=} h_n(\mu_{\hbar}^{\Sigma}).
 \end{aligned}$$

In ergodic theory, the last term is called the entropy of the measure μ_{\hbar}^{Σ} with respect to the partition of Σ into n -cylinders. Before using the results of the previous section, we choose the parameters ν, γ appearing in Proposition 4.1 such that $\nu = \gamma = \delta'/2$, where δ' is the small parameter in Proposition 2.9. This ensures that the time $2T_{\nu, \gamma, \hbar} \geq n_E(\hbar)$ (see (2.10)).

We then have, for any n_o and n such that $n + n_o \leq T_{\nu, \gamma, \hbar}$,

$$(4.4) \quad \begin{aligned} h_{n_o+n}(\mu_{\hbar}^{\Sigma}) &\leq h_{n-1}(\mu_{\hbar}^{\Sigma}) + h_{n_o}(\sigma_{\sharp}^n \mu_{\hbar}^{\Sigma}) \\ &= h_{n-1}(\mu_{\hbar}^{\Sigma}) + h_{n_o}(\mu_{\hbar}^{\Sigma}) + \mathcal{O}_{n_o}(\hbar^{\delta'/4}). \end{aligned}$$

The notation \mathcal{O}_{n_o} means that the last term is bounded by $C_{n_o} \hbar^{\delta'/4}$, with a constant C_{n_o} depending on n_o . The first inequality is a general property of the entropy, due to the concavity of the logarithm. The second equality comes from the almost invariance of μ_{\hbar}^{Σ} (Proposition 4.1) and the continuity of the function $x \mapsto -x \log x$. The pressure for ψ_{\hbar} (see (2.13)) also involves sums of the type

$$\sum_{\epsilon = \epsilon_0 \dots \epsilon_{n_o+n}} \mu_{\hbar}^{\Sigma}([\epsilon]) \log J_{n_o+n}^u(\epsilon) \stackrel{\text{def}}{=} \mu_{\hbar}^{\Sigma}(\log J_{n_o+n}^u).$$

Using the factorization (2.16) of the Jacobian, this sum can be split into

$$(4.5) \quad \begin{aligned} \mu_{\hbar}^{\Sigma}(\log J_{n_o+n}^u) &= \mu_{\hbar}^{\Sigma}(\log J_{n-1}^u) + \sigma_{\sharp}^{n-1} \mu_{\hbar}^{\Sigma}(\log J_1^u) + \sigma_{\sharp}^n \mu_{\hbar}^{\Sigma}(\log J_{n_o}^u) \\ &= \mu_{\hbar}^{\Sigma}(\log J_{n-1}^u) + \mu_{\hbar}^{\Sigma}(\log J_1^u) + \mu_{\hbar}^{\Sigma}(\log J_{n_o}^u) + \mathcal{O}_{n_o}(\hbar^{\delta'/4}). \end{aligned}$$

We used once more the quasi-invariance of μ_{\hbar}^{Σ} to get the second equality. Combining the inequalities (4.4, 4.5) with (4.3), we obtain the Proposition 2.9 with the constant

$$R = 3 \max_{\rho \in \mathcal{E}^{\epsilon}} |\log J_1^u(\rho)|. \quad \square$$

5. Some results of pseudodifferential calculus

5.1. Pseudodifferential calculus on a manifold

In this section we present the standard Weyl quantization of observables defined on the cotangent of the compact d -dimensional manifold M (see for instance [13]). The manifold can be equipped with an atlas $\{f_{\ell}, V_{\ell}\}$, such that the V_{ℓ} form an open cover of M , and for each ℓ , f_{ℓ} is a diffeomorphism from V_{ℓ} to a bounded open set $W_{\ell} \subset \mathbb{R}^d$. Each f_{ℓ} induces a pullback

$f_\ell^* : C^\infty(W_\ell) \rightarrow C^\infty(V_\ell)$. We denote by \tilde{f}_ℓ the induced canonical map between T^*V_ℓ and T^*W_ℓ :

$$(x, \xi) \in T^*V_\ell \mapsto \tilde{f}_\ell(x, \xi) = (f_\ell(x), (Df_\ell(x))^{-1})^T \xi \in T^*W_\ell,$$

(A^T is the transposed of A) and by $\tilde{f}_\ell^* : C^\infty(T^*W_\ell) \rightarrow C^\infty(T^*V_\ell)$ the corresponding pull-back. One then chooses a smooth partition of unity on M adapted to the charts $\{V_\ell\}$, namely a set of functions $\phi_\ell \in C_c^\infty(V_\ell)$ such that $\sum_\ell \phi_\ell = 1$ on M .

Any observable $a \in C^\infty(T^*M)$ can now be split into $a = \sum_j a_\ell$, with $a_\ell = \phi_\ell a$, each term being pushed to $\tilde{a}_\ell = (\tilde{f}_\ell^{-1})^* a_\ell \in C^\infty(T^*W_\ell)$. If a belongs to a nice class of functions (possibly depending on \hbar), for instance the space of symbols

$$(5.1) \quad a \in S^{m,k} = S^k(\langle \xi \rangle^m) \\ \stackrel{\text{def}}{=} \left\{ a = a_\hbar \in C^\infty(T^*M), |\partial_x^\alpha \partial_\xi^\beta a| \leq C_{\alpha,\beta} \hbar^{-k} \langle \xi \rangle^{m-|\beta|} \right\},$$

then Weyl-quantization associates to each \tilde{a}_ℓ a pseudodifferential operator on $\mathcal{S}(\mathbb{R}^d)$:

$$(5.2) \quad \forall u \in \mathcal{S}(\mathbb{R}^d), \text{Op}_\hbar^w(\tilde{a}_\ell)u(x) = \frac{1}{(2\pi\hbar)^d} \int e^{\frac{i}{\hbar}\langle x-y, \xi \rangle} \tilde{a}_\ell\left(\frac{x+y}{2}, \xi; \hbar\right) u(y) dy d\xi.$$

To pull this pseudodifferential operator back on $C^\infty(V_\ell)$, one takes a smooth cutoff $\psi_\ell \in C_c^\infty(V_\ell)$ such that $\psi_\ell(x) = 1$ close to $\text{supp } \phi_\ell$. The quantization of $a \in S^{m,k}$ is finally defined as follows:

$$(5.3) \quad \forall u \in C^\infty(M), \text{Op}_\hbar(a)u = \sum_\ell \psi_\ell \times f_\ell^* \circ \text{Op}_\hbar^w(\tilde{a}_\ell) \circ (f_\ell^{-1})^*(\psi_\ell \times u).$$

The space of pseudodifferential operators image of $S^{m,k}$ through this quantization is denoted by $\Psi^{m,k}(M)$. The quantization obviously depends on the cutoffs ϕ_ℓ, ψ_ℓ . However, this dependence only appears at second order in \hbar , and the principal symbol map $\sigma : \Psi^{m,k}(M) \rightarrow S^{m,k}/S^{m,k-1}$ is intrinsically defined. All microlocal properties of pseudodifferential operators on \mathbb{R}^d are carried over to $\Psi^{m,k}(M)$. The Laplacian $-\hbar^2 \Delta$ belongs to $\Psi^{2,0}(M)$, with principal symbol $\sigma(-\hbar^2 \Delta) = |\xi|_x^2$.

We actually need to consider symbols more general than (5.1). Following [10], for any $0 \leq \epsilon < 1/2$ we introduce the symbol class

$$(5.4) \quad S_\epsilon^{m,k} \stackrel{\text{def}}{=} \left\{ a \in C^\infty(T^*M), |\partial_x^\alpha \partial_\xi^\beta a| \leq C_{\alpha,\beta} \hbar^{-k-\epsilon|\alpha+\beta|} \langle \xi \rangle^{m-|\beta|} \right\}.$$

The induced functions \tilde{a}_ℓ will then belong to the corresponding class on T^*W_ℓ , for which we can use the results of [10]. For instance, the quantization of any $a \in S_\epsilon^{0,0}$ leads to a bounded operator on $L^2(M)$ (the norm being bounded uniformly in \hbar).

5.2. Egorov theorem up to logarithmic times

We need analogous estimates to Bouzouina-Robert’s [7] concerning the quantum-classical equivalence for long times. Our setting is more general, since we are interested in observables on T^*M for an arbitrary manifold M . On the other hand, we will only be interested in the first order term in the Egorov theorem, whereas [7] described the complete asymptotic expansion in powers of \hbar .

The evolution is given by the propagator U^t on $L^2(M)$, which quantizes the flow g^t on T^*M . We will consider smooth observables $a \in C_c^\infty(T^*M)$ supported in a thin neighbourhood of the energy layer \mathcal{E} , say inside the energy strip $\mathcal{E}^\nu = \mathcal{E}([1/2 - \nu, 1/2 + \nu])$ for some small $\nu > 0$. This strip is invariant through the flow, so the evolved observable $a_t = a \circ g^t$ will remain supported inside \mathcal{E}^ν . If λ_{\max} is the maximal expansion rate of the flow on \mathcal{E} (see the definition in Theorem 1.2), then by homogeneity the maximal expansion rate inside \mathcal{E}^ν is $\sqrt{1 + 2\nu}\lambda_{\max}$. If we let $\lambda_\nu \stackrel{\text{def}}{=} (1 + \nu)\lambda_{\max}$, the successive derivatives of the flow on \mathcal{E}^ν are controlled as follows:

$$(5.5) \quad \forall t \in \mathbb{R}, \forall \rho \in \mathcal{E}^\nu, \|\partial_\rho^\alpha g^t(\rho)\| \leq C_\alpha e^{\lambda_\nu |\alpha t|}.$$

Obviously, the derivatives of the evolved observable also satisfy

$$(5.6) \quad \forall t \in \mathbb{R}, \forall \rho \in \mathcal{E}^\nu, \|\partial^\alpha a_t(\rho)\| \leq C_{\alpha,\alpha} e^{\lambda_\nu |\alpha t|}.$$

For times of the order of $|\log \hbar|$, each derivative is bounded by some power of \hbar^{-1} . More precisely, for any $\gamma \in (0, 1]$ and any $\hbar \in (0, 1/2)$, we call $T_{\nu,\gamma,\hbar}$ the following time:

$$(5.7) \quad T_{\nu,\gamma,\hbar} = \frac{(1 - \gamma)|\log \hbar|}{2\lambda_\nu} = \frac{(1 - \gamma)|\log \hbar|}{2(1 + \nu)\lambda_{\max}}.$$

Starting from a smooth observable $a = a_0$, the bounds (5.6) show that the family of function $\{a_t = a \circ g^t : |t| \leq T_{\nu,\gamma,\hbar}\}$ remains in the symbol class $S_\epsilon^{-\infty,0}$, with $\epsilon = \frac{1-\gamma}{2}$. Furthermore, any quasi-norm is uniformly bounded within the family. To prove a Egorov estimate, we start as usual from the

identity

$$(5.8) \quad U^{-t} \text{Op}_{\hbar}(a)U^t - \text{Op}_{\hbar}(a \circ g^t) = \int_0^t ds U^{-s} (\text{Diff } a_{t-s}) U^s,$$

$$\text{with } \text{Diff } a_t \stackrel{\text{def}}{=} \frac{i}{\hbar} [-\hbar^2 \Delta, \text{Op}_{\hbar}(a_t)] - \text{Op}_{\hbar}(\{H, a_t\}).$$

Since $-\hbar^2 \Delta$ belongs to $\Psi^{2,0} \subset \Psi_{\epsilon}^{2,0}$ and $\text{Op}_{\hbar}(a_t) \in \Psi_{\epsilon}^{-\infty,0}$ for times $|t| \leq T_{\nu,\gamma,\hbar}$, the semiclassical calculus of [10, Prop. 7.7] (performed locally on each chart V_j) shows that $\text{Diff } a_t \in \Psi_{\epsilon}^{-\infty,-\alpha}$, with $\alpha = 1 - \epsilon = \frac{1+\gamma}{2}$. From the Calderon-Vaillancourt theorem on $\Psi_{\epsilon}^{-\infty,-\alpha}$ [10, Thm. 7.11], we extract a constant $C_a > 0$ such that, for any small enough $\hbar > 0$ and any time $|t| \leq T_{\nu,\gamma,\hbar}$,

$$\|\text{Diff } a_t\| \leq C_a \hbar^{\alpha} = C_a \hbar^{\frac{1+\gamma}{2}}.$$

We can finally combine the above estimate in (5.8) and use the unitarity of U^t (Duhamel’s principle) to obtain the following Egorov estimate.

PROPOSITION 5.1. — *Fix $\nu, \gamma \in (0, 1]$. Let a be a smooth, \hbar -independent observable supported in \mathcal{E}^{ν} . Then, there is a constant C_a such that, for any time $|t| \leq T_{\nu,\gamma,\hbar}$, one has*

$$(5.9) \quad \|U^{-t} \text{Op}_{\hbar}(a)U^t - \text{Op}_{\hbar}(a \circ g^t)\| \leq C_a |t| \hbar^{\frac{1+\gamma}{2}}.$$

Let us now consider two observables $a, b \in C_c^{\infty}(\mathcal{E}^{\nu})$, evolve one in the future, the other in the past. The calculus in $S_{\epsilon}^{-\infty,0}$ (with again $\epsilon = \frac{1-\gamma}{2}$) shows that, for any time $|t| \leq T_{\nu,\gamma,\hbar}$, one has

$$[\text{Op}_{\hbar}(a \circ g^t), \text{Op}_{\hbar}(b \circ g^{-t})] \in S_{\epsilon}^{-\infty,-\gamma}.$$

Together with the above Egorov estimate and the Calderon-Vaillancourt theorem on $\Psi_{\epsilon}^{-\infty,-\gamma}$, this shows the following

LEMMA 5.2. — *Fix $\nu, \gamma \in (0, 1]$. Let $a, b \in C_c^{\infty}(\mathcal{E}^{\nu})$ be independent of \hbar . Then there is a constant $C > 0$ such that, for small \hbar and any time $|t| \leq T_{\nu,\gamma,\hbar}$,*

$$\|[U^{-t} \text{Op}_{\hbar}(a)U^t, U^t \text{Op}_{\hbar}(b)U^{-t}]\| \leq C \hbar^{\gamma}.$$

5.3. Cutoff in a thin energy strip

As explained in §2.2.3, we need an energy cutoff $\chi^{(0)}$ localizing in the energy strip of width $\sim \hbar^{\epsilon}$ around \mathcal{E} , with $\epsilon \in [0, 1)$ arbitrary close to 1. As a result, the m -th derivatives of χ transversally to \mathcal{E} will grow like $\hbar^{-m\epsilon}$. The symbol classes (5.4) introduced in the previous sections do not include such

functions if $\epsilon \geq 1/2$. Yet, because the fluctuations occur close to \mathcal{E} and only transversally, it is possible to work with a “second-microlocal” pseudodifferential calculus which includes such fast-varying, anisotropic symbols. We summarize here the treatment of this problem performed in [28, Section 4].

5.3.1. Local behavior of the anisotropic symbols

For any $\epsilon \in [0, 1)$, we introduce a class of symbols $S_{\mathcal{E}, \epsilon}^{m, k}$, made of functions $a = a_{\hbar}$ satisfying the following properties:

- for any family of smooth vectors fields V_1, \dots, V_{l_1} tangent to \mathcal{E} , and of smooth vector fields W_1, \dots, W_{l_2} , one has in each energy strip $\mathcal{E}^\nu = \mathcal{E}([1/2 - \nu, 1/2 + \nu])$:

$$\sup_{\rho \in \mathcal{E}^\nu} |V_1 \cdots V_{l_1} W_1 \cdots W_{l_2} a(\rho)| = \mathcal{O}(\hbar^{-k - \epsilon l_2}).$$

- away from \mathcal{E} , we have $|\partial_x^\alpha \partial_\xi^\beta a(\rho)| = \mathcal{O}(\hbar^{-k} \langle \xi \rangle^{m - |\beta|})$.

Notice that $S^{m, k} \subset S_{\mathcal{E}, \epsilon'}^{m, k} \subset S_{\mathcal{E}, \epsilon}^{m, k}$ if $1 > \epsilon > \epsilon' \geq 0$.

To quantize this class of symbols, we cover a certain neighbourhood \mathcal{E}^ν of \mathcal{E} by a family of bounded open sets \mathcal{V}_j , such that for each j , \mathcal{V}_j is mapped by a canonical diffeomorphism κ_j to a bounded open set $\mathcal{W}_j \subset T^*\mathbb{R}^d$, with $(0, 0) \in \mathcal{W}_j$. We will denote by (x, ξ) the local coordinates on $\mathcal{V}_j \subset T^*M$, and (y, η) the image coordinates on \mathcal{W}_j . The canonical map κ_j is chosen such that $H \circ \kappa_j^{-1} = \eta_1 + 1/2$. In particular, the image of $\mathcal{E} \cap \mathcal{V}_j$ is a piece of the hyperplane $\{\eta_1 = 0\}$.

We consider a smooth cutoff function ϕ supported inside \mathcal{E}^ν , with $\phi \equiv 1$ in $\mathcal{E}^{\nu/2}$, and a smooth partition of unity (φ_j) such that $1 = \sum_j \varphi_j$ on $\cup_j \mathcal{V}_j$, and $\text{supp } \varphi_j \Subset \mathcal{V}_j$. For any symbol $a \in S_{\mathcal{E}, \epsilon}^{m, k}$, the function $a(1 - \phi)$ is supported outside $\mathcal{E}^{\nu/2}$, and it belongs to the standard class $S^{m, k}$ of (5.1). On the other hand, for each index j the function

$$a_j \stackrel{\text{def}}{=} (a\phi\varphi_j) \circ \kappa_j^{-1}$$

is compactly supported inside $\mathcal{W}_j \subset T^*\mathbb{R}^d$.

That function can be Weyl-quantized as in (5.2). Although $a_j(y, \eta)$ can oscillate at a rate $\hbar^{-\epsilon}$ along the coordinate η_1 near $\{\eta_1 = 0\}$, for $a, b \in S_{\mathcal{E}, \epsilon}^{m, k}$ the product $\text{Op}_\hbar^w(a_j) \text{Op}_\hbar^w(b_j)$ is still of the form $\text{Op}_\hbar^w(c_j)$, where the function $c_j(y, \eta)$ is given by the Moyal product $a_j \sharp b_j$ and satisfies an asymptotic expansion in powers of $\hbar^{1-\epsilon}$ and \hbar .

Mimicking the proof of the Calderon-Vaillancourt theorem in [10, Thm. 7.11], we use the isometry (in $L^2(\mathbb{R}^d)$) between $\text{Op}_\hbar^w(A)$ and $\text{Op}_1^w(A \circ T_\hbar)$,

where the rescaling

$$T_{\hbar}(y, \eta) = (y_1 \hbar^{\frac{1-\epsilon}{2}}, y' \hbar^{1/2}; \eta_1 \hbar^{\frac{1+\epsilon}{2}}, \eta' \hbar^{1/2})$$

ensures that the derivatives of $a_j \circ T_{\hbar}$ are uniformly bounded in \hbar . As a consequence we get the following

PROPOSITION 5.3. — *There exist N_d and $C > 0$ such that the following bound holds. For any symbol $a \in S_{\mathcal{E}, \epsilon}^{m, k}$ and any j , the operator $\text{Op}_{\hbar}^w(a_j)$ acts continuously on $L^2(\mathbb{R}^d)$, and its norm is bounded as follows:*

$$\|\text{Op}_{\hbar}^w(a_j)\| \leq \|a_j\|_{L^\infty} + C \sum_{1 \leq |\alpha| + |\beta| \leq N_d} \hbar^{\frac{1}{2}(|\alpha'| + |\beta'| + (1-\epsilon)\alpha_1 + (1+\epsilon)\beta_1)} \times \|\partial_y^\alpha \partial_\eta^\beta a_j\|_{L^\infty}.$$

5.3.2. Global quantization of the anisotropic symbols

We now glue together the various pieces of $a \in S_{\mathcal{E}, \epsilon}^{m, k}$ to define its global quantization. First of all, since $a(1 - \phi)$ belongs to the standard class $S^{m, k}$ of (5.1), we can quantize it as in §5.1.

Then, for each index j we select a Fourier integral operator $U_{\kappa_j} : L^2(\pi(\mathcal{V}_j)) \rightarrow L^2(\pi(\mathcal{W}_j))$, elliptic near $\text{supp } \varphi_j \times \kappa_j(\text{supp } \varphi_j) \subset \mathcal{V}_j \times \mathcal{W}_j$, and associated with the diffeomorphism κ_j (an explicit expression is given in §3.2.1). Since a_j describes the symbol a in the coordinates (y, η) , it makes sense to pull $\text{Op}_{\hbar}^w(a_j)$ back to the original coordinates (x, ξ) using U_{κ_j} . The quantization of the global symbol $a \in S_{\mathcal{E}, \epsilon}^{m, k}$ is then defined as follows:

$$(5.10) \quad \text{Op}_{\mathcal{E}, \hbar}(a) \stackrel{\text{def}}{=} \text{Op}_{\hbar}(a(1 - \phi)) + \sum_j U_{\kappa_j}^* \text{Op}_{\hbar}^w(a_j) U_{\kappa_j}.$$

The Fourier integral operators (U_{κ_j}) can and will be chosen such that $\text{Op}_{\mathcal{E}, \hbar}(1) = \text{Id} + \mathcal{O}_{L^2 \rightarrow L^2}(\hbar^\infty)$. The operators $\text{Op}_{\mathcal{E}, \hbar}(a)$ make up a space $\Psi_{\mathcal{E}, \epsilon}^{m, k}$ of pseudodifferential operators on M . The quantization $\text{Op}_{\mathcal{E}, \hbar}$ depends on the choice of the cutoffs ϕ, φ_j , the diffeomorphisms κ_j and the associated FIOs (U_{κ_j}) . It is equal to the quantization Op_{\hbar} for symbols a supported outside the energy strip \mathcal{E}^ν ; otherwise, it differs from Op_{\hbar} by higher-order terms.

The space $\Psi_{\mathcal{E}, \epsilon}^{-\infty, k}$ is invariant under conjugation by FIOs which preserve the energy layer \mathcal{E} . We will apply that property to the propagator $U = e^{i\hbar\Delta/2}$, which quantizes the flow g^1 . One actually has a Egorov property

$$U^{-1} \text{Op}_{\mathcal{E}, \hbar}(a) U = \text{Op}_{\mathcal{E}, \hbar}(b), \text{ with } b - a \circ g \in S_{\mathcal{E}, \epsilon}^{-\infty, k-1+\epsilon}.$$

One is naturally lead to the definition of an \hbar -dependent essential support of a symbol $a_{\hbar} \in S_{\mathcal{E}, \epsilon}^{m, k}$ (we will only consider the finite part of the

essential support, the infinite part at $|\xi| = \infty$ being irrelevant for our purposes). A family of sets $(V_{\hbar} \subset T^*M)_{\hbar \rightarrow 0}$ does not intersect $\text{ess - supp } a_{\hbar}$ iff there exists $\chi_{\hbar} \in S_{\mathcal{E}, \epsilon}^{-\infty, 0}$, with $\chi_{\hbar} \geq 1$ on V_{\hbar} , such that $\chi_{\hbar} a_{\hbar} \in S_{\mathcal{E}, \epsilon}^{-\infty, -\infty}$. The essential support of a_{\hbar} is also the wavefront set of its quantization, $WF_{\hbar}(\text{Op}_{\mathcal{E}, \hbar}(a_{\hbar}))$.

The above Egorov property can be iterated to all orders, showing that the wavefront set of an operator $A \in \Psi_{\mathcal{E}, \epsilon}^{-\infty, k}$ is transported classically:

$$(5.11) \quad WF_{\hbar}(U^{-1}AU) = g^{-1}(WF_{\hbar}(A)).$$

5.4. Properties of the energy cutoffs

Take some small $\delta > 0$ and $C_{\delta} > 0$ as in §2.2.3, and define $\epsilon = 1 - \delta$. One can easily check that the cutoffs $\chi^{(n)}$ defined in (2.6), with $n \leq C_{\delta} |\log \hbar|$, all belong to the symbol class $S_{\mathcal{E}, \epsilon}^{-\infty, 0}$. From the above results, their quantizations $\text{Op}(\chi^{(n)}) = \text{Op}_{\mathcal{E}, \epsilon}(\chi^{(n)})$ are continuous operators on $L^2(M)$, of norms

$$(5.12) \quad \|\text{Op}(\chi^{(n)})\| = 1 + \mathcal{O}(\hbar^{\delta/2}),$$

with an implied constant independent of n . We want to check that these cutoffs have little influence on an eigenstate ψ_{\hbar} satisfying (2.2). For this, we invoke the ellipticity of $(-\hbar^2 \Delta - 1) \in \Psi_{\mathcal{E}, \epsilon}^{2, 0} \subset \Psi_{\mathcal{E}, \epsilon}^{2, 0}$ away from \mathcal{E} . Using [28, Prop. 4.1], one can adapt the standard division lemma to show the following

PROPOSITION 5.4. — *For $\hbar > 0$ small enough and any $n \in \mathbb{N}$, $0 \leq n \leq C |\log \hbar|$, there exists $A_{\hbar}^{(n)} \in \Psi_{\mathcal{E}, \epsilon}^{-2, \epsilon}$ and $R_{\hbar}^{(n)} \in \Psi_{\mathcal{E}, \epsilon}^{-\infty, -\infty}$ such that*

$$\text{Op}_{\mathcal{E}, \hbar}(1 - \chi^{(n)}) = A_{\hbar}^{(n)}(-\hbar^2 \Delta - 1) + R_{\hbar}^{(n)}.$$

As a result, for any eigenstate $\psi_{\hbar} = -\hbar^2 \Delta \psi_{\hbar}$, one has

$$\|\psi_{\hbar} - \text{Op}_{\mathcal{E}, \hbar}(\chi^{(n)})\psi_{\hbar}\| = \mathcal{O}(\hbar^{\infty})\|\psi_{\hbar}\|.$$

The implied constant is uniform with respect to n .

This result contains in particular the estimate (2.8).

We end this section by proving some properties of the cutoffs $\chi^{(n)}$. The general idea is that an eigenstate ψ_{\hbar} is localized in an energy strip of width \hbar , so that inserting cutoffs $\chi^{(n)}$ in expressions of the type $\text{Op}(a)\psi_{\hbar}$ has a negligible effect.

LEMMA 5.5. — *The following estimates are uniform for $\hbar > 0$ small enough and $0 \leq n \leq C_\delta |\log \hbar|$:*

$$\begin{aligned} & \| (1 - \text{Op}(\chi^{(n+1)})) U \text{Op}(\chi^{(n)}) \| = \mathcal{O}(\hbar^\infty), \\ \forall k = 1, \dots, K, & \| (1 - \text{Op}(\chi^{(n+1)})) U P_k \text{Op}(\chi^{(n)}) \| = \mathcal{O}(\hbar^\infty). \end{aligned}$$

Here P_k is any element of the partition of unity (2.4).

Proof. — For the symbols $\chi^{(n)}$ the essential support (which has been defined above in a rather indirect way) coincides with the support. The first statement of the Lemma uses the classical transport of the wavefront set (5.11), applied to $\text{Op}(\chi^{(n)})$. Since $\chi^{(n)}$ is invariant through the geodesic flow, $U \text{Op}(\chi^{(n)}) U^{-1}$ has the same wavefront set as $\text{Op}(\chi^{(n)})$. From the definition (2.6), the support of $(1 - \chi^{(n+1)})$ is at a distance $\geq C \hbar^\epsilon$ from the support of $\chi^{(n)}$. The calculus on $S_{\mathcal{E}, \epsilon}^{0,0}$ then implies that the product $(1 - \text{Op}(\chi^{(n+1)})) \text{Op}(\chi^{(n)})$ is in $\Psi_{\mathcal{E}, \epsilon}^{-\infty, -\infty}$.

The second statement is a consequence of the first: the calculus on $\Psi_{\mathcal{E}, \epsilon}^{0,0}$, which contains the cutoffs $\text{Op}(\chi^{(n)})$ and the multiplication operators P_k , shows that the wavefront set of $P_k \text{Op}(\chi^{(n)})$ is contained in that of $\text{Op}(\chi^{(n)})$. □

We draw from this Lemma two properties which we use in the text (see (2.5) for the definition of P_ϵ).

COROLLARY 5.6. — *For any sequence ϵ of length $n \leq C_\delta |\log \hbar|$, one has*

$$\| (1 - \text{Op}(\chi^{(n)})) P_\epsilon \text{Op}(\chi^{(0)}) \| = \mathcal{O}(\hbar^\infty).$$

For any two sequence ϵ, ϵ' of length $n \leq C_\delta |\log \hbar|/4$, one has

$$\| (1 - \text{Op}(\chi^{(4n)})) P_{\epsilon'}^* U^n P_\epsilon \text{Op}(\chi^{(n)}) \| = \mathcal{O}(\hbar^\infty).$$

6. The entropic uncertainty principle: an application of complex interpolation

In this section we prove the weighted entropic uncertainty principle, namely theorem 2.1, by adapting the original proof of [25].

We consider a complex Hilbert space $(\mathcal{H}, \langle \cdot, \cdot \rangle)$, and denote the associated norm by $\|\psi\| = \sqrt{\langle \psi, \psi \rangle}$. The same notation $\|\cdot\|$ will also be used for the operator norm on $\mathcal{L}(\mathcal{H})$.

Let $(\alpha_k)_{k=1, \dots, \mathcal{N}}$ be a family of positive numbers. We consider the weighted l_p -norms on $\mathcal{H}^{\mathcal{N}} \ni \Psi = (\Psi_1, \dots, \Psi_{\mathcal{N}})$:

$$(6.1) \quad \|\Psi\|_p^{(\alpha)} \stackrel{\text{def}}{=} \left(\sum_{k=1}^{\mathcal{N}} \alpha_j^{p-2} \|\Psi_k\|^p \right)^{1/p}, \quad 1 \leq p < \infty,$$

and

$$\|\Psi\|_\infty^{(\alpha)} \stackrel{\text{def}}{=} \max_k (\alpha_k \|\Psi_k\|).$$

For $p = 2$, this norm does not depend on (α_k) and coincides with the Hilbert norm deriving from the scalar product

$$\langle \Psi, \Phi \rangle_{\mathcal{H}^{\mathcal{N}}} = \sum_k \langle \Psi_k, \Phi_k \rangle_{\mathcal{H}}.$$

If $\Psi \in \mathcal{H}^{\mathcal{N}}$ has Hilbert norm unity, we define its entropy as

$$h(\Psi) = - \sum_{k=1}^{\mathcal{N}} \|\Psi_k\|^2 \log \|\Psi_k\|^2,$$

and its pressure with respect to the weights (α_k) is defined by

$$(6.2) \quad p_\alpha(\Psi) = - \sum_{k=1}^{\mathcal{N}} \|\Psi_k\|^2 \log \|\Psi_k\|^2 - \sum_{k=1}^{\mathcal{N}} \|\Psi_k\|^2 \log \alpha_k^2.$$

This is the derivative of $\|\Psi\|_p^{(\alpha)}$ with respect to p , evaluated at $p = 2$.

Similarly, let $(\beta_j)_{j=1, \dots, \mathcal{M}}$ be a family of weights. They induce the following $l_p^{(\beta)}$ -norms on $\mathcal{H}^{\mathcal{M}} \ni \Phi = (\Phi_1, \dots, \Phi_{\mathcal{M}})$:

$$\|\Phi\|_p^{(\beta)} \stackrel{\text{def}}{=} \left(\sum_{j=1}^{\mathcal{M}} \beta_j^{p-2} \|\Phi_j\|^p \right)^{1/p}, \quad 1 \leq p < \infty,$$

and

$$\|\Phi\|_\infty^{(\beta)} \stackrel{\text{def}}{=} \max_j (\beta_j \|\Phi_j\|).$$

We can define the entropy of a normalized vector $\Phi \in \mathcal{H}^{\mathcal{M}}$, and its pressure $p_\beta(\Phi)$ with respect to the weights $(\beta_j)_{j=1, \dots, \mathcal{M}}$. The standard $l_p - l_q$ duality [12, Thm.IV.8.1] reads as follows in the present context:

PROPOSITION 6.1. — *For any $1 < p, q < \infty$ such that $\frac{1}{p} + \frac{1}{q} = 1$, then*

$$(6.3) \quad \sup_{\|\Psi\|_p^{(\alpha)}=1} |\langle \Lambda, \Psi \rangle| = \|\Lambda\|_q^{(\alpha)}.$$

6.1. Complex interpolation

A bounded operator $T : \mathcal{H}^{\mathcal{N}} \rightarrow \mathcal{H}^{\mathcal{M}}$ can be represented by a $\mathcal{M} \times \mathcal{N}$ matrix (T_{jk}) of bounded operators on \mathcal{H} . For $1 \leq p, q \leq \infty$ we denote by $\|T\|_{p,q}^{(\alpha,\beta)}$ the norm of T from $l_p^{(\alpha)}(\mathcal{H}^{\mathcal{N}})$ to $l_q^{(\beta)}(\mathcal{H}^{\mathcal{M}})$. We assume that $\|T\|_{2,2} = 1$, which implies in particular that $\|T_{jk}\| \leq 1$ for all k, j .

Example 6.2. — Suppose we have two partitions of unity $(\pi_k)_{k=1}^{\mathcal{N}}$ and $(\tau_j)_{j=1}^{\mathcal{M}}$ on \mathcal{H} , that is, two families of operators such that

$$(6.4) \quad \sum_{k=1}^{\mathcal{N}} \pi_k \pi_k^* = \text{Id}, \quad \sum_{j=1}^{\mathcal{M}} \tau_j \tau_j^* = \text{Id}.$$

The main example we have in mind is the case where \mathcal{U} is a unitary operator on \mathcal{H} and $T_{jk} \stackrel{\text{def}}{=} \tau_j^* \mathcal{U} \pi_k$.

Let O be a bounded operator on \mathcal{H} , and let $\vartheta \geq 0$. We will be interested in the action of T on the cone

$$\mathcal{C}(O, \vartheta) = \{ \Psi \in \mathcal{H}^{\mathcal{N}}, \|O\Psi_k - \Psi_k\| \leq \vartheta \|\Psi\|_2 \text{ for all } k = 1, \dots, \mathcal{N} \} \subset \mathcal{H}^{\mathcal{N}}.$$

Notice that the cone $\mathcal{C}(O, \vartheta)$ coincides with $\mathcal{H}^{\mathcal{N}}$ in the special case $O = \text{Id}$, $\vartheta = 0$, which is already an interesting case.

We introduce the positive number

$$c_O(T) = \max_{j,k} \alpha_k \beta_j \|T_{jk} O\|_{\mathcal{L}(\mathcal{H})},$$

and also $A = \max_k \alpha_k$, $B = \max_j \beta_j$. The following theorem extends the result of [25].

THEOREM 6.3. — *For all $\Psi \in \mathcal{C}(O, \vartheta)$ such that $\|\Psi\|_2 = 1$ and $\|T\Psi\|_2 = 1$, we have*

$$p_\beta(T\Psi) + p_\alpha(\Psi) \geq -2 \log (c_O(T) + \mathcal{N}AB\vartheta).$$

The proof of this theorem follows the standard proof of the Riesz-Thorin theorem [12, sec.VI.10]. In particular, one uses the following convexity property of complex analytic functions.

LEMMA 6.4 (3-circle theorem). — *Let $f(z)$ be analytic and bounded in the strip $\{0 < x < 1\}$, and continuous on the closed strip. Then, the function $\log M(x) = \log \sup_{y \in \mathbb{R}} |f(x + iy)|$ is convex in the interval $0 \leq x \leq 1$.*

We will define an appropriate analytic function in the unit strip. Let $\Psi \in \mathcal{C}(O, \vartheta)$ with $\|\Psi\|_2 = 1$. Fix $t \in [0, 1]$, close to 0, and let

$$\tilde{\Psi} = \frac{\Psi}{\|\Psi\|_{\frac{2}{1+t}}^{(\alpha)}}.$$

From the definition of the norm and Hölder’s inequality, we have

$$\|\Psi\|_{\frac{2}{1+t}}^{(\alpha)} \geq A^{-t}.$$

Consider any state $\Phi \in \mathcal{H}^{\mathcal{M}}$ such that $\|\Phi\|_{\frac{2}{1+t}}^{(\beta)} \leq 1$. For each $z = x + iy$ in the strip $\{0 \leq x \leq 1\}$, we define

$$a(z) = \frac{1+z}{1+t},$$

and the states

$$\begin{aligned} \tilde{\Psi}(z) &= \left(\tilde{\Psi}(z)_k = \tilde{\Psi}_k \|\tilde{\Psi}_k\|^{a(z)-1} \alpha_k^{a(z)-1} \right)_{k=1, \dots, \mathcal{N}}, \\ \Phi(z) &= \left(\Phi(z)_j = \Phi_j \|\Phi_j\|^{a(z)-1} \beta_j^{a(z)-1} \right)_{j=1, \dots, \mathcal{M}}. \end{aligned}$$

By construction, we have

$$\forall z = x + iy, \|\tilde{\Psi}(z)\|_{\frac{2}{1+x}}^{(\alpha)} = 1 \text{ and } \|\Phi(z)\|_{\frac{2}{1+x}}^{(\beta)} \leq 1.$$

In particular, for any $y \in \mathbb{R}$ we have

$$(6.5) \quad \|\tilde{\Psi}(iy)\|_2 = 1 \text{ and } \|\Phi(iy)\|_2 \leq 1 \implies \left| \langle T\tilde{\Psi}(iy), \Phi(iy) \rangle \right| \leq \|T\|_{2,2}.$$

Similarly, for any $y \in \mathbb{R}$,

$$\|\Phi(1+iy)\|_1^{(\beta)} \leq 1 \implies \left| \langle T\tilde{\Psi}(1+iy), \Phi(1+iy) \rangle \right| \leq \|T\tilde{\Psi}(1+iy)\|_{\infty}^{(\beta)}.$$

We decompose the right hand side by inserting the operator O :

$$\begin{aligned} \|T\tilde{\Psi}(1+iy)\|_{\infty}^{(\beta)} &= \max_j \beta_j \left\| \sum_k T_{jk} \tilde{\Psi}(1+iy)_k \right\| \\ &\leq \max_j \beta_j \left\| \sum_k T_{jk} O\tilde{\Psi}(1+iy)_k \right\| + \max_j \beta_j \left\| \sum_k T_{jk} (\text{Id} - O)\tilde{\Psi}(1+iy)_k \right\|. \end{aligned}$$

The first term on the right hand side is bounded above by $c_O(T)\|\tilde{\Psi}(1+iy)\|_1^{(\alpha)} = c_O(T)$. For the second term, we remark that

$$\|\tilde{\Psi}(1+iy)_k\| = |\alpha_k|^{\frac{1-t}{1+t}} \|\tilde{\Psi}_k\|_{\frac{2}{1+t}}^{\frac{2}{1+t}} = \frac{|\alpha_k|^{\frac{1-t}{1+t}} \|\Psi_k\|_{\frac{2}{1+t}}^{\frac{2}{1+t}}}{\left(\|\Psi\|_{\frac{2}{1+t}}^{(\alpha)}\right)^{\frac{2}{1+t}}}.$$

On the one hand, $\|\Psi_k\| \leq \|\Psi\|_2 \leq 1$ and $|\alpha_k|^{\frac{1-t}{1+t}} \leq A^{\frac{1-t}{1+t}}$. On the other hand we have already stated that $\|\Psi\|_{\frac{2}{1+t}}^{(\alpha)} \geq A^{-t}$. Putting these bounds together and using the fact that $\Psi \in \mathcal{C}(O, \vartheta)$, we get

$$\forall k = 1, \dots, \mathcal{N}, \quad \|(\text{Id} - O)\tilde{\Psi}(1+iy)_k\| \leq A\vartheta.$$

Summing over k and using $\|T_{jk}\| \leq 1$, we find

$$\max_j \beta_j \left\| \sum_{k=1}^{\mathcal{N}} T_{jk} (\text{Id} - O)\tilde{\Psi}(1+iy)_k \right\| \leq \mathcal{N}AB\vartheta.$$

We have proved that for all $y \in \mathbb{R}$,

$$(6.6) \quad \left| \langle T\tilde{\Psi}(1 + iy), \Phi(1 + iy) \rangle \right| \leq c_O(T) + \mathcal{N}AB\vartheta.$$

The function $z \mapsto \langle T\tilde{\Psi}(z), \Phi(z) \rangle$ is bounded and analytic in the strip $\{0 \leq x \leq 1\}$: this is the function to which we apply the 3-circle theorem (Lemma 6.4). Taking into account (6.5, 6.6), we obtain for any $x \in [0, 1]$, $y \in \mathbb{R}$,

$$\begin{aligned} \log \left| \langle T\tilde{\Psi}(x + iy), \Phi(x + iy) \rangle \right| &\leq (1 - x) \log \|T\|_{2,2} + x \log(c_O(T) + \mathcal{N}AB\vartheta) \\ &\leq x \log (c_O(T) + \mathcal{N}AB\vartheta). \end{aligned}$$

The last inequality is due to our assumption $\|T\|_{2,2} = 1$. In particular, taking $x + iy = t$, and exponentiating, we get

$$\left| \langle T\tilde{\Psi}, \Phi \rangle \right| \leq (c_O(T) + \mathcal{N}AB\vartheta)^t.$$

Taking the supremum over $\left\{ \Phi \in \mathcal{H}^M, \|\Phi\|_{\frac{2}{1+t}}^{(\beta)} \leq 1 \right\}$ and using the $l_{\frac{2}{1+t}}^{(\beta)}$ - $l_{\frac{2}{1-t}}^{(\beta)}$ duality (Prop. 6.1), we obtain

$$\|T\tilde{\Psi}\|_{\frac{2}{1-t}}^{(\beta)} \leq (c_O(T) + \mathcal{N}AB\vartheta)^t,$$

and by homogeneity

$$(6.7) \quad \|T\Psi\|_{\frac{2}{1-t}}^{(\beta)} \leq (c_O(T) + \mathcal{N}AB\vartheta)^t \|\Psi\|_{\frac{2}{1+t}}^{(\alpha)}.$$

We may now take the limit $t \rightarrow 0$ in this inequality. Using the assumption $\|\Psi\|_2 = 1$, we notice that

$$\begin{aligned} \log \|\Psi\|_{\frac{2}{1+t}}^{(\alpha)} &\sim \frac{1+t}{2} \log \left(\sum_k \|\Psi_k\|^2 \exp\{-t \log \|\Psi_k\|^2 - t \log \alpha_k^2\} \right) \\ &\sim \frac{t}{2} p_\alpha(\Psi). \end{aligned}$$

Similarly, $\log \|T\Psi\|_{\frac{2}{1-t}}^{(\beta)} \sim -\frac{t}{2} p_\beta(T\Psi)$. Therefore, in the limit $t \rightarrow 0$, (6.7) implies Theorem 6.3. □

6.2. Specialization to particular operators T and states Ψ

We now come back to the case of Example 6.2.

LEMMA 6.5. — *Let $\mathcal{U} : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded operator. Using the two partitions of Example 6.2, we construct the operator $T : \mathcal{H}^{\mathcal{N}} \rightarrow \mathcal{H}^{\mathcal{M}}$ through its components $T_{jk} = \tau_j^* \mathcal{U} \pi_k$. Then the two following norms are equal:*

$$\|T\|_{2,2} = \|\mathcal{U}\|_{\mathcal{L}(\mathcal{H})}.$$

Proof. — The operator T may be described as follows. Consider a line and column vectors of operators on \mathcal{H} :

$$L \stackrel{\text{def}}{=} (\pi_1, \dots, \pi_N), \text{ respectively } C = \begin{pmatrix} \tau_1^* \\ \vdots \\ \tau_M^* \end{pmatrix}.$$

We can write $T = CUL$. We insert this formula in the identity

$$\|T\|_{2,2}^2 = \|T^*T\|_{\mathcal{L}(\mathcal{H}^{\mathcal{N}})} = \|L^* \mathcal{U}^* C^* CUL\|_{\mathcal{L}(\mathcal{H}^{\mathcal{N}})}$$

Using the resolution of identity of the τ_j , we notice that $C^*C = \text{Id}_{\mathcal{H}}$, so that the above norm reads

$$\|L^* \mathcal{U}^* \mathcal{U} L\|_{\mathcal{L}(\mathcal{H}^{\mathcal{N}})}.$$

Then, using the resolution of identity of the π_k , we get

$$\begin{aligned} \|(\mathcal{U}L)^*(\mathcal{U}L)\|_{\mathcal{L}(\mathcal{H}^{\mathcal{N}})} &= \|(\mathcal{U}L)(\mathcal{U}L)^*\|_{\mathcal{L}(\mathcal{H})} = \|(\mathcal{U}L)L^* \mathcal{U}^*\|_{\mathcal{L}(\mathcal{H})} \\ &= \|\mathcal{U} \mathcal{U}^*\|_{\mathcal{L}(\mathcal{H})}. \end{aligned} \quad \square$$

Therefore, if \mathcal{U} is contracting (resp. $\|\mathcal{U}\|_{\mathcal{L}(\mathcal{H})} = 1$) one has $\|T\|_{2,2} \leq 1$ (resp. $\|T\|_{2,2} = 1$).

We also specialize the vector $\Psi \in \mathcal{H}^{\mathcal{N}}$ by taking $\Psi_k = \pi_k^* \psi$ for some normalized $\psi \in \mathcal{H}$. From the resolution of identity on the π_k , we check that $\|\Psi\|_2 = \|\psi\|$, and also $(T\Psi)_j = \tau_j^* \mathcal{U} \psi$. Thus, if $\|\mathcal{U} \psi\| = 1$, the second resolution of identity induces $\|T\Psi\|_2 = \|\mathcal{U} \psi\| = 1$. With this choice for T and Ψ , Theorem 6.3 reads as follows:

THEOREM 6.6. — *We consider the setting of Example 6.2. Let \mathcal{U} be an isometry on \mathcal{H} . Define $c_O^{(\alpha,\beta)}(U) \stackrel{\text{def}}{=} \sup_{j,k} \alpha_k \beta_j \|\tau_j^* \mathcal{U} \pi_k O\|_{\mathcal{L}(\mathcal{H})}$.*

Then, for any normalized $\psi \in \mathcal{H}$ satisfying

$$\forall k = 1, \dots, \mathcal{N}, \quad \|(\text{Id} - O)\pi_k^* \psi\| \leq \vartheta,$$

and defining the pressures as in (6.2), we have

$$p_\beta((\tau_j^* \mathcal{U} \psi)_{j=1, \dots, \mathcal{M}}) + p_\alpha((\pi_k^* \psi)_{k=1, \dots, \mathcal{N}}) \geq -2 \log(c_O^{(\alpha,\beta)}(U) + \mathcal{N} A B \vartheta).$$

This theorem implies Theorem 2.1, if we take the same partition $\pi = \tau$ (in particular $\mathcal{N} = \mathcal{M}$), and if we remark that the pressures $p_\alpha((\pi_k^* \psi)_{k=1, \dots, \mathcal{N}})$ and $p_\beta((\pi_j^* \mathcal{U} \psi)_{j=1, \dots, \mathcal{N}})$ are the same as the quantities $p_{\pi, \alpha}(\psi)$, $p_{\pi, \beta}(U\psi)$ appearing in the theorem.

BIBLIOGRAPHY

- [1] N. ANANTHARAMAN, H. KOCH & S. NONNENMACHER, “Entropy of eigenfunctions”, to appear in the Proceedings of the International Congress on Mathematical Physics - Rio de Janeiro, August 6 - 11, 2006.
- [2] N. ANANTHARAMAN, “Entropy and the localization of eigenfunctions”, to appear in *Ann. of Math.*, 2008.
- [3] N. ANANTHARAMAN & S. NONNENMACHER, “Entropy of semiclassical measures of the Walsh-quantized baker’s map”, *Ann. Henri Poincaré* **8** (2007), no. 1, p. 37-74.
- [4] P. H. BÉRARD, “On the wave equation on a compact Riemannian manifold without conjugate points”, *Math. Z.* **155** (1977), no. 3, p. 249-276.
- [5] M. V. BERRY, “Regular and irregular semiclassical wavefunctions”, *J. Phys. A* **10** (1977), no. 12, p. 2083-2091.
- [6] O. BOHIGAS, “Random matrix theories and chaotic dynamics”, in *Chaos et physique quantique (Les Houches, 1989)*, North-Holland, Amsterdam, 1991, p. 87-199.
- [7] A. BOUZOUINA & D. ROBERT, “Uniform semiclassical estimates for the propagation of quantum observables”, *Duke Math. J.* **111** (2002), no. 2, p. 223-252.
- [8] Y. COLIN DE VERDIÈRE, “Ergodicité et fonctions propres du laplacien”, *Comm. Math. Phys.* **102** (1985), no. 3, p. 497-502.
- [9] Y. COLIN DE VERDIÈRE & B. PARISSÉ, “Équilibre instable en régime semi-classique. I. Concentration microlocale”, *Comm. Partial Differential Equations* **19** (1994), no. 9-10, p. 1535-1563.
- [10] M. DIMASSI & J. SJÖSTRAND, *Spectral asymptotics in the semi-classical limit*, London Mathematical Society Lecture Note Series, vol. 268, Cambridge University Press, Cambridge, 1999.
- [11] H. DONNELLY, “Quantum unique ergodicity”, *Proc. Amer. Math. Soc.* **131** (2003), no. 9, p. 2945-2951 (electronic).
- [12] N. DUNFORD & J. T. SCHWARTZ, *Linear operators. I. General theory*, With the assistance of W. G. Bade and R. G. Bartle. Pure and Applied Mathematics, Vol. 7, Interscience Publishers, Inc., New York, 1958.
- [13] L. EVANS & M. WORSKI, “Lectures on semiclassical analysis” (version 0.2), <http://math.berkeley.edu/~zworski>.
- [14] F. FAURE & S. NONNENMACHER, “On the maximal scarring for quantum cat map eigenstates”, *Comm. Math. Phys.* **245** (2004), no. 1, p. 201-214.
- [15] F. FAURE, S. NONNENMACHER & S. DE BIÈVRE, “Scarred eigenstates for quantum cat maps of minimal periods”, *Comm. Math. Phys.* **239** (2003), no. 3, p. 449-492.
- [16] J. HADAMARD, *Le problème de Cauchy et les équations aux dérivées partielles linéaires hyperboliques*, Hermann, Paris, 1932.
- [17] L. HÖRMANDER, *The analysis of linear partial differential operators. I*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 256, Springer-Verlag, Berlin, 1983, Distribution theory and Fourier analysis.

- [18] A. KATOK & B. HASSELBLATT, *Introduction to the modern theory of dynamical systems*, Encyclopedia of Mathematics and its Applications, vol. 54, Cambridge University Press, Cambridge, 1995, With a supplementary chapter by Katok and Leonardo Mendoza.
- [19] D. KELMER, “Arithmetic quantum unique ergodicity for symplectic linear maps of the multidimensional torus”, to appear in *Ann. of Math.*, [math-ph/0510079](#).
- [20] ———, “Scarring on invariant manifolds for perturbed quantized hyperbolic toral automorphisms”, preprint, [arXiv:math-ph/0607033](#).
- [21] W. KLINGENBERG, “Riemannian manifolds with geodesic flow of Anosov type”, *Ann. of Math. (2)* **99** (1974), p. 1-13.
- [22] K. KRAUS, “Complementary observables and uncertainty relations”, *Phys. Rev. D* (3) **35** (1987), no. 10, p. 3070-3075.
- [23] F. LEDRAPPIER & L.-S. YOUNG, “The metric entropy of diffeomorphisms. I. Characterization of measures satisfying Pesin’s entropy formula”, *Ann. of Math. (2)* **122** (1985), no. 3, p. 509-539.
- [24] E. LINDENSTRAUSS, “Invariant measures and arithmetic quantum unique ergodicity”, *Ann. of Math. (2)* **163** (2006), no. 1, p. 165-219.
- [25] H. MAASSEN & J. B. M. UFFINK, “Generalized entropic uncertainty relations”, *Phys. Rev. Lett.* **60** (1988), no. 12, p. 1103-1106.
- [26] S. NONNENMACHER & M. ZWORSKI, “Quantum decay rates in chaotic scattering”, preprint, [arXiv:0706.3242](#), 2007.
- [27] Z. RUDNICK & P. SARNAK, “The behaviour of eigenstates of arithmetic hyperbolic manifolds”, *Comm. Math. Phys.* **161** (1994), no. 1, p. 195-213.
- [28] J. SJÖSTRAND & M. ZWORSKI, “Asymptotic distribution of resonances for convex obstacles”, *Acta Math.* **183** (1999), no. 2, p. 191-253.
- [29] A. I. ŠNIREL’MAN, “Ergodic properties of eigenfunctions”, *Uspehi Mat. Nauk* **29** (1974), no. 6(180), p. 181-182.
- [30] A. VOROS, “Semiclassical ergodicity of quantum eigenstates in the Wigner representation”, in *Stochastic behavior in classical and quantum Hamiltonian systems (Volta Memorial Conf., Como, 1977)*, Lecture Notes in Phys., vol. 93, Springer, Berlin, 1979, p. 326-333.
- [31] S. A. WOLPERT, “The modulus of continuity for $\Gamma_0(m)\backslash\mathbb{H}$ semi-classical limits”, *Comm. Math. Phys.* **216** (2001), no. 2, p. 313-323.
- [32] S. ZELDITCH, “Uniform distribution of eigenfunctions on compact hyperbolic surfaces”, *Duke Math. J.* **55** (1987), no. 4, p. 919-941.

Manuscrit reçu le 4 octobre 2006,
 accepté le 31 mai 2007.

Nalini ANANTHARAMAN
 École Normale Supérieure
 Unité de Mathématiques Pures et Appliquées
 6, allée d’Italie
 69364 LYON Cedex 07 (France)
nanantha@umpa.ens-lyon.fr
 Stéphane NONNENMACHER
 CEA/DSM/PhT
 Service de Physique Théorique
 Unité de recherche associé CNRS
 CEA/Saclay
 91191 Gif-sur-Yvette (France)
snonnenmacher@cea.fr