


Les cahiers du laboratoire Leibniz



A generalization of the Pentomino Exclusion
Problem : the Delta-dislocation in graphs.

Sylvain Gravier, Julien Moncel et Charles Payan.

Laboratoire Leibniz-IMAG, 46 av. Félix Viallet, 38000 GRENOBLE, France -
ISSN : 1298-020X

n° 72
Jan. 2003

Site internet : <http://www-leibniz.imag.fr/LesCahiers/>

A generalization of the Pentomino Exclusion Problem : the Δ -dislocation in graphs.

Sylvain Gravier*, Julien Moncel*, Charles Payan *

January 10, 2003

Abstract

In this paper we investigate the Pentomino Exclusion Problem, due to Golomb. We solve this problem on the $5 \times n$ grid and we give some lower and upper bounds for the $k \times n$ grid for all k, n .

A generalization of this in graphs leads to a new combinatorial problem, the Δ -dislocation problem : find the minimum number of vertices to be removed from the graph so as the remaining connected components have cardinality at most Δ . We investigate the algorithmic aspects of the Δ -dislocation problem : we first prove the problem is NP-Complete, then we give a sublinear algorithm which solves the problem on a restricted class of graphs which includes the $k \times n$ grid graphs when k is not apart of the input.

1 The Pentomino Exclusion Problem

1.1 Introduction

A *polyomino* is a pattern formed by the connection of a specified number of equal-sized squares along common edges (see [5]). A *pentomino* is a polyomino composed of 5 squares.

Golomb [5] proposed the following *Pentomino Exclusion Problem*, denoted $PEP_{k \times n}$: *Find the minimum number of unit squares to be placed on a $k \times n$ chessboard so as to exclude all pentominos.*

For convenience, we will denote this number $\kappa_{k \times n}$.

Bosch [1, 2] proposed a Linear Integer Programming approach to compute

*CNRS - UJF, GeoD research Group - laboratoire Leibniz, 46, avenue Félix Viallet, 38031 Grenoble Cedex (France)

$\kappa_{n \times n}$, and determined this number when $n \leq 12$ [1]. In [6], the authors determined $\kappa_{k \times n}$ for all n and $k \leq 4$ (see Theorem 1) and gave some results about the infinite case (\mathbb{Z}^2). Using the asymptotical results in [6], we will here determine $\kappa_{5 \times n}$ for all n and give some upper and lower bounds of this number for all k and all n .

Theorem 1

$$\kappa_{k \times n} = \begin{cases} \lfloor \frac{n}{5} \rfloor & \text{if } k = 1 \\ 2 \lfloor \frac{n}{3} \rfloor & \text{if } k = 2 \\ n & \text{if } k = 3 \text{ and } n \geq 2 \\ \lceil \frac{3n}{2} \rceil - 1 & \text{if } k = 4, n \geq 4 \end{cases} \quad \square$$

1.2 Case of the grid $5 \times n$

In this section, we investigate the problem $PEP_{5 \times n}$.

We denote by $G_{k,n}$ the $k \times n$ grid. For given k and n , C_1, \dots, C_n (respectively R_1, \dots, R_k) denote the *columns* (resp. the *rows*) of $G_{k,n}$. The squares of $G_{k,n}$ are denoted by $s_{i,j}$ where $\{s_{i,j}\} = R_i \cap C_j$.

Lemma 1 $\kappa_{5 \times n} \geq 2n - 2$.

Proof:

The proof works by induction on n .

The cases $n \leq 2$ are obvious.

Assume now that $n \geq 3$. Let S be a solution of $PEP_{5 \times n}$.

If $|S \cap C_1| \geq 2$, then by the induction hypothesis applied on $\cup_{i=2 \dots n} C_i$ we obtain $|S| \geq 2 + 2(n-1) - 2 = 2n - 2$.

Assume now, $|S \cap C_1| = 1$.

Let $j \geq 2$ the smallest integer such that $|S \cap C_j| \neq 2$.

If $|S \cap C_j| \geq 3$, then $|S \cap (C_1 \cup C_2 \cup \dots \cup C_j)| \geq 2j$, and by the induction hypothesis applied on $\cup_{i=j+1, \dots, n} C_i$ we have $|S \cap (C_{j+1} \cup C_{j+2} \cup \dots \cup C_n)| \geq 2(n-j) - 2$, so that $|S| \geq 2n - 2$.

Assume now that $|S \cap C_j| = 1$

If $j = 2$, then it is easy to see that $S \cap C_1 = \{s_{3,1}\}$ and $S \cap C_2 = \{s_{3,2}\}$ (see Figure 1). So we obtain $|S \cap C_3| \geq 4$, hence $|S| \geq 1 + 1 + 4 + 2(n-3) - 2 = 2n - 2$.

If $j = n$, then $|S| = 1 + 2(n-2) + 1 = 2n - 2$.

C_1	C_2	C_3

Figure 1: The case $|S \cap C_1| = |S \cap C_2| = 1$

If $2 < j < n$, then we have to study the position of the element in $S \cap C_j$.

If $S \cap C_j \in \{s_{1,j}; s_{5,j}\}$, then $|S \cap C_{j-1}| \geq 4$, which is a contradiction with the definition of j .

If $S \cap C_j \in \{s_{2,j}; s_{4,j}\}$, then since $|S \cap C_{j-1}| = 2$, it is easy to see that it implies that $|S \cap C_{j+1}| \geq 4$. Then by induction hypothesis applied on $\cup_{i=j+2 \dots n} C_i$, we obtain $|S| \geq 1 + 2(j-2) + 1 + 4 + 2(n-j-1) - 2 = 2n-2$.

Thus, we may assume that $S \cap C_j = \{s_{3,j}\}$.


We have $|S \cap (R_1 \cup R_2) \cap (C_{j-2} \cup C_{j-1} \cup C_j)| \geq 2$ and $|S \cap (R_4 \cup R_5) \cap (C_{j-2} \cup C_{j-1} \cup C_j)| \geq 2$. Since $|C_{j-1} \cap S| = 2$, we get also $|C_{j-2} \cap S| = 2$, which implies that $j-2 > 1$ and $(C_{j-2} \cup C_{j-1}) \cap R_3 = \emptyset$.

The squares $s_{2,j-1}, s_{4,j-1}$ must belong to S , otherwise the set $\{s_{1,j}, s_{2,j}, s_{2,j-1}, s_{3,j-1}, s_{3,j-2}, s_{4,j-1}, s_{4,j}, s_{5,j}\}$ contains a pentomino. From now, an easy exhaustive exploration of cases (see Figure 2) shows that $|S \cap C_{j-3}| \geq 3$, which is a contradiction with the definition of j . \square

Theorem 2

$$\kappa_{5 \times n} = \begin{cases} 2n-1 & \text{if } n \in \{1 \dots 4\} \\ 2n-2 & \text{if } n \geq 5 \end{cases}$$

Proof: The case $n \in \{1 \dots 4\}$ holds by Theorem 1. To conclude we exhibit solutions for $n \geq 5$ (see Figure 3) satisfying $|S| = 2n-2$. \square



At least 3 more squares in the first column !!

Figure 2: The case $S \cap C_k = \{s_{3,k}\}$.

1.3 General Bounds

We need the following Theorem due to S. Gravier and C. Payan [6] :

Theorem 3 *The density¹ of an optimal solution of $PEP_{\mathbb{Z}^2}$ is $\frac{3}{7}$. \square*

Theorem 4 *For all $k, n \geq 5$ we have :*

$$\frac{3}{7}kn + \frac{3}{7}(k+n) - \left(\left\lceil \frac{4n}{5} \right\rceil + \left\lceil \frac{4k}{5} \right\rceil\right) - \frac{4}{7} \leq \kappa_{k \times n} \leq \frac{3}{7}(k-4)n + 22\left\lceil \frac{n}{14} \right\rceil$$

Proof:

For the upper bound, we know, by Theorem 3, that the optimal density of $PEP_{\mathbb{Z}^2}$ is $\frac{3}{7}$. In Figure 4, we give such a solution \mathcal{T} . This solution is periodic and can be described as a translation of a given row. Here we will use a more precise description. For a periodic solution \mathcal{S} of $PEP_{\mathbb{Z}^2}$, each row R_i can be encoded by the *upper-word* $u_1 \dots u_t$ (respectively the *down-word* $d_1 \dots d_t$) where the integer u_s (resp. d_s) means that the corresponding square belongs to a polyomino of size u_s (resp. d_s) in $\cup_{j \geq i} R_j \setminus \mathcal{S}$ (resp. $\cup_{j \leq i} R_j \setminus \mathcal{S}$). Therefore, a square belongs to \mathcal{S} if and only if it is labelled 0.

For instance, the solution \mathcal{T} in Figure 4 admits the upper-word $UW = 01040403303301$ and the down-word $DW = 04010103303304$.

¹The density of a solution S in a finite set T is given by $d(S, T) = \frac{|S \cap T|}{|T|}$. Roughly speaking, the density of a solution S in \mathbb{Z}^2 is the limit (when there exists one) of $d(S, T)$ when T ‘grows’. For a more precise definition we refer the reader to [6].

Now, from \mathcal{T} , we will construct a solution of $\text{PEP}_{k \times n}$ for any $k, n \geq 5$, as follows : first replace the rows $\cup_{j \leq 2} R_j$ by two rows corresponding to a solution of $\text{PEP}_{2 \times \mathbb{Z}\mathbb{Z}}$ where the row R_2 has a down-word $DW2 = 10401040101040$ (see Figure 5). The density of $R_1 \cup R_2$ is $\frac{11}{14}$ which is better than the density of \mathcal{T} ($\frac{12}{14}$).

Observe that this gives a solution \mathcal{T}' (see Figure 5) of $\text{PEP}_{\mathbb{Z}\mathbb{Z}^+ \times \mathbb{Z}\mathbb{Z}}$. Indeed, each coordinate of the word $UW + DW2$ is smaller than 5.

Moreover, since for the solution \mathcal{T} each polymino in $\mathbb{Z}\mathbb{Z}^2 \setminus \mathcal{T}$ has cardinality 4, then every row R_j of \mathcal{T}' with $j \geq 3$ has an down-word for which each coordinate is smaller or equal to the ‘corresponding’ coordinate of DW . In fact, in our solution of $\text{PEP}_{\mathbb{Z}\mathbb{Z}^+ \times \mathbb{Z}\mathbb{Z}}$, the down-word of R_j is again DW whenever $j \geq 4$ and the down-word of R_3 is 01010103303301.

Let $k \geq 5$ be an integer and let DW' be the down-word of \mathcal{T}' associated to R_{k-2} .

Replace the rows $\cup_{j \geq k-2} R_j$ by two rows corresponding to the solution of $\text{PEP}_{2 \times \mathbb{Z}\mathbb{Z}}$ where the row R_{k-1} has the upper-word $UW2 = 10401040101040$ (see Figure 5).

This construction gives a solution S' of $\text{PEP}_{k \times \mathbb{Z}\mathbb{Z}}$, since each coordinate of the word $DW' + UW2$ is smaller than 5.

Now, to obtain a solution S^* for $\text{PEP}_{k \times n}$ it is enough to take n consecutive columns C_1, \dots, C_n of S' for which the density of $(R_3 \cup \dots \cup R_{k-2}) \cap (C_1 \cup \dots \cup C_n)$ is $\leq \frac{3}{7}$.

Finally, S^* has cardinality :

- In the first (last) 2 rows, less or equal to $11 \lceil \frac{n}{14} \rceil$.
- In $(R_3 \cup \dots \cup R_{k-2}) \cap (C_1 \cup \dots \cup C_n)$, less or equal to $\frac{3}{7}(k-4)n$.

This gives the desired upper bound :

$$\kappa(k \times n) \leq \frac{3}{7}(k-4)n + 22 \lceil \frac{n}{14} \rceil.$$

For the lower bound, let S be an optimal solution of $\text{PEP}_{k \times n}$. First we claim that :

We may assume that there is at most 2 consecutive elements of S in (C_1, C_n, R_1, R_k) . (1)

Indeed, each solution S of the $\text{PEP}_{k \times n}$ can be transformed into another solution S' satisfying the claim by the following transformation : given 3 consecutive elements $s_{1,i-1}, s_{1,i}, s_{1,i+1}$ of $S \cap R_1$; set $S' = S \cup \{s_{2,i}\} \setminus \{s_{1,i}\}$. Clearly, repeating this process in C_1, C_n, R_1, R_k we will obtain a solution S' of $\text{PEP}_{k \times n}$ satisfying Claim 1 and such that $|S'| \leq |S|$.

Let S be an optimal solution of $\text{PEP}_{k \times n}$ satisfying Claim 1. We will construct a solution S^* of $\text{PEP}_{(2k+2) \times (2n+2)}$ such that $\{s + (a, (2n+2)), b, (2k+2) \mid s \in S^*, a, b \in \mathbb{Z}\}$ is a solution of $\text{PEP}_{\mathbb{Z}^2}$. We set $S^* = S \cup S_1 \cup S_2 \cup S_3 \cup B_1 \cup B_2 \cup B_3 \cup B_4 \cup B_5 \cup B_6 \cup B_7 \cup B_8 \cup T$ (see Figure 7), where :

- $B_1 = \{s_{i,n+1} \mid i \in \{1, \dots, k\}, s_{i,n} \notin S\}$, and $B_2 = \{s_{k+1,j} \mid j \in \{1, \dots, n\}, s_{k,j} \notin S\}$. Due to Claim 1, B_1 and B_2 satisfy the pentomino exclusion property (and even the trimino exclusion property).
- S_1 is the symmetric of S according to axis B_1 , i.e. $S_1 = \{s_{i,2n+2-j} \mid s_{i,j} \in S\}$. Analogously, define S_2 as the symmetric of S according to axis B_2 , i.e. $S_2 = \{s_{2k+2-i,j} \mid s_{i,j} \in S\}$.
- In the same way, $B_3 = \{s_{k+1,j} \mid j \in \{n+2, \dots, 2n+1\}, s_{k,j} \notin S_1\}$ and $B_4 = \{s_{i,n+1} \mid i \in \{k+2, \dots, 2k+1\}, s_{k,j} \notin S_2\}$.
- S_3 is the symmetric of S_1 according to axis B_3 , i.e. $S_3 = \{s_{2k+2-i,j} \mid s_{i,j} \in S_1\}$. Notice that, by construction, S_3 is also the symmetric of S_2 according to axis B_4 .
- $B_5 = \{s_{i,2n+2} \mid i \in \{1, \dots, k\}, s_{i,2n+1} \notin S_1\}$, $B_6 = \{s_{i,2n+2} \mid i \in \{k+2, \dots, 2k+1\}, s_{i,2n+1} \notin S_3\}$, $B_7 = \{s_{2k+2,j} \mid j \in \{1, \dots, n\}, s_{2k+1,j} \notin S_2\}$ and $B_8 = \{s_{2k+2,j} \mid j \in \{n+2, \dots, 2n+1\}, s_{2k+1,j} \notin S_3\}$.
- $T = \{s_{k+1,n+1}, s_{k+1,2n+2}, s_{2k+2,n+1}, s_{2k+2,2n+2}\}$.

The set S^* is a solution of $\text{PEP}_{(2k+2) \times (2n+2)}$. The corresponding $(2k+2) \times (2n+2)$ rectangle tiles \mathbb{Z}^2 by translation of vectors $(0, 2k+2)$ and $(2n+2, 0)$ which gives a solution of $\text{PEP}_{\mathbb{Z}^2}$. As the optimal density for $\text{PEP}_{\mathbb{Z}^2}$ is $\frac{3}{7}$, then the density of S^* is greater or equal to $\frac{3}{7}$.

Since the first and last rows and columns of S have density at least $\frac{1}{5}$, then the B_i 's have density at most $\frac{4}{5}$. As the S_i 's are symmetric of S then $|S_i| = |S| = \kappa_{k,n}$ for all i . Moreover $|T| = 4$, so finally we have :

$$\frac{3}{7}(2k+2)(2n+2) \leq |S^*| \leq 4\kappa_{k,n} + 4\left(\left\lceil \frac{4n}{5} \right\rceil + \left\lceil \frac{4k}{5} \right\rceil\right) + 4$$

Which leads to the desired bound :

$$\kappa_{k,n} \geq \frac{3}{7}kn + \frac{3}{7}(k+n) - \left(\left\lceil \frac{4n}{5} \right\rceil + \left\lceil \frac{4k}{5} \right\rceil\right) - \frac{4}{7}.$$

Observe that in the proof of the lower bound we need only the assumption that $k, n \geq 2$. \square

Let us mention as a direct consequence of the previous theorem, a maybe more explicit corollary which shows how close are our bounds :

Corollary 1 *For all $k, n \geq 5$. There are two constant C_1 and C_2 , such that :*

$$\frac{3}{7}kn - \frac{26}{70}(k+n) - C_1 \leq \kappa_{k \times n} \leq \frac{3}{7}kn - \frac{5}{70}(k+n) + C_2. \quad \square$$

2 The Δ -dislocation problem

2.1 Introduction

In this section we present a generalization of the PEP.

For a graph $G = (V, E)$ and a positive integer Δ , we will say that a subset S of vertices of G is a Δ -dislocation set of G if and only if all the connected components of $G - S$ have at most Δ vertices.

One can notice that if G is the grid $k \times n$ and $\Delta = 4$, then a Δ -dislocation set of G is a solution of $\text{PEP}_{k \times n}$.

Since $S = V(G)$ is always a set of Δ -dislocation of G for any Δ , then the corresponding optimisation problem is to find an S of minimum cardinality:

Δ -DISLOCATION

Instance : a graph G and an integer k

Question : is there a Δ -dislocation set S in G of cardinality $\leq k$?

Remark that for a graph G and $\Delta = 1$, a Δ -dislocation set S of G is a transversal of G , i.e. the complementary of S is an independent set of G . Since the problem TRANSVERSAL is NP-Complete[4], then for $\Delta = 1$ the problem Δ -DISLOCATION is NP-Complete. In the next subsection we will

prove that for any fixed Δ this problem is still NP-Complete.

Motivated by the $\text{PEP}_{k \times n}$ problem, we present in subsection 3 a polynomial time algorithm for solving Δ -dislocation problem in the grid graph $G_{k,n}$ whenever k is not apart of the input. More generally, our algorithm works for a largest class of graphs namely fasciagraphs. This algorithm was inspired by the work of Žerovnik and Klavžar on fasciagraphs [7].

2.2 Complexity of the Δ -dislocation problem

In this section we investigate the complexity of the Δ -dislocation problem.

Theorem 5 *For any Δ , we have that Δ -DISLOCATION problem is NP-Complete.*

Proof: The problem is clearly in NP. We will reduce it to TRANSVERSAL.

TRANSVERSAL

Instance : A graph G and an integer k

Question : Is there an transversal T in G of size $\leq k$?

Given a graph G , let us construct a graph G' such that G has a transversal of size $\leq k$ if and only if G' has a set of Δ -dislocation of size $\leq k$.

G' is constructed in the following way :

- To each vertex $v_i \in V(G)$ corresponds a clique K_i on Δ vertices in G' .
- In each clique K_i in G' let us choose one vertex v_i^*
- To each edge $v_i v_j \in E(G)$ corresponds one edge $v_i^* v_j^* \in E(G')$

If S is a subset $\{v_i\}_i$ of $V(G)$, then S^* will denote the corresponding subset $\{v_i^*\}_i$ of $V(G')$. Conversely for a subset $S^* = \{v_i^*\}_i$ of $V(G')$, S will denote the subset $\{v_i\}_i$ of $V(G)$.

★ Assume that G has a transversal T of size $\leq k$. Then T^* is a set of Δ -dislocation of G' . Indeed, as T is a transversal of G , then the connected components of G' are :

- the $K_i - v_i^*$ for all $v_i^* \in T^*$, of cardinality $\Delta - 1$
- the K_j for all $v_j^* \notin T^*$, of cardinality Δ

This shows that T^* is a set of Δ -dislocation of G .

★ On the other hand, assume that G' has a set S of Δ -dislocation of size $\leq k$. Let us show that G has an transversal of size $\leq k$.

Let us consider the set $T^* \subseteq V(G')$ defined by :

$$v_i^* \in T^* \iff K_i \cap S \neq \emptyset$$

As S is of size $\leq k$, then T^* is of size $\leq k$.

Consider the set T in $V(G)$ corresponding to T^* . Then we claim that T is a transversal of G . Indeed, if there exist an edge $v_i^* v_j^* \in E(G')$ such that $v_i^*, v_j^* \notin T^*$, it means that $K_i \cap T = K_j \cap T = \emptyset$, so that $K_i \cup K_j$ is contained in a connected component of $G' - T^*$ of size $\geq 2\Delta$, a contradiction. \square

2.3 Case of the Fasciagraphs

2.3.1 Fasciagraphs : definition

Let G_1, \dots, G_n be disjoint graphs and X_1, \dots, X_n a sequence of sets of edges such that an edge of X_i joins a vertex of $V(G_i)$ with a vertex of $V(G_{i+1})$, where G_{n+1} denotes the graph G_1 .

A *polygraph* $\Omega_n(G_1, \dots, G_n; X_1, \dots, X_n)$ over the *monographs* G_1, \dots, G_n is defined in the following way :

- $V(\Omega_n) = V(G_1) \cup V(G_2) \cup \dots \cup V(G_n)$
- $E(\Omega_n) = E(G_1) \cup X_1 \cup E(G_2) \cup \dots \cup X_{n-1} \cup E(G_n) \cup X_n$

The monographs G_j are also called the *fibers* of the polygraph.

Assume that for all $i = 1 \dots n$, G_i is isomorphic to a fixed graph G . In addition, let the sets X_i , $i = 1 \dots n - 1$ be equal to a fixed edge set X and $X_n = \emptyset$. Then we call the polygraph a *fasciagraph* and we denote it $\Psi_n(G, X)$.

2.3.2 Algorithm for fasciagraphs

Let G be a graph on k vertices v_1, \dots, v_k and let $\Psi_n(G, X)$ be a fasciagraph of fiber G . We will denote v_i^j the vertex v_i in the j -th fiber G_j of $\Psi_n(G, X)$. In this section we denote by Ψ_j , $j = 1 \dots n$ the subgraph of $\Psi_n(G, X)$ induced by the j -th first fibers G_1, \dots, G_j of $\Psi_n(G, X)$.

For a fixed $j \leq n$, a word (a_1, a_2, \dots, a_k) of $\{0, \dots, \Delta\}^k$ is said to be *valid* (for G_j) if there exists a set S of Δ -dislocation of Ψ_j such that :

$a_i = t$ iff v_i^j belongs to a connected component of cardinality k of $\Psi_j - S$

We will construct an auxiliary digraph $\mathcal{G}(G, X)$.

- The vertices of $\mathcal{G}(G, X)$ are the valid words of $\{0, \dots, \Delta\}^k$, plus two additional vertices B and E , that we call respectively *begin* and *end* vertices of $\mathcal{G}(G, X)$.
- There is an arc between B and any valid word $V \neq E$ for G_1 .
- There is an arc between any vertex $V \neq B$ and E .
- There is an arc between two valid words V and $W \neq B, E$ if and only if there exists a set S of Δ -dislocation and an integer j for which V is valid for G_j and W is valid for G_{j+1} .

Notice that there is a one to one mapping between the sets of Δ -dislocation of $\Psi_n(G, X)$ and the pathes on $n + 2$ vertices from B to E of $\mathcal{G}(G, X)$.

In addition, the graph is labeled in the following way :

- For any V and any $W \neq E$, the arc (V, W) is labeled by the number of zeros of the vector W .
- For any V , the arc (V, E) is labeled by 0.

Remark that for a given path on $n+2$ vertices $P = (B, V_1, V_2, \dots, V_n, E)$, the length of P is the sum of the number of zeros of V_1, V_2, \dots, V_n . So the length of the path is the cardinality of the corresponding set of Δ -dislocation. Then, determine the minimum cardinality of a Δ -dislocation set of $\Psi_n(G, X)$ is equivalent to compute the minimum length of a (B, E) -path on $n + 2$ vertices in $\mathcal{G}(G, X)$. This can be done in $O(K \log n)$ time, where K depends only on the size of $\mathcal{G}(G, X)$ (see for example [3] or [8]).

Theorem 6 *For a fixed k , one can compute the Δ -dislocation number of a fasciagraph $\Psi_n(G, X)$ in $O(\log n)$ time whenever $|V(G)| \leq k$.*

Proof: First observe that the size of $\mathcal{G}(G, X)$ depends only on k and Δ , since $\mathcal{G}(G, X)$ has at most $(\Delta + 1)^k$ vertices.

It remains, now, to prove that one can construct $\mathcal{G}(G, X)$ in $O(\log n)$ time. This comes from the fact that determining if a word of $\{0, \dots, \Delta\}^k$ is valid, and checking if two valid words are adjacent, can be done by computing all the Δ -dislocation sets of $\Psi_\Delta(G, X)$. Indeed, for $j > \Delta$, a word w is valid for G_j in $\Psi_n(G, X)$ if and only if w is valid for G_Δ in $\Psi_\Delta(G, X)$. \square

3 Open Problems

Roughly speaking, Theorem 4 shows that $\kappa_{k \times n}$ is approximately $\frac{3}{7}$ times the area of the $k \times n$ grid minus an improvement on the boundary $(k + n)$. For $k = 5$ and when 14 divides n , the upper bound of Theorem 4 gives $\kappa_{5 \times n} \leq 2n$. Moreover, Theorem 2 tells us that $\kappa_{5 \times n} = 2n - 2$. Therefore, one may ask :

Conjecture 1 *For any k and n ‘sufficiently large’ (with $n \geq k$), we have*

$$\kappa_{k \times n} = \lfloor \frac{3}{7}(k - 4)n \rfloor + 22 \lfloor \frac{n}{14} \rfloor - O(1).$$

From the algorithmic point of view, one may consider a weaker question related to the grid graph. Theorem 6 proves that determining $\kappa_{k \times n}$ for a grid graph is a polynomial task whenever k is fixed. So a weaker version of Conjecture 1 is :

Conjecture 2 *It is polynomially solvable to decide whether $\kappa_{k \times n} \leq P$ for given integers P, k and n .*

References

- [1] R.A. Bosch, *A Pentomino Exclusion Problem*. Mathematical Programming Newsletter, Optima 60 (december 1998), 3.
- [2] R.A. Bosch, *Peaceably Coexisting Armies of Queens*. Mathematical Programming Newsletter, Optima 62 (june 1999), 6.
- [3] B. Carré, *Graphs and Networks*. Clarendon Press, Oxford, (1979).
- [4] M.R. Garey, D.S. Johnson, *Computers and Intractability a Guide to the Theory of NP-Completeness* A Series of Books in the mathematical Sciences. San Francisco: W. H. Freeman and Company (1979).

- [5] S.W. Golomb, *Polyominoes - Puzzles, Patterns, Problems, and Packings*. Princeton Science Library, (1994).
- [6] S. Gravier, C. Payan, *On the Pentomino Exclusion Problem*. Disc. and Comput. Geom. 26 (2001), 375-386.
- [7] S. Klavžar, J. Žerovnik, *Algebraic Approach to Fasciagraphs and Rota-graphs*. Discrete Applied Mathematics 60 (1996) 93-100.
- [8] U. Zimmermann, *Linear and Combinatorial Optimization in Ordered Algebraic Structures*. Ann. Discrete Math. 10 (1981).




Figure 3: Some solutions of $PEP_{5 \times n}$ for $n \geq 5$.




Figure 4: Optimal solution of $\mathbb{Z}\mathbb{Z}^2$.




Figure 5: Construction of a solution of $\text{PEP}_{k \times \mathbb{Z}}$.




Figure 6: Proof of Claim 1.




Figure 7: Construction of S^* from S .




Figure 8: Construction of G' from G .




Figure 9: An example of fasciagraph over monographs K_3 .

Le laboratoire Leibniz est fortement pluridisciplinaire. Son activité scientifique couvre un large domaine qui comprend aussi bien des thèmes fondamentaux que des thèmes très liés aux applications, aussi bien en mathématiques qu'en informatique.

Les recherches sur les Environnements Informatiques d'Apprentissage Humain et la didactique des mathématiques ouvrent cette pluridisciplinarité sur les sciences humaines, elles jouent un rôle particulier en favorisant les coopérations entre différentes composantes du laboratoire.

- * mathématiques discrètes et recherche opérationnelle
- * logique et mathématique pour l'informatique
- * informatique de la connaissance
- * EIAH et didactique des mathématiques

Les cahiers du laboratoire Leibniz ont pour vocation la diffusion des rapports de recherche, des séminaires ou des projets de publication réalisés par des membres du laboratoire. Au-delà, Les cahiers peuvent accueillir des textes de chercheurs qui ne sont pas membres du laboratoire Leibniz mais qui travaillent sur des thèmes proches et ne disposent pas de tels supports de publication. Dans ce dernier cas, les textes proposés sont l'objet d'une évaluation par deux membres du Comité de Rédaction.

Comité de rédaction

- * mathématiques discrètes et recherche opérationnelle
Gerd Finke, András Sebő
- * logique et mathématique pour l'informatique
Ricardo Caferra, Rachid Echahed
- * informatique de la connaissance
Yves Demazeau, Daniel Memmi,
- * EIAH et didactique des mathématiques
Nicolas Balacheff, Jean-Luc Dorier, Denise Grenier

Contact Gestion & Réalisation : Jacky Coutin
Directeur de la publication : Nicolas Balacheff
ISSN : 1298-020X - © laboratoire Leibniz