

Logarithmic stability in determination of a 3D viscoelastic coefficient and a numerical example

Maya de Buhan^{1,2} and Axel Osses²

¹ UPMC Univ Paris 06, UMR 7598, Laboratoire J.L. Lions, F-75005 Paris, France

² Departamento de Ingeniería Matemática and Centro de Modelamiento Matemático (UMI 2807 CNRS), FCFM Universidad de Chile, Casilla 170/3-Correo 3, Santiago, Chile

E-mail: debuhan@ann.jussieu.fr and axosses@dim.uchile.cl

Received 6 December 2009, in final form 5 May 2010

Published 9 July 2010

Online at stacks.iop.org/IP/26/095006

Abstract

We prove a Carleman estimate and a logarithmic stability estimate for an inverse problem in three-dimensional viscoelasticity. More precisely, we obtain logarithmic stability for the inverse problem of recovering the spatial part of a viscoelastic coefficient of the form $p(x)h(t)$ from a unique measurement on an arbitrary part of the boundary. The main assumptions are $h'(0) = 0$, $h(0) \neq 0$, p is known in a neighborhood of the boundary and regularity and sensitivity of the reference trajectory. We propose a method to solve the problem numerically and illustrate the theoretical result by a numerical example.

(Some figures in this article are in colour only in the electronic version)

1. Introduction and main results

This paper is concerned with the inverse problem of determining an unknown coefficient in the 3D viscoelasticity system. We set Ω an open bounded domain of \mathbb{R}^3 with a sufficiently smooth boundary $\partial\Omega$. Then the viscoelasticity system, endowed with initial and boundary conditions, is as follows:

$$\begin{cases} \mathcal{P}u(x, t) = f(x, t), & \forall (x, t) \in \Omega \times (0, +\infty), \\ u(x, 0) = \bar{u}_0(x), & \forall x \in \Omega, \\ \partial_t u(x, 0) = \bar{u}_1(x), & \forall x \in \Omega, \\ u(x, t) = 0, & \forall (x, t) \in \partial\Omega \times (0, +\infty), \end{cases} \quad (1)$$

where \mathcal{P} is the integro-differential hyperbolic operator defined as

$$\begin{aligned} \mathcal{P}u(x, t) = & \partial_t^2 u(x, t) - \nabla \cdot (\mu(x)(\nabla u(x, t) + \nabla u(x, t)^T) + \lambda(x)(\nabla \cdot u)(x, t)I) \\ & + \int_0^t \nabla \cdot (\tilde{\mu}(x, s)(\nabla u(x, t-s) + \nabla u(x, t-s)^T) + \tilde{\lambda}(x, s)(\nabla \cdot u)(x, t-s)I) ds. \end{aligned} \quad (2)$$

This system models the dynamics of a 3D viscoelastic material subjected to a load f , u being the displacement vector, \bar{u}_0 and \bar{u}_1 the initial displacement and velocity, (λ, μ) the Lamé coefficients and $(\tilde{\lambda}, \tilde{\mu})$ the viscosity coefficients, respectively. The well-posedness nature of the direct problem (1)–(2) is guaranteed by the following proposition.

Proposition 1 (Theorem 4.2 in [1]). *Under regularity assumptions on the coefficients $(\lambda, \mu, \tilde{\lambda}, \tilde{\mu})$, if $\bar{u}_0 \in H_0^1(\Omega)^3$, $\bar{u}_1 \in L^2(\Omega)^3$ and $f \in L^\infty(0, +\infty; L^2(\Omega)^3) \cap L^2(0, +\infty; L^2(\Omega)^3)$, then there exists a unique weak solution $u \in L^\infty(0, +\infty; H_0^1(\Omega)^3) \cap W^{1,\infty}(0, +\infty; L^2(\Omega)^3) \cap W^{2,\infty}(0, +\infty; H^{-1}(\Omega)^3)$ to the problem (1)–(2).*

If we now assume that the coefficient $\tilde{\mu}$ can be decomposed as follows:

$$\tilde{\mu}(x, t) = p(x)h(t), \quad \forall (x, t) \in \Omega \times (0, +\infty), \quad (3)$$

then the inverse problem we are interested in is formulated accordingly.

Definition 1 (Inverse problem). *Given $(\lambda, \mu, \tilde{\lambda}, h, u_0, u_1, f)$, recover $p(x)$, for all $x \in \Omega$, from measurements of*

$$u(x, t), \quad \forall (x, t) \in \Gamma \times (0, T),$$

where Γ is a part of $\partial\Omega$ and $T > 0$.

Several authors have dealt with the problem of recovering the coefficients of the viscoelasticity system (1)–(2). Some of them, e.g. Grasselli [2], Janno *et al* [3], von Wolfersdorf [4], Cavaterra *et al* [5], recovered the time dependence h of the coefficient $\tilde{\mu}$ by reducing the problem to a nonlinear Volterra integral equation by using Fourier's method to solve the direct problem and by applying the contraction principle. Others, e.g. Lorenzi [6], Lorenzi and Romanov [7], recovered the space dependence p of the coefficient $\tilde{\mu}$ by using the method of Bukhgeim and Klibanov [8] based on Carleman estimates [9]. Thus, in 2007, Lorenzi *et al* [10] recovered the coefficient p from three measurements, relying on the assumption that all the coefficients in the operator (2) are independent of the third space variable. Here, unlike the latter, we recover p from a unique measurement, assuming that it is known in a neighborhood of the boundary.

Furthermore, theorem 1 establishes the logarithmic stability with respect to a unique measurement on an arbitrary part of the boundary for the inverse problem of recovering the coefficient p . Its proof is given in section 2. It relies on a Carleman estimate (theorem 2) and a sharp unique continuation result (theorem 3) for the operator (2). We prove these results in sections 3 and 4 respectively. In section 5, we propose an adaptive spectral method to approximate the solution of the inverse problem numerically.

1.1. Stability estimate

Inverse problems are ill-posed in the classical sense [11]. Stability estimates play thus a special role in the theory. Bukhgeim and Klibanov [8] developed a remarkable method based on Carleman estimates [9] to prove the uniqueness and stability for inverse problems associated with partial differential equations. In [12], using this method, we proved a Hölder stability result with a unique internal measurement for the recovery of p in the system (1)–(3). In this paper, we extend this result using a unique continuation estimate and obtain a logarithmic stability result. Although this estimate is weaker, it is nonetheless related to a measurement on an arbitrary part of the boundary. A similar result was proved in [13] for the hyperbolic scalar equation and in [14] for the Lamé system. We follow their method but adapt it to take into account the additive integral term of operator (2). Thus, our result can be seen as an extension

of the paper [14]. Nevertheless, the presence of the integral term modifies the structure of the equation. Firstly, the latter is no longer reversible because of the energy dissipation associated with the viscous phenomenon. In practice, this prevents the extension of the solution to $[-T, 0]$ as usually performed. This explains why we need to use a point-wise Carleman estimate (see section 1.2). Secondly, the coefficients of the equation now depend on time and not only on space as for the Lamé system. This implies that the Fourier–Bros–Iagolnitzer (FBI) transform classically used to prove the unique continuation result (see section 1.3) is useless. Hence, we have to introduce a new transformation in order to deal with convolution terms. We now need to define the following condition on a scalar function q .

Condition 1. *The scalar function q is said to satisfy condition 1 if*

- (i) *there exists $K > 0$ such that $\forall x \in \overline{\Omega}$, $q(x) \geq K$,*
- (ii) *there exists $x_0 \in \mathbb{R}^3 \setminus \overline{\Omega}$ such that $\forall x \in \overline{\Omega}$, $\frac{1}{2}q(x) - \nabla q(x) \cdot (x - x_0) \geq 0$.*

We can now state the main result.

Theorem 1 (Logarithmic stability). *Let u (respectively \bar{u}) be the solution of the system (1)–(3), associated with the coefficient p (respectively \bar{p}). We assume that*

- (H1) *$(\lambda, \mu) \in C^2(\overline{\Omega})^2$ and $(\tilde{\lambda}, \tilde{\mu}) \in C^2(\overline{\Omega \times (0, +\infty)})^2$ are such that the solutions u and $\bar{u} \in W^{8,\infty}(\Omega \times (0, +\infty))^3$,*
- (H2) *μ and $\lambda + 2\mu$ satisfy condition 1 with the same x_0 ;*
- (H3) *$p = \bar{p}$ is known in a neighborhood ω of $\partial\Omega$;*
- (H4) *$h(0) \neq 0$, $h'(0) = 0$;*
- (H5) *there exists $M > 0$ such that $\forall x \in \overline{\Omega \setminus \omega}$, $|\nabla \bar{u}_0 + \nabla \bar{u}_0^T(x) \cdot (x - x_0)| \geq M$.*

Then, for $\Gamma \subset \partial\Omega$ arbitrarily small, there exist $\kappa \in (0, 1)$ and $T_0 > 0$ such that, for all $T > T_0$, the following estimate holds:

$$\|p - \bar{p}\|_{H^2(\Omega)} \leq C \left[\log \left(2 + \frac{C}{\sum_{1 \leq |\alpha| \leq 2} \|\partial_x^\alpha (u - \bar{u})\|_{L^2(\Gamma \times (0, T))}^2} \right) \right]^{-\kappa}$$

where $C > 0$ depends on the $C^2(\overline{\Omega})$ -norm of p and \bar{p} and on the $W^{8,\infty}(\Omega \times (0, T))^3$ -norm of u and \bar{u} .

An example of an initial datum \bar{u}_0 verifying (H5) corresponds to the choice $\bar{u}_0 \in H_0^1(\Omega)^3$ such that $\bar{u}_0(x) = x$, $\forall x \in \overline{\Omega \setminus \omega}$. Thus,

$$|\nabla \bar{u}_0 + \nabla \bar{u}_0^T(x) \cdot (x - x_0)| = |x - x_0| \geq d_0 > 0, \quad \forall x \in \overline{\Omega \setminus \omega}.$$

From theorem 1, we immediately deduce corollary 1.

Corollary 1 (Uniqueness). *Under all the hypothesis of theorem 1, we have then*

$$\begin{aligned} \partial_x^\alpha u(x, t) &= \partial_x^\alpha \bar{u}(x, t), & \forall (x, t) \in \Gamma \times (0, T), & \quad |\alpha| = 1, 2 \\ \implies p(x) &= \bar{p}(x), & \forall x \in \Omega. \end{aligned}$$




Figure 1. The Carleman weight function $\varphi(x, t)$ (left) and its isovalues $\varphi(x, t) = \delta$ (right) for $x_0 = 0$ and $\beta = 1$.

1.2. A Carleman estimate

In a pioneering work [9], Carleman introduced what is now commonly known as a Carleman estimate, in the context of proving the uniqueness in the Cauchy elliptic problem. Since then, the theory of Carleman estimates has been extensively studied. As for a general treatment of Carleman estimates, see Hörmander [15], Isakov [16], Tataru [17], Imanuvilov and Yamamoto [18]. In 2006, Cavaterra *et al* [19] modified the point-wise Carleman inequality of Klibanov and Timonov [20] for a hyperbolic scalar equation and integrated, thanks to a change of variable, the integral term of (2). In 2008, Imanuvilov and Yamamoto [21] proved a Carleman estimate for the 3D Lamé system—operator (2) without the integral term. Their approach consisted in decoupling the system by writing the equations satisfied by u , $\nabla \wedge u$ and $\nabla \cdot u$ in view of applying the well-known results for hyperbolic scalar equations [22]. In this paper, we prove a Carleman estimate for the complete integro-differential operator (2) by the combination of these two techniques. We start by introducing the following Carleman weight function.

Definition 2 (A Carleman weight function). For $x_0 \in \mathbb{R}^3 \setminus \overline{\Omega}$ and $\beta > 0$, we introduce a function φ in the following way:

$$\varphi(x, t) = |x - x_0|^2 - \beta t^2, \quad \forall (x, t) \in \Omega \times (0, +\infty). \quad (4)$$

This function is plotted in figure 1. Let us now introduce some notations used in what follows.



- Distances (see figure 2): for $x_0 \in \mathbb{R}^3 \setminus \overline{\Omega}$, we note

$$d_0 = \inf_{x \in \Omega} |x - x_0| \quad \text{and} \quad d = \sup_{x \in \Omega} |x - x_0|^2.$$

- Domains (see figure 3): for $\varepsilon > 0$ and $\delta > 0$, we note

$$\begin{aligned} \Omega(\varepsilon) &= \{x \in \Omega, \text{dist}(x, \partial\Omega) > \varepsilon\}, & \Omega(0) &= \Omega, \\ \mathcal{Q}(\varepsilon, \delta) &= \{(x, t) \in \Omega(\varepsilon) \times (0, +\infty), \varphi(x, t) > \delta\}, & \mathcal{Q}(0, 0) &= \mathcal{Q}. \end{aligned}$$

We note that $\mathcal{Q} \subset \Omega \times (0, T)$ if $T > \frac{d}{\sqrt{\beta}}$.

**Figure 2.** Definitions of the distances d_0 and d .**Figure 3.** Definitions of the domains $\Omega(\varepsilon)$ and $Q(\varepsilon, \delta)$.

- Norms: for $Q \subset \mathbb{R}^3 \times \mathbb{R}$, $\sigma > 0$ and $k \in \mathbb{N}$, we introduce the following norms:

$$\begin{aligned} \|\star\|_{H^{1,\sigma}(Q)}^2 &= \|\nabla_{x,t} \star\|_{L^2(Q)}^2 + \sigma^2 \|\star\|_{L^2(Q)}^2, \\ \|\star\|_{H_x^{k,\sigma}(Q)}^2 &= \sum_{|\alpha| \leq k} \sigma^{2(k-|\alpha|)} \|\partial_x^\alpha \star\|_{L^2(Q)}^2. \end{aligned}$$

- Constants: C and C_j , $1 \leq j \leq 6$, are generic positive constants.

Equipped with these definitions, we can now state the following result.

Theorem 2 (Carleman estimate). *Let \mathcal{P} be the operator defined by (1). We assume that*

(H1) $(\lambda, \mu) \in C^2(\overline{\Omega})^2$ and $(\tilde{\lambda}, \tilde{\mu}) \in C^2(\overline{\Omega} \times (0, +\infty))^2$,

(H2) μ and $\lambda + 2\mu$ satisfy condition 1 with the same x_0 .

Then, there exist $\beta_0 > 0$ and $\sigma_0 > 0$ such that, for all $\beta \leq \beta_0$, for all $\sigma \geq \sigma_0$, for all $\varepsilon > 0$, for all $0 < \delta < d_0^2$ and for all $u \in H^2(Q)^3$ satisfying $u(x, 0) = 0$ or $\partial_t u(x, 0) = 0$, $\forall x \in \Omega$, the following estimate holds:

$$\begin{aligned} \frac{C}{\sigma} \|u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 &\leq \|(\mathcal{P}u) e^{\sigma\varphi}\|_{L^2(Q)}^2 + \|\nabla(\mathcal{P}u) e^{\sigma\varphi}\|_{L^2(Q)}^2 \\ &\quad + e^{C\sigma} \|u\|_{H^2(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,\delta))}^2 + \sigma^3 e^{2\sigma\delta} \|u\|_{H^2(Q)}^2, \end{aligned}$$

where φ is defined by (4) and $C > 0$ depends on the $C^2(\overline{Q})$ -norm of the coefficient $\tilde{\mu}$ but is independent of σ .

The last term of this estimate is rather uncommon since it is global but it will disappear thanks to a clever choice of σ .

1.3. A unique continuation result

In [23], Bellassoued proved a sharp unique continuation result for the Lamé system. He applied a method initially developed by Robbiano [24] which uses the FBI transform [25] to change the problem near the boundary into a problem for which elliptic estimates can be applied. In this paper, we show a similar result for the system (1)–(2) by adapting these techniques. In particular, we propose a new transformation inspired from the FBI transform but which is able to deal with the additive convolution term of the operator (2). The result we prove is as follows.

Theorem 3 (Unique continuation). *Let u be the vector solution of*

$$\begin{cases} \mathcal{P}u(x, t) = R(x, t), & \forall (x, t) \in \Omega \times (0, +\infty), \\ u(x, 0) = 0, & \forall x \in \Omega, \\ \partial_t u(x, 0) = 0, & \forall x \in \Omega, \\ u(x, t) = 0, & \forall (x, t) \in \partial\Omega \times (0, +\infty). \end{cases}$$

We assume that

(H1) $(\lambda, \mu) \in C^2(\overline{\Omega})^2$ and $(\tilde{\lambda}, \tilde{\mu}) \in C^2(\overline{\Omega \times (0, +\infty)})^2$ are such that the solution $u \in W^{4,\infty}(\Omega \times (0, +\infty))^3$,

(H6) $R(x, t) = 0, \forall (x, t) \in \omega \times (0, +\infty)$ where ω is a neighborhood of $\partial\Omega$.

Then, for $\Gamma \subset \partial\Omega$ arbitrarily small, there exists $T_0 > 0$ such that, for all $T \geq T_0$ and for all $\rho > 0$, the following estimate holds:

$$\|u\|_{H^2(\omega \times (0, \frac{T}{2} - \rho))}^2 \leq C \left[\log \left(2 + \frac{C}{\sum_{1 \leq |\alpha| \leq 2} \|\partial_x^\alpha u\|_{L^2(\Gamma \times (0, 3T))}^2} \right) \right]^{-1}$$

where $C > 0$ depends on the $C^2(\overline{\Omega \times (0, T)})$ -norm of the coefficient $\tilde{\mu}$ and on the norm $W^{4,\infty}(\Omega \times (0, T))^3$ of u .

1.4. Numerical results

Section 5 presents a numerical example to illustrate one of our theoretical results [12]. Here, we propose a non-quadratic functional to solve the inverse problem. Moreover, as a regularization method [26], we use a spectral basis, adapted in space and in frequency to the solution. This idea, combined with mesh adaptation, allows us to improve the accuracy of the method by minimizing the numerical error. The unknown parameter is successfully recovered at each vertex of the discretized domain.

2. Proof of theorem 1

This section presents the proof of the stability result (theorem 1). The idea of the proof is the following. Firstly, we bring the unknown parameter p to the source by writing the equation satisfied by $\hat{v} = \partial_t(u - \bar{u})$. Then, we use the method of Bukhgeim and Klivanov; i.e. we differentiate the previous equation to bring the parameter in the initial condition and we apply the Carleman estimate of theorem 2 to the new variable $\partial_t \hat{v}$ in order to bound the initial energy. Thanks to a Carleman estimate for a first-order operator (lemma 2), we come back to the coefficient in the estimate. Thus, we obtain a first Hölder stability result with an internal measurement on a neighborhood of the boundary. Finally, we combine this result with the unique continuation relation of theorem 3 to conclude.

Let u (respectively \bar{u}) be the solution of the system (1)–(3), associated with the coefficient p (respectively \bar{p}). We suppose that hypotheses (H1)–(H5) hold and we fix $\Gamma \subset \partial\Omega$ arbitrarily small. By linearity and without loss of generality, we can suppose that $\bar{u}_1 = 0$.

2.1. Bring the unknown parameter p to the source

We suppose (\bar{p}, \bar{u}) to be known and we introduce $\hat{p} = p - \bar{p}$ and $\hat{u} = u - \bar{u}$ which satisfy the following equation:

$$\begin{aligned} \mathcal{P}\hat{u}(x, t) &= \partial_t^2 \hat{u}(x, t) - \nabla \cdot (\mu(x)(\nabla \hat{u}(x, t) + \nabla \hat{u}(x, t)^T) + \lambda(x)(\nabla \cdot \hat{u})(x, t)I) \\ &\quad + \int_0^t \nabla \cdot (h(s)p(x)(\nabla \hat{u}(x, t-s) + \nabla \hat{u}(x, t-s)^T) \\ &\quad + \tilde{\lambda}(x, s)(\nabla \cdot \hat{u})(x, t-s)I) \, ds \\ &= - \int_0^t h(s) \nabla \cdot (\hat{p}(x)(\nabla \bar{u}(x, t-s) + \nabla \bar{u}(x, t-s)^T)) \, ds, \\ &\quad \forall (x, t) \in \Omega \times (0, +\infty), \end{aligned}$$

with null initial and boundary conditions. We note that the coefficient \hat{p} we wish to recover appears in the source term. The problem is that this source term vanishes at the initial time $t = 0$, so it does not satisfy the hypothesis of the method of Bukhgeim and Klivanov [8]. To overcome this problem, we derive the equation in time and set

$$\hat{v}(x, t) = \partial_t \hat{u}(x, t), \quad \forall (x, t) \in \Omega \times (0, +\infty).$$

Then, \hat{v} satisfies the equation

$$\begin{aligned} \mathcal{P}\hat{v}(x, t) &= - \int_0^t h(s) \nabla \cdot (\hat{p}(x)(\nabla \bar{v}(x, t-s) + \nabla \bar{v}(x, t-s)^T)) \, ds \\ &\quad - h(t) \nabla \cdot (\hat{p}(x)(\nabla \bar{u}_0(x) + \nabla \bar{u}_0(x)^T)), \quad \forall (x, t) \in \Omega \times (0, +\infty), \end{aligned} \quad (5)$$

with null initial and boundary conditions. As a consequence, the unknown coefficient is still in the source term but it no longer vanishes at the initial time.

2.2. Use the method of Bukhgeim and Klivanov

We derive (5) and set

$$\hat{w}(x, t) = \partial_t \hat{v}(x, t) = \partial_t^2 \hat{u}(x, t), \quad \forall (x, t) \in \Omega \times (0, +\infty),$$

which satisfies then

$$\begin{aligned} \mathcal{P}\hat{w}(x, t) &= - \int_0^t h(s) \nabla \cdot (\hat{p}(x)(\nabla \bar{w}(x, t-s) + \nabla \bar{w}(x, t-s)^T)) \, ds \\ &\quad - h'(t) \nabla \cdot (\hat{p}(x)(\nabla \bar{u}_0(x) + \nabla \bar{u}_0(x)^T)), \quad \forall (x, t) \in \Omega \times (0, +\infty), \end{aligned} \quad (6)$$

with the following initial and boundary conditions:

$$\begin{cases} \hat{w}(x, 0) = 0, & \forall x \in \Omega, \\ \partial_t \hat{w}(x, 0) = -h(0) \nabla \cdot (\hat{p}(x)(\nabla \bar{u}_0(x) + \nabla \bar{u}_0(x)^T)), \\ \hat{w}(x, t) = 0, & \forall (x, t) \in \partial\Omega \times (0, +\infty). \end{cases} \quad (7)$$

Therefore, the coefficient \hat{p} that we want to recover appears now in the initial conditions. We are going to write an inequality by bounding the energy of the system (7) by the source and the observations. To do that, we apply the Carleman estimate of theorem 2 to \hat{w} and to its time derivatives.

2.3. Apply the Carleman estimate

Theorem 2 states that for φ defined by equation (4) with β sufficiently small, for σ sufficiently large, for $\varepsilon > 0$ and $0 < \delta < d_0^2$, the following estimate holds:

$$\begin{aligned} \frac{1}{\sigma} \|\hat{w} e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 &\leq C \left(e^{C\sigma} \|\hat{w}\|_{H^2(Q(\varepsilon,\delta)\setminus Q(2\varepsilon,\delta))}^2 + \sigma^3 e^{2\sigma\delta} \|\hat{w}\|_{H^2(Q)}^2 \right. \\ &\quad + \left. \int_Q \left| \int_0^t h(s) \nabla \cdot (\hat{p}(x) (\nabla \bar{w}(x, t-s) + \nabla \bar{w}(x, t-s)^T)) \, ds \right|^2 \right. \\ &\quad \times e^{2\sigma\varphi(x,t)} \, dx \, dt + \int_Q \left| \int_0^t h(s) \nabla (\nabla \cdot (\hat{p}(x) (\nabla \bar{w}(x, t-s) \right. \\ &\quad + \nabla \bar{w}(x, t-s)^T)) \, ds \left. \right|^2 e^{2\sigma\varphi(x,t)} \, dx \, dt + \|h' \nabla \cdot (\hat{p}(\nabla \bar{u}_0 + \nabla \bar{u}_0^T)) e^{\sigma\varphi}\|_{L^2(Q)}^2 \\ &\quad \left. + \|h' \nabla (\nabla \cdot (\hat{p}(\nabla \bar{u}_0 + \nabla \bar{u}_0^T))) e^{\sigma\varphi}\|_{L^2(Q)}^2 \right). \end{aligned} \quad (8)$$

We now need a result to bound the integral terms. This is given by the following lemma.

Lemma 1 (Lemma 3.1.1 in [20]). *Let φ be defined by (4). There exists $C > 0$ such that, for all $\sigma > 0$ and $u \in L^2(Q)$,*

$$\int_Q \left(\int_0^t |u(x, s)| \, ds \right)^2 e^{2\sigma\varphi(x,t)} \, dx \, dt \leq \frac{C}{\sigma} \int_Q |u(x, t)|^2 e^{2\sigma\varphi(x,t)} \, dx \, dt.$$

We use the result of lemma 1 and the fact that (\hat{p}, h) is bounded in $\Omega \times (0, +\infty)$ (H1) to write

$$\begin{aligned} \frac{C}{\sigma} \|\hat{w} e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 &\leq e^{C\sigma} \|\hat{w}\|_{H^2(Q(\varepsilon,\delta)\setminus Q(2\varepsilon,\delta))}^2 + \sigma^3 e^{2\sigma\delta} \|\hat{w}\|_{H^2(Q)}^2 \\ &\quad + \|\nabla \cdot (\hat{p}(\nabla \bar{u}_0 + \nabla \bar{u}_0^T)) e^{\sigma\varphi}\|_{L^2(Q)}^2 + \|\nabla (\nabla \cdot (\hat{p}(\nabla \bar{u}_0 + \nabla \bar{u}_0^T))) e^{\sigma\varphi}\|_{L^2(Q)}^2. \end{aligned}$$

Thus, taking into account the fact that $\bar{u}_0 \in W^{8,\infty}(\overline{\Omega \times (0, +\infty)})$ (H1), we obtain

$$\begin{aligned} \frac{C}{\sigma} \|\hat{w} e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 &\leq e^{C\sigma} \|\partial_t^2 \hat{u}\|_{H^2(Q(\varepsilon,\delta)\setminus Q(2\varepsilon,\delta))}^2 + \sigma^3 e^{2\sigma\delta} \|\partial_t^2 \hat{u}\|_{H^2(Q)}^2 \\ &\quad + \sum_{|\alpha| \leq 2} \|(\partial_x^\alpha \hat{p}) e^{\sigma\varphi}\|_{L^2(Q)}^2. \end{aligned}$$

Similarly, we obtain, for $i = 1, 2$,

$$\begin{aligned} \frac{C}{\sigma} \|(\partial_t^i \hat{w}) e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 &\leq e^{C\sigma} \|\partial_t^{2+i} \hat{u}\|_{H^2(Q(\varepsilon,\delta)\setminus Q(2\varepsilon,\delta))}^2 + \sigma^3 e^{2\sigma\delta} \|\partial_t^{2+i} \hat{u}\|_{H^2(Q)}^2 \\ &\quad + \sum_{|\alpha| \leq 2} \|(\partial_x^\alpha \hat{p}) e^{\sigma\varphi}\|_{L^2(Q)}^2. \end{aligned} \quad (9)$$

In order to apply the Carleman estimate, we verified that the functions $\partial_t \hat{w}$ and $\partial_t^2 \hat{w}$ have one of their initial conditions null. This requirement is true since we supposed $h'(0) = 0$ in (H4). We will now bound the initial energy of the system (7).




Figure 4. Definition of the cut-off function χ_1 .

2.4. Bound the initial energy

We define the cut-off function $\chi_1 \in C^\infty(\mathbb{R}^3 \times \mathbb{R})$ by $0 \leq \chi_1 \leq 1$ and

$$\chi_1(x, t) = \begin{cases} 1, & \text{if } (x, t) \in Q(0, 2\delta), \\ 0, & \text{if } (x, t) \in Q \setminus \overline{Q(0, \delta)}, \end{cases}$$

and we set

$$w^*(x, t) = \chi_1(x, t) \hat{w}(x, t), \quad \forall (x, t) \in \Omega \times (0, +\infty).$$

This function is illustrated in figure 4.

Since the time derivative of w^* is null outside $Q(0, \delta)$, i.e. for

$$t \geq t_x^\delta = \sqrt{\frac{|x - x_0|^2 - \delta}{\beta}}$$

which is well defined for all $x \in \Omega$ because $\delta < d_0^2$, we can write

$$\begin{aligned} \int_{\Omega(\varepsilon)} |\partial_t w^*(x, 0)|^2 e^{2\sigma\varphi(x, 0)} dx &= - \int_{\Omega(\varepsilon)} \int_0^{t_x^\delta} \partial_t (|\partial_t w^*(x, t)|^2 e^{2\sigma\varphi(x, t)}) dx dt \\ &= \int_{Q(\varepsilon, \delta)} (4\beta t \sigma |\partial_t w^*(x, t)|^2 e^{2\sigma\varphi(x, t)} \\ &\quad - 2\partial_t w^*(x, t) \cdot \partial_t^2 w^*(x, t) e^{2\sigma\varphi(x, t)}) dx dt. \end{aligned}$$

In addition, we have

$$\partial_t w^*(x, t) = \chi_1(x, t) \partial_t \hat{w}(x, t) + \partial_t \chi_1(x, t) \hat{w}(x, t), \quad \forall (x, t) \in \Omega \times (0, +\infty),$$

$$\partial_t^2 w^*(x, t) = \chi_1(x, t) \partial_t^2 \hat{w}(x, t) + 2\partial_t \chi_1(x, t) \partial_t \hat{w}(x, t) + \partial_t^2 \chi_1(x, t) \hat{w}(x, t).$$

Therefore, we obtain

$$\begin{aligned} \int_{\Omega(\varepsilon)} |\partial_t w^*(x, 0)|^2 e^{2\sigma\varphi(x, 0)} dx \\ \leq C \left(\int_{Q(\varepsilon, \delta)} (\sigma |\hat{w}(x, t)|^2 + \sigma |\partial_t \hat{w}(x, t)|^2 + |\partial_t^2 \hat{w}(x, t)|^2) e^{2\sigma\varphi(x, t)} dx dt \right). \end{aligned}$$

We do the same work for the derivatives in x for $|\alpha| \leq 2$ and thus we obtain

$$\begin{aligned} \sum_{|\alpha| \leq 2} \sigma^{2(1-|\alpha|)} \int_{\Omega(\varepsilon)} |\partial_x^\alpha (\partial_t \hat{w})(x, 0)|^2 e^{2\sigma\varphi(x,0)} dx \\ \leq \frac{C}{\sigma} \left(\|\hat{w} e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 + \|(\partial_t \hat{w}) e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 + \|(\partial_t^2 \hat{w}) e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 \right) \\ \leq C \left(\sum_{|\alpha| \leq 2} \|(\partial_x^\alpha \hat{p}) e^{\sigma\varphi}\|_{L^2(Q)}^2 + e^{C\sigma} \|\hat{u}\|_{H^6(Q(\varepsilon,2\delta) \setminus Q(2\varepsilon,\delta))}^2 + \sigma^3 e^{2\sigma\delta} \|\hat{u}\|_{H^6(Q)}^2 \right), \end{aligned}$$

where we used (9) in the last inequality. We have to go back to the coefficient on the left-hand side.

2.5. Return to the coefficient in the estimate

To this end, we need a Carleman estimate for a first-order operator. It is given by the following lemma.

Lemma 2 (Lemma 3.2 in [18]). *We consider the following first-order partial differential operator:*

$$\mathcal{R}(x)\star = a(x) \cdot \nabla \star + a_0(x)\star, \quad \forall x \in \Omega,$$

with

$$\begin{aligned} a_0 \in C^0(\overline{\Omega}), \quad \|a_0\|_{C^2(\overline{\Omega})} \leq M_1 \quad \text{and} \quad a \in C^1(\overline{\Omega})^n, \quad \|a\|_{C^2(\overline{\Omega})} \leq M_2, \\ \exists x_0, \quad |a(x) \cdot (x - x_0)| \geq M_3 > 0, \quad \varphi_0(x) = |x - x_0|^2, \quad \forall x \in \overline{\Omega}. \end{aligned}$$

Then, there exist $\sigma_0 > 0$ and $C > 0$ such that, for all $\sigma \geq \sigma_0$ and for all $q \in C_0^2(\Omega)$,

$$\sigma^2 \sum_{|\alpha| \leq 2} \int_{\Omega} |\partial_x^\alpha q(x)|^2 e^{2\sigma\varphi_0(x)} dx \leq C \sum_{|\alpha| \leq 2} \int_{\Omega} |\partial_x^\alpha (\mathcal{R}(x)q(x))|^2 e^{2\sigma\varphi_0(x)} dx.$$

Since the coefficient \hat{p} satisfies the following first-order system:

$$h(0)\nabla \cdot (\hat{p}(x)(\nabla \bar{u}_0(x) + \nabla \bar{u}_0(x)^T)) = -\partial_t w^*(x, 0),$$

we can apply lemma 2 to each equation of the system with

$$\begin{aligned} q(x) &= \hat{p}(x), \quad \mathcal{R}(x)q(x) = -\partial_t w^*(x, 0), \quad \varphi_0(x) = \varphi(x, 0), \\ a_0(x) &= h(0)\nabla \cdot (\nabla \bar{u}_0(x) + \nabla \bar{u}_0(x)^T), \quad a(x) = h(0)(\nabla \bar{u}_0(x) + \nabla \bar{u}_0(x)^T), \end{aligned}$$

which satisfy (H5) and we obtain then

$$\sum_{|\alpha| \leq 2} \int_{\Omega(\varepsilon)} |\partial_x^\alpha \hat{p}(x)|^2 e^{2\sigma\varphi(x,0)} dx \leq C \sum_{|\alpha| \leq 2} \sigma^{2(1-|\alpha|)} \int_{\Omega(\varepsilon)} |\partial_x^\alpha (\partial_t w^*)(x, 0)|^2 e^{2\sigma\varphi(x,0)} dx.$$

We supposed in (H3) that we know p in a neighborhood ω of the boundary $\partial\Omega$, i.e. $\hat{p} = 0$ in ω and so we can assume ε to be sufficiently small such that $\Omega \setminus \Omega(\varepsilon) \subset \omega$. This allows us to integrate on Ω on the left-hand side. Then, we have

$$\begin{aligned} \sum_{|\alpha| \leq 2} \int_{\Omega} |\partial_x^\alpha \hat{p}(x)|^2 e^{2\sigma\varphi(x,0)} dx \\ \leq C \left(\sum_{|\alpha| \leq 2} \|(\partial_x^\alpha \hat{p}) e^{\sigma\varphi}\|_{L^2(Q)}^2 + e^{C\sigma} \|\hat{u}\|_{H^6(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,\delta))}^2 + \sigma^3 e^{2\sigma\delta} \|\hat{u}\|_{H^6(Q)}^2 \right). \end{aligned} \tag{10}$$

We can absorb the first term of the right-hand side of (10) on the left-hand side, thanks to the Carleman weights. Indeed,

$$\sum_{|\alpha| \leq 2} \|(\partial_x^\alpha \hat{p}) e^{\sigma \varphi}\|_{L^2(Q)}^2 = \sum_{|\alpha| \leq 2} \int_{\Omega} |\partial_x^\alpha \hat{p}(x)|^2 e^{2\sigma \varphi(x,0)} \left(\int_0^{t_x^0} e^{2\sigma(\varphi(x,t) - \varphi(x,0))} dt \right) dx,$$

with, $\forall x \in \Omega$:

$$\int_0^{t_x^0} e^{2\sigma(\varphi(x,t) - \varphi(x,0))} dt \leq \int_0^{+\infty} e^{-2\sigma \beta t^2} dt = \frac{1}{\sqrt{2\sigma \beta}} \int_0^{+\infty} e^{-z^2} dz = \frac{C}{\sqrt{\sigma}}.$$

Then,

$$\sum_{|\alpha| \leq 2} \|(\partial_x^\alpha \hat{p}) e^{\sigma \varphi}\|_{L^2(Q)}^2 \leq \frac{C}{\sqrt{\sigma}} \sum_{|\alpha| \leq 2} \int_{\Omega} |\partial_x^\alpha \hat{p}(x)|^2 e^{2\sigma \varphi(x,0)} dx.$$

Finally, we obtain

$$\begin{aligned} e^{2\sigma d_0^2} \|\hat{p}\|_{H^2(\Omega)}^2 &\leq \sum_{|\alpha| \leq 2} \int_{\Omega} |\partial_x^\alpha \hat{p}(x)|^2 e^{2\sigma \varphi(x,0)} dx \\ &\leq C(e^{C\sigma} \|\hat{u}\|_{H^6(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^2 + \sigma^3 e^{2\sigma \delta} \|\hat{u}\|_{H^6(Q)}^2), \end{aligned}$$

and thus, we conclude that

$$\|\hat{p}\|_{H^2(\Omega)}^2 \leq C(e^{C\sigma} \|\hat{u}\|_{H^6(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^2 + e^{-\sigma(d_0^2 - \delta)}),$$

with $l = d_0^2 - \delta > 0$, since $\sup_{\sigma} (\sigma^3 e^{-\sigma l}) < +\infty$ and since we assumed in (H1) that \hat{u} is in $W^{8,\infty}(Q)$. In order to have

$$e^{C\sigma} \|\hat{u}\|_{H^6(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^2 = e^{-\sigma l},$$

we propose to set

$$\sigma = -\frac{1}{l+C} \log(\|\hat{u}\|_{H^6(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^2). \quad (11)$$

We have $\sigma > 0$ if we suppose that $\|\hat{u}\|_{H^6(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^2 < 1$. Thus, the right-hand side of the equation can be written as follows:

$$e^{C\sigma} \|\hat{u}\|_{H^6(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^2 + e^{-\sigma l} = 2 \left(\|\hat{u}\|_{H^6(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^2 \right)^{\frac{1}{l+C}}.$$

We now use the following interpolation result.

Lemma 3 (Proposition 4 in [27]). *For all $m \in \mathbb{N}^*$, there exists a constant $C > 0$ such that, for all r satisfying $0 \leq r \leq m$ and for all $u \in H^m(Q)$, we have*

$$\|u\|_{H^r(Q)} \leq C \|u\|_{L^2(Q)}^{1-\frac{r}{m}} \|u\|_{H^m(Q)}^{\frac{r}{m}}.$$

We apply lemma 3 to the second derivative of $\hat{u} \in H^8(Q)$ with $m = 6$ and $r = 2$ in order to write

$$\|\hat{u}\|_{H^6(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))} \leq C \|\hat{u}\|_{H^2(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^{1/3} \|\hat{u}\|_{H^8(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^{2/3}.$$

Hence,

$$\|\hat{p}\|_{H^2(\Omega)} \leq C \left(\|\hat{u}\|_{H^2(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^{1/3} \right)^{\frac{l}{l+C}}. \quad (12)$$

So we already have a stability result with observation in $Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta)$ which is included in $(\Omega(\varepsilon) \setminus \Omega(2\varepsilon)) \times (0, T)$ if $T > \frac{d}{\sqrt{\beta}}$.

2.6. Use the unique continuation result

We can now apply theorem 3 to \hat{u} . We verify that hypothesis (H6) holds, i.e. that

$$R(x, t) = - \int_0^t h(s) \nabla \cdot (\hat{p}(x) (\nabla \bar{u}(x, t-s) + \nabla \bar{u}(x, t-s)^T)) \, ds$$

vanishes in $\Omega \setminus \Omega(2\varepsilon) \subset \omega$, and choose $\rho < \delta$. We deduce that if T is large enough, we have

$$\|\hat{u}\|_{H^2(Q(\varepsilon, \delta) \setminus Q(2\varepsilon, \delta))}^2 \leq C \left[\log \left(2 + \frac{C}{\sum_{|\alpha|=1}^2 \|\partial_x^\alpha \hat{u}\|_{L^2(\Gamma \times (0, 6T))}^2} \right) \right]^{-1}.$$

Therefore,

$$\|\hat{p}\|_{H^2(\Omega)} \leq C \left[\log \left(2 + \frac{C}{\sum_{|\alpha|=1}^2 \|\partial_x^\alpha \hat{u}\|_{L^2(\Gamma \times (0, 6T))}^2} \right) \right]^{-\frac{l}{6(l+C)}}.$$

We set $\kappa = \frac{l}{6(l+C)} \in (0, 1)$ and change $6T$ by T . This achieves to prove theorem 1.

3. Proof of theorem 2

This section is devoted to the proof of the Carleman estimate (theorem 2). The proof consists in decoupling the system of equations as in [21] by writing the equations satisfied by u , $\nabla \wedge u$ and $\nabla \cdot u$. Then, the idea is to introduce a change of variable as in [19] to reduce the problem to a scalar hyperbolic equation for which the Carleman estimate is well known (corollary 2). Finally, we come back to the initial variable by a series of inequalities assuming the regularity of the coefficients and using the fundamental lemma 1.

Let u be the solution of the system

$$\begin{cases} \mathcal{P}u(x, t) = F(x, t), & \forall (x, t) \in \Omega \times (0, +\infty), \\ u(x, 0) = 0 & \text{or} \quad \partial_t u(x, 0) = 0, & \forall x \in \Omega, \\ u(x, t) = 0, & \forall (x, t) \in \partial\Omega \times (0, +\infty), \end{cases} \quad (13)$$

with

$$\begin{aligned} \mathcal{P}u(x, t) = & \partial_t^2 u(x, t) - \nabla \cdot (\mu(x) (\nabla u(x, t) + \nabla u(x, t)^T) + \lambda(x) (\nabla \cdot u)(x, t) I) \\ & + \int_0^t \nabla \cdot (\tilde{\mu}(x, s) (\nabla u(x, t-s) + \nabla u(x, t-s)^T) \\ & + \tilde{\lambda}(x, s) (\nabla \cdot u)(x, t-s) I) \, ds. \end{aligned}$$

Let us assume that hypotheses (H1) and (H2) are satisfied. First we have to decouple the equations.

3.1. Decouple the system of equations

To this end, we develop the system (13) as follows:

$$\begin{aligned} \partial_t^2 u(x, t) - \mu(x) \Delta u(x, t) - (\mu + \lambda)(x) \nabla (\nabla \cdot u)(x, t) - (\nabla \cdot u)(x, t) \nabla \lambda(x) \\ - (\nabla u(x, t) + \nabla u(x, t)^T) \cdot \nabla \mu(x) + \int_0^t \tilde{\mu}(x, t-s) \Delta u(x, s) \, ds \\ + \int_0^t (\nabla \cdot u)(x, s) \nabla \tilde{\lambda}(x, t-s) \, ds + \int_0^t (\tilde{\mu} + \tilde{\lambda})(x, t-s) \nabla (\nabla \cdot u)(x, s) \, ds \end{aligned}$$

$$+ \int_0^t (\nabla u(x, s) + \nabla u(x, s)^T) \cdot \nabla \tilde{\mu}(x, t - s) \, ds = F(x, t),$$

$$\forall (x, t) \in \Omega \times (0, +\infty).$$

Note the change of variable in the convolution terms. Then, we take the curl and the divergence of this system. We introduce the vectors $u_1 = u$, $u_3 = \nabla \wedge u$ and the scalar $u_2 = \nabla \cdot u$ which satisfy then the following system of seven equations:

$$\begin{aligned} \partial_t^2 u_i(x, t) - q_i(x) \Delta u_i(x, t) + \int_0^t \tilde{q}_i(x, t - s) \Delta u_i(x, s) \, ds \\ = F_i(x, t) + A_i(u_1, u_2, u_3)(x, t), \quad \forall i \in \{1, 2, 3\}, \end{aligned} \quad (14)$$

where we set

$$\begin{aligned} q_1 = q_2 = \mu, \quad q_3 = \lambda + 2\mu, \quad \tilde{q}_1 = \tilde{q}_2 = \tilde{\mu}, \quad \tilde{q}_3 = \tilde{\lambda} + 2\tilde{\mu}, \\ F_1 = F, \quad F_2 = \nabla \wedge F \quad \text{and} \quad F_3 = \nabla \cdot F. \end{aligned}$$

Here, the coupling terms A_i are first-order integro-differential operators in x and t , with coefficients bounded in $\Omega \times (0, +\infty)$, according to hypothesis (H1). These equations are coupled only at order 1; therefore, we can apply the results known for the scalar equations. However, the problem is now that the trace of the functions $\nabla \cdot u$ and $\nabla \wedge u$ on $\partial\Omega$ are no longer defined. We need now to change the integro-differential hyperbolic equation (14) into a hyperbolic equation in order to apply the classical results.

3.2. Use a change of variable

We introduce the following change of variable:

$$\tilde{u}_i(x, t) = q_i(x) u_i(x, t) - \int_0^t \tilde{q}_i(x, t - s) u_i(x, s) \, ds, \quad \forall i \in \{1, 2, 3\}. \quad (15)$$

Then, we have, for all $i \in \{1, 2, 3\}$ and for all $(x, t) \in \Omega \times (0, +\infty)$,

$$\begin{aligned} \partial_t^2 \tilde{u}_i(x, t) = q_i(x) \partial_t^2 u_i(x, t) - \int_0^t \partial_t^2 \tilde{q}_i(x, t - s) u_i(x, s) \, ds \\ + \partial_t \tilde{q}_i(x, 0) u_i(x, t) + \tilde{q}_i(x, 0) \partial_t u_i(x, t) \end{aligned}$$

and

$$\begin{aligned} \Delta \tilde{u}_i(x, t) = q_i(x) \Delta u_i(x, t) - \int_0^t \tilde{q}_i(x, t - s) \Delta u_i(x, s) \, ds + 2 \nabla q_i(x) \cdot \nabla u_i(x, t) \\ + \Delta q_i(x) u_i(x, t) - 2 \int_0^t \nabla \tilde{q}_i(x, t - s) \cdot \nabla u_i(x, s) \, ds \\ - \int_0^t \Delta \tilde{q}_i(x, t - s) u_i(x, s) \, ds. \end{aligned}$$

Therefore, for all $i \in \{1, 2, 3\}$ and for all $(x, t) \in \Omega \times (0, +\infty)$, the \tilde{u}_i satisfy a hyperbolic system of the type

$$\partial_t^2 \tilde{u}_i(x, t) - q_i(x) \Delta \tilde{u}_i(x, t) = q_i(x) (F_i(x, t) + A_i(u_1, u_2, u_3)(x, t)) + L_i(u_i)(x, t),$$

where the L_i are first-order integro-differential operators, the coefficients of which are bounded in $\Omega \times (0, +\infty)$. We now need a Carleman estimate for a scalar hyperbolic equation.




Figure 5. Definition of the domain $\Omega_t(\varepsilon, \delta)$ and the time t_x^δ .

3.3. Use a Carleman estimate for a hyperbolic scalar equation

The equations we consider are valid in $(0, +\infty)$ and because of the presence of the integral term, we cannot extend the solution to $(-\infty, 0)$. That is why we cannot use a classical global Carleman estimate found in the literature. Therefore, we will start with the following point-wise Carleman estimate stated in $Q(0, \delta)$:

Lemma 4 (Theorem 2.2.4 in [20]). *Let $q \in C^2(\overline{\Omega})$ satisfy condition 1 and φ be defined by (4). Then, there exist $\beta_0 > 0$ and $\sigma_0 > 0$ such that, for all $\beta \leq \beta_0$, for all $\sigma \geq \sigma_0$, for all $\delta > 0$ and for all $u \in H^2(Q(0, \delta))$, we have*

$$(\sigma |\nabla_{x,t} u(x, t)|^2 + \sigma^3 |u(x, t)|^2) e^{2\sigma\varphi(x,t)} + \nabla \cdot U(x, t) + \partial_t V(x, t) \leq C |\partial_t^2 u(x, t) - q(x) \Delta u(x, t)|^2 e^{2\sigma\varphi(x,t)}, \quad \forall (x, t) \in Q(0, \delta). \quad (16)$$

Here, (U, V) is a vector-valued function and satisfies

$$|U(x, t)| + |V(x, t)| \leq C (\sigma |\nabla_{x,t} u(x, t)|^2 + \sigma^3 |u(x, t)|^2) e^{2\sigma\varphi(x,t)}, \quad \forall (x, t) \in Q(0, \delta).$$

Moreover, $V(x, 0) = 0$, $\forall x \in \Omega$, if $u(x, 0) = 0$ or $\partial_t u(x, 0) = 0$, $\forall x \in \Omega$.

From this lemma, we deduce the following Carleman estimate.

Corollary 2. *Let $q \in C^2(\overline{\Omega})$ satisfy condition 1 and φ be defined by (4), with $\beta > 0$ sufficiently small. Then, there exist $\beta_0 > 0$ and $\sigma_0 > 0$ such that, for all $\beta \leq \beta_0$, for all $\sigma \geq \sigma_0$, for all $\delta > 0$ and for all $u \in H^2(Q(0, \delta))$ satisfying $u(x, 0) = 0$ or $\partial_t u(x, 0) = 0$, $\forall x \in \Omega$, we have*

$$\sigma \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(0,\delta))}^2 \leq C (\|(\partial_t^2 u - q \Delta u) e^{\sigma\varphi}\|_{L^2(Q(0,\delta))}^2 + \sigma \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(\partial Q(0,\delta) \setminus (\Omega \times \{0\}))}^2).$$

Proof. We integrate (16) over $Q(0, \delta)$:

$$\begin{aligned} \int_{Q(0,\delta)} (\sigma |\nabla_{x,t} u(x, t)|^2 + \sigma^3 |u(x, t)|^2) e^{2\sigma\varphi(x,t)} dx dt &\leq - \int_{Q(0,\delta)} (\nabla \cdot U(x, t) - \partial_t V(x, t) \\ &\quad + C |\partial_t^2 u(x, t) - \mu(x) \delta u(x, t)|^2 e^{2\sigma\varphi(x,t)}) dx dt. \end{aligned}$$

We now define (see figure 5) $\forall t \geq 0$, $\Omega_t(\varepsilon, \delta) = Q(\varepsilon, \delta) \cap (\Omega \times \{t\})$ and

$$\forall x \in \Omega, \quad t_x^\delta = \begin{cases} \sqrt{\frac{|x-x_0|^2 - \delta}{\beta}} & \text{if } \delta < |x - x_0|^2, \\ 0 & \text{else.} \end{cases}$$




Figure 6. Definition of the cut-off function χ_2 .

We note that

$$\begin{aligned} \int_{Q(0,\delta)} \nabla \cdot U(x, t) \, dx \, dt &= \int_0^\infty \int_{\Omega_t(0,\delta)} \nabla \cdot U(x, t) \, dx \, dt = \int_0^\infty \int_{\partial \Omega_t(0,\delta)} U(x, t) \cdot n \, dx \, dt \\ &= \int_{\partial Q(0,\delta) \setminus (\Omega \times \{0\})} U(x, t) \cdot n \, dx \, dt \end{aligned}$$

and that

$$\begin{aligned} \int_{Q(0,\delta)} \partial_t V(x, t) \, dx \, dt &= \int_\Omega \int_0^{t_x^\delta} \partial_t V(x, t) \, dt \, dx = \int_\Omega (V(x, t_x^\delta) - V(x, 0)) \, dx \\ &= \int_{\partial Q(0,\delta) \setminus (\Omega \times \{0\})} V(x, t) \, dx \, dt. \end{aligned}$$

Thus,

$$\begin{aligned} - \int_{Q(0,\delta)} \nabla \cdot U(x, t) \, dx \, dt - \int_{Q(0,\delta)} \partial_t V(x, t) \, dx \, dt \\ \leq \int_{\partial Q(0,\delta) \setminus (\Omega \times \{0\})} (|U(x, t)| + |V(x, t)|) \, dx \, dt \\ \leq C \int_{\partial Q(0,\delta) \setminus (\Omega \times \{0\})} (\sigma |\nabla_{x,t} u(x, t)|^2 + \sigma^3 |u(x, t)|^2) e^{2\sigma \varphi(x,t)} \, dx \, dt. \end{aligned}$$

This achieves to prove the expected result. \square

For $\varepsilon > 0$ and $\delta > 0$, we introduce then the cut-off function $\chi_2 \in C^\infty(\mathbb{R}^3 \times \mathbb{R})$ which satisfies $0 \leq \chi_2 \leq 1$ and is such that

$$\chi_2(x, t) = \begin{cases} 1, & \text{if } (x, t) \in Q(2\varepsilon, 2\delta), \\ 0, & \text{if } (x, t) \in Q \setminus \overline{Q(\varepsilon, \delta)}. \end{cases} \quad (17)$$

Figure 6 illustrates this cut-off function. Then, we set

$$u_i^*(x, t) = \chi_2(x, t) \tilde{u}_i(x, t), \quad \forall i \in \{1, 2, 3\}, \quad \forall (x, t) \in \Omega \times (0, +\infty).$$

Thus, u_i^* satisfies the equation

$$\begin{aligned} \partial_t^2 u_i^*(x, t) - q_i(x) \Delta u_i^*(x, t) &= \chi(x, t) (\partial_t^2 \tilde{u}_i(x, t) - \mu(x) \Delta \tilde{u}_i(x, t)) \\ &\quad + 2\partial_t \chi(x, t) \partial_t \tilde{u}_i(x, t) + \tilde{u}_i(x, t) (\partial_t^2 \chi(x, t) - \mu(x) \Delta \chi(x, t)) \\ &\quad - 2\mu(x) \nabla \chi(x, t) \cdot \nabla \tilde{u}_i(x, t) \\ &= \chi(x, t) (q_i(x) (F_i(x, t) + A_i(u_1, u_2, u_3)(x, t)) + L_i(u_i)(x, t)) + \tilde{L}_i(\tilde{u}_i)(x, t), \end{aligned}$$

where \tilde{L}_i is a first-order integro-differential operator with coefficients bounded in $\Omega \times (0, +\infty)$. Since we supposed that $u(x, 0) = 0$ or $\partial_t u(x, 0) = 0$, $\forall x \in \Omega$, and that, according to (H2), the coefficient q_i satisfies condition 1, we can apply corollary 2 to u_i^* in $Q(0, \delta/2)$, i.e. there exists $\beta > 0$ sufficiently small such that, for $\sigma > 0$ sufficiently large, we have

$$\sigma \|u_i^* e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta))}^2 \leq C \|(\partial_t^2 u_i^* - q_i \Delta u_i^*) e^{\sigma\varphi}\|_{L^2(Q(\varepsilon,\delta))}^2,$$

without the boundary term since u_i^* is null in $Q \setminus \overline{Q(\varepsilon, \delta)}$.

3.4. Return to the initial variable u

3.4.1. Upper bound for the right-hand side. We take into account the fact that

- all the coefficients $\lambda, \mu, \tilde{\lambda}, \tilde{\mu}$ and χ are bounded in $\Omega \times (0, +\infty)$ according to (H1);
- all the operators A_i, L_i and \tilde{L}_i are of order inferior or equal to 1;
- lemma 1 holds

to calculate

$$\begin{aligned} \|(\partial_t^2 u_i^* - q_i \Delta u_i^*) e^{\sigma\varphi}\|_{L^2(Q(\varepsilon,\delta))}^2 &= \|(\chi(\mu(F_i + A_i(u_1, u_2, u_3)) + L_i(u_i)) \\ &\quad + \tilde{L}_i(\tilde{u}_i)) e^{\sigma\varphi}\|_{L^2(Q(\varepsilon,\delta))}^2. \end{aligned}$$

Then,

$$\sigma \|u_i^* e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta))}^2 \leq C \|F_i e^{\sigma\varphi}\|_{L^2(Q)}^2 + C \sum_{1 \leq j \leq 3} \|u_j e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta))}^2. \quad (18)$$

3.4.2. Lower bound for the left-hand side. We use again the change of variable (15), for all $(x, t) \in \Omega \times (0, +\infty)$:

$$u_i(x, t) = \frac{1}{q_i(x)} \tilde{u}_i(x, t) + \int_0^t \frac{\tilde{q}_i(x, t-s)}{q_i(x)} u_i(x, s) ds, \quad \forall i \in \{1, 2, 3\}.$$

Then, we can write, taking into account (H2) and thanks to lemma 1, $\forall i \in \{1, 2, 3\}$:


$$\begin{aligned} \int_{Q(\varepsilon,\delta)} |u_i(x, t)|^2 e^{2\sigma\varphi(x,t)} dx dt &\leq C \int_{Q(\varepsilon,\delta)} |\tilde{u}_i(x, t)|^2 e^{2\sigma\varphi(x,t)} dx dt \\ &\quad + \frac{C}{\sigma} \int_{Q(\varepsilon,\delta)} |u_i(x, t)|^2 e^{2\sigma\varphi(x,t)} dx dt. \end{aligned}$$

For σ sufficiently large, we obtain

$$\int_{Q(\varepsilon,\delta)} |u_i(x, t)|^2 e^{2\sigma\varphi(x,t)} dx dt \leq C \int_{Q(\varepsilon,\delta)} |\tilde{u}_i(x, t)|^2 e^{2\sigma\varphi(x,t)} dx dt.$$

In the same way,

$$\begin{aligned} \int_{Q(\varepsilon,\delta)} |\nabla_{x,t} u_i(x, t)|^2 e^{2\sigma\varphi(x,t)} dx dt \\ \leq C \int_{Q(\varepsilon,\delta)} (|\tilde{u}_i(x, t)|^2 + |\nabla_{x,t} \tilde{u}_i(x, t)|^2) e^{2\sigma\varphi(x,t)} dx dt \end{aligned}$$

Figure 7. Definition of the open \mathcal{O} .

which allows to write, $\forall i \in \{1, 2, 3\}$:

$$\|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta))}^2 \leq C \|\tilde{u}_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta))}^2. \quad (19)$$

In addition, $\chi_2 = 1$ sur $Q(2\varepsilon, 2\delta)$, so we have $\tilde{u}_i = u_i^*$. Hence, $\forall i \in \{1, 2, 3\}$,

$$\|\tilde{u}_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta))}^2 = \|u_i^* e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(2\varepsilon,2\delta))}^2 + \|\tilde{u}_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,2\delta))}^2. \quad (20)$$

Finally, using (18), (19) and (20) and again lemma 1, we obtain

$$\begin{aligned} \sum_{1 \leq i \leq 3} \sigma \|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta))}^2 \\ \leq C \sum_{1 \leq i \leq 3} \left(\|F_i e^{\sigma\varphi}\|_{L^2(Q)}^2 + \|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta))}^2 + \sigma \|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,2\delta))}^2 \right), \end{aligned}$$

and thanks to the Carleman weights, we absorb the second term of the right-hand side on the left-hand side. Finally, we need the following lemma.

Lemma 5. Let φ be defined by (4). There exist $C > 0$ and $\sigma_0 > 0$ such that, for all $\sigma \geq \sigma_0$ and for all $u \in H^2(Q)$, we have

$$\begin{aligned} \frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(2\varepsilon,2\delta))}^2 \\ \leq C \left(\sigma \|u e^{\sigma\varphi}\|_{H_x^{1,\sigma}(Q(\varepsilon,\delta))}^2 + \|\nabla(\nabla \cdot u) e^{\sigma\varphi}\|_{L^2(Q(\varepsilon,\delta))}^2 + \|\nabla(\nabla \wedge u) e^{\sigma\varphi}\|_{L^2(Q(\varepsilon,\delta))}^2 \right). \end{aligned}$$

Proof. With χ_2 defined in (17), we compute, for all $(x, t) \in Q$,

$$\begin{aligned} \Delta(\chi_2(x, t) u(x, t) e^{\sigma\varphi(x, t)}) &= 2(\sigma \chi_2(x, t) \nabla \varphi(x, t) + \nabla \chi_2(x, t)) \cdot \nabla u(x, t) e^{\sigma\varphi(x, t)} \\ &\quad + (\sigma^2 \chi_2(x, t) |\nabla \varphi(x, t)|^2 + \sigma \chi_2(x, t) \Delta \varphi(x, t) + \Delta \chi_2(x, t)) u(x, t) e^{\sigma\varphi(x, t)} \\ &\quad + (\sigma \nabla \varphi(x, t) \cdot \nabla \chi_2(x, t)) u(x, t) e^{\sigma\varphi(x, t)} + \chi_2(x, t) \Delta u(x, t) e^{\sigma\varphi(x, t)}. \end{aligned}$$

For all $t \geq 0$, we can find an open \mathcal{O} (see figure 7) with a smooth boundary (at least $C^{1,1}$) such that $\Omega_t(\varepsilon, \delta) \subset \mathcal{O} \subset \Omega_t(0, 0)$. Since, for t fixed, the function $\chi_2(\cdot, t) u(\cdot, t) e^{\sigma\varphi(\cdot, t)}$ is in $H^2(\mathcal{O}) \cap H_0^1(\mathcal{O})$, we can write

$$\|\chi_2(\cdot, t) u(\cdot, t) e^{\sigma\varphi(\cdot, t)}\|_{H^2(\mathcal{O})} \leq C \|\Delta(\chi_2(\cdot, t) u(\cdot, t) e^{\sigma\varphi(\cdot, t)})\|_{L^2(\mathcal{O})}. \quad (21)$$

If we integrate inequality (21) for $t \in (0, +\infty)$ and note that $\chi_1 = 0$ in $\mathcal{O} \setminus \Omega_t(\varepsilon, \delta)$, we obtain

$$\begin{aligned} \|\chi_2 u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 &\leq \|u e^{\sigma\varphi}\|_{L^2(0,+\infty;H^2(\Omega_t(\varepsilon,\delta)))}^2 + \sigma^2 \|\nabla(u e^{\sigma\varphi})\|_{L^2(Q(\varepsilon,\delta))}^2 + \sigma^4 \|u e^{\sigma\varphi}\|_{L^2(Q(\varepsilon,\delta))}^2 \\ &\leq C(\|e^{\sigma\varphi} \Delta u\|_{L^2(Q(\varepsilon,\delta))}^2 + \sigma^2 \|e^{\sigma\varphi} \nabla u\|_{L^2(Q(\varepsilon,\delta))}^2 + \sigma^4 \|u e^{\sigma\varphi}\|_{L^2(Q(\varepsilon,\delta))}^2). \end{aligned}$$

We use the classical formula

$$\Delta u = \nabla(\nabla \cdot u) - \nabla \wedge (\nabla \wedge u),$$

and the fact that $\chi_2 = 1$ in $Q(2\varepsilon, 2\delta)$ to write

$$\begin{aligned} \|u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(2\varepsilon,2\delta))}^2 &\leq \|\chi_2 u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 \\ &\leq C(\|e^{\sigma\varphi} \nabla(\nabla \cdot u)\|_{L^2(Q(\varepsilon,\delta))}^2 + \|e^{\sigma\varphi} \nabla(\nabla \wedge u)\|_{L^2(Q(\varepsilon,\delta))}^2 \\ &\quad + \sigma^2 \|u e^{\sigma\varphi}\|_{H_x^{1,\sigma}(Q(\varepsilon,\delta))}^2). \end{aligned}$$

This is the expected result. \square

Using this lemma, we can finally write

$$\begin{aligned} \frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(2\varepsilon,2\delta))}^2 &\leq C(\sigma \|u_1 e^{\sigma\varphi}\|_{H_x^{1,\sigma}(Q(\varepsilon,\delta))}^2 + \|\nabla u_2 e^{\sigma\varphi}\|_{L^2(Q(\varepsilon,\delta))}^2 + \|\nabla u_3 e^{\sigma\varphi}\|_{L^2(Q(\varepsilon,\delta))}^2) \\ &\leq C \sum_{1 \leq i \leq 3} \sigma \|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta))}^2 \\ &\leq C \sum_{1 \leq i \leq 3} (\|F_i e^{\sigma\varphi}\|_{L^2(Q)}^2 + \sigma \|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,2\delta))}^2); \end{aligned}$$

then

$$\begin{aligned} \frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 &= \frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(2\varepsilon,2\delta))}^2 + \frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,2\delta))}^2 \\ &\leq C \sum_{1 \leq i \leq 3} (\|F_i e^{\sigma\varphi}\|_{L^2(Q)}^2 + \sigma \|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,2\delta))}^2). \end{aligned}$$

Moreover, we have

$$\begin{aligned} \sum_{1 \leq i \leq 3} \sigma \|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,2\delta))}^2 &= \sum_{1 \leq i \leq 3} (\sigma \|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,\delta))}^2 + \sigma \|u_i e^{\sigma\varphi}\|_{H^{1,\sigma}(Q(\varepsilon,\delta) \setminus Q(\varepsilon,2\delta))}^2) \\ &\leq e^{C\sigma} \|u\|_{H^2(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,\delta))}^2 + \sigma^3 e^{2\sigma\delta} \|u\|_{H^2(Q)}^2, \end{aligned}$$

since

$$\varphi(x, t) \leq \delta, \quad \forall (x, t) \in Q(\varepsilon, \delta) \setminus Q(\varepsilon, 2\delta).$$

Thus

$$\begin{aligned} \frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H_x^{2,\sigma}(Q(\varepsilon,\delta))}^2 &\leq C(\|F e^{\sigma\varphi}\|_{L^2(Q)}^2 + \|(\nabla F) e^{\sigma\varphi}\|_{L^2(Q)}^2 \\ &\quad + e^{C\sigma} \|u\|_{H^2(Q(\varepsilon,\delta) \setminus Q(2\varepsilon,\delta))}^2 + \sigma^3 e^{2\sigma\delta} \|u\|_{H^2(Q)}^2), \end{aligned}$$

and this concludes the proof of theorem 2.




Figure 8. Definition of the neighborhood ω of $\partial\Omega$ and the cut-off function χ_3 .

4. Proof of theorem 3

In this section, we prove the unique continuation result of theorem 3. It consists in transforming the integro-differential hyperbolic system (1)–(2) into an elliptic one thanks to a FBI-type transform. Then, we show a Carleman estimate (theorem 4) for the resulting integro-differential elliptic operator, using the same techniques as in the proof of theorem 2. And we use this Carleman estimate to obtain interpolation inequalities which link the value of the solution in an interior domain near the boundary to the one on the boundary. Finally, we come back to the solution of the initial problem by a series of inequalities.

Let u be the solution of

$$\mathcal{P}u(x, t) = R(x, t), \quad \forall (x, t) \in \Omega \times (0, +\infty), \quad (22)$$

with null initial and boundary conditions and let us assume that (H1) and (H6) are satisfied.

4.1. Transform the hyperbolic system into an elliptic one

Let us choose $\varepsilon > 0$ such that $\Omega \setminus \Omega(3\varepsilon) \subset \omega$.

We introduce the cut-off function $\chi_3 \in C_0^\infty(\mathbb{R}^3)$ (see figure 8) which satisfies $0 \leq \chi_3 \leq 1$ and is such that

$$\chi_3(x) = \begin{cases} 0, & \text{if } x \in \Omega(4\varepsilon), \\ 1, & \text{if } x \in \Omega \setminus \overline{\Omega(3\varepsilon)}, \end{cases}$$

and we set

$$u^*(x, t) = \chi_3(x)u(x, t), \quad \forall (x, t) \in \Omega \times (0, +\infty).$$

Thus, the new variable u^* satisfies the following equation:

$$\begin{aligned} \mathcal{P}u^*(x, t) &= \chi_3(x) \mathcal{P}u(x, t) + [\mathcal{P}, \chi_3]u(x, t) \\ &\implies \partial_t^2 u^*(x, t) - \mathcal{L}(x)u^*(x, t) + \int_0^t \tilde{\mathcal{L}}(x, s)u^*(x, t-s) \, ds \\ &= [\mathcal{P}(x, t), \chi_3]u(x, t), \end{aligned} \quad (23)$$

since, according to (H6), $R(x, t)$ is equal to zero in ω . We have introduced the following operators:

$$\begin{aligned} \mathcal{L}(x)\star &= \nabla \cdot (\mu(x)(\nabla \star + \nabla \star^T) + \lambda(x)(\nabla \cdot \star)I), \\ \tilde{\mathcal{L}}(x, t)\star &= \nabla \cdot (p(x)h(t)(\nabla \star + \nabla \star^T) + \tilde{\lambda}(x, t)(\nabla \cdot \star)I). \end{aligned}$$




Figure 9. Definition of the cut-off function θ .

We define a special transformation which is inspired from the classical FBI transform [25]:

Definition 3. Let $T > 1$ and $0 < \eta < \frac{1}{6}$. We introduce $\tilde{Q} = \Omega \times (-\frac{T}{2}, \frac{5T}{2}) \times (-3\eta, 3\eta)$. Then, $\forall (x, t, r) \in \tilde{Q}$, we define

$$(\mathcal{F}_\gamma u^*)(x, t, r) = \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} \theta(t-y) u^*(x, y) dy,$$

where $\theta \in C^\infty(\mathbb{R})$, $0 \leq \theta \leq 1$,

$$\theta(z) = \begin{cases} 1, & \text{if } z > -\frac{T}{2} + 3\eta, \\ 0, & \text{if } z < -\frac{T}{2} + \frac{3\eta}{2}. \end{cases}$$

This function is plotted in figure 9.

Thanks to this transform, we will convert locally the hyperbolic system (23) into an elliptic system. Firstly, we note that, $\forall (x, t, r) \in \tilde{Q}$,

$$\begin{aligned} \partial_r(\mathcal{F}_\gamma u^*)(x, t, r) &= \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} -i\gamma(t+ir-y) e^{-\frac{\gamma}{2}(t+ir-y)^2} \theta(t-y) u^*(x, y) dy \\ &= \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} -i\partial_y(e^{-\frac{\gamma}{2}(t+ir-y)^2}) \theta(t-y) u^*(x, y) dy \\ &= \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} i e^{-\frac{\gamma}{2}(t+ir-y)^2} \partial_y(\theta(t-y) u^*(x, y)) dy. \end{aligned}$$

The boundary terms of the integration by parts vanish because $u^*(x, 0) = 0$ and $e^{-\frac{\gamma}{2}(t+ir-y)^2} \rightarrow 0$ if $y \rightarrow \infty$. Moreover, $\theta(t-y) = 0$ if $y \geq T$. Then,

$$\begin{aligned} \partial_r^2(\mathcal{F}_\gamma u^*)(x, t, r) &= -\sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} \partial_y^2(\theta(t-y) u^*(x, y)) dy \\ &= -(\mathcal{F}_\gamma(\partial_y^2 u^*))(x, t, r) - F_\gamma(x, t, r), \quad \forall (x, t, r) \in \tilde{Q}, \end{aligned}$$

where

$$F_\gamma(x, t, r) = \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} (\theta''(t-y) u^*(x, y) - 2\theta'(t-y) \partial_y u^*(x, y)) dy.$$

Similarly, we can write

$$\partial_t^2(\mathcal{F}_\gamma u^*)(x, t, r) = (\mathcal{F}_\gamma(\partial_y^2 u^*))(x, t, r), \quad \forall (x, t, r) \in \tilde{Q},$$

and it is easy to see that

$$\mathcal{L}(x)(\mathcal{F}_\gamma u^*)(x, t, r) = (\mathcal{F}_\gamma(\mathcal{L}(x)u^*))(x, t, r), \quad \forall (x, t, r) \in \tilde{Q}.$$

We now have to look at the integral term

$$\begin{aligned} \mathcal{F}_\gamma \left(\int_0^y \tilde{\mathcal{L}}(x, s) u^*(x, y-s) ds \right) (x, t, r) \\ &= \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} \theta(t-y) \int_0^y \tilde{\mathcal{L}}(x, s) u^*(x, y-s) ds dy \\ &= \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} \tilde{\mathcal{L}}(x, s) \int_s^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} \theta(t-y) u^*(x, y-s) dy ds \\ &= \int_0^{+\infty} \tilde{\mathcal{L}}(x, s) \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} e^{-\frac{\gamma}{2}(t-s+ir-z)^2} \theta(t-s-z) u^*(x, z) dz ds \\ &= \int_0^{+\infty} \tilde{\mathcal{L}}(x, s) (\mathcal{F}_\gamma u^*)(x, t-s, r) ds = \int_{-T/2}^t \tilde{\mathcal{L}}(x, t-s) (\mathcal{F}_\gamma u^*)(x, s, r) ds. \end{aligned}$$

The main advantage of this transformation with respect to the FBI transform is that it transforms the convolution product of two functions into the convolution of the first function by the transform of the second function. The function $u_\gamma = \mathcal{F}_\gamma u^*$, function of three variables (x, t, r) , then satisfies a system of elliptic integro-differential equations, $\forall (x, t, r) \in \tilde{Q}$:

$$\begin{aligned} \mathcal{Q}u_\gamma(x, t, r) &= -2\partial_r^2 u_\gamma(x, t, r) - \partial_t^2 u_\gamma(x, t, r) - \mathcal{L}(x)u_\gamma(x, t, r) \\ &\quad + \int_{-T/2}^t \tilde{\mathcal{L}}(x, t-s) u_\gamma(x, s, r) ds = 2F_\gamma(x, t, s) + G_\gamma(x, t, s), \end{aligned} \quad (24)$$

where

$$G_\gamma(x, t, r) = \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} \theta(t-y) [\mathcal{P}(x, t), \chi_3] \hat{u}(x, y) dy,$$

with the boundary conditions:

$$\begin{cases} u_\gamma(x, t, r) = 0, & \forall (x, t, r) \in \partial\Omega \times \left(-\frac{T}{2}, \frac{5T}{2}\right) \times (-3\eta, 3\eta), \\ u_\gamma(x, t, r) = 0, & \forall (x, t, r) \in \Omega \times \left\{-\frac{T}{2}\right\} \times (-3\eta, 3\eta). \end{cases}$$

We note then that

$$G_\gamma(x, t, r) = 0, \quad \forall x \in \Omega \setminus \overline{\Omega(3\varepsilon)}, \quad (25)$$

because $[\mathcal{P}(x, t), \chi_3]$ only involves the derivatives of χ_3 and its support is in $\overline{\Omega(3\varepsilon)} \setminus \Omega(4\varepsilon)$. We also have

$$\|F_\gamma\|_{H^1(\tilde{Q})} \leq C e^{-m\gamma T} \|u^*\|_{H^2(\Omega \times (0, 3T))} \quad (26)$$

as well as

$$\|u_\gamma\|_{H^2(\tilde{Q})} \leq C e^{C\gamma} \|u^*\|_{H^2(\Omega \times (0, 3T))}, \quad (27)$$

where C depends on η, T, Ω but not on γ and where m is independent of T . Indeed,

$$\begin{aligned} \|F_\gamma\|_{L^2(\tilde{Q})}^2 &= \frac{\gamma}{2\pi} \int_{\tilde{Q}} \left| \int_0^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} (\theta''(t-y) u^*(x, y) \right. \\ &\quad \left. - 2\theta'(t-y) \partial_y u^*(x, y)) dy \right|^2 dr dt dx \end{aligned}$$

with

$$\begin{aligned}
 & \left| \int_0^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} (\theta''(t-y)u^*(x, y) - 2\theta'(t-y)\partial_y u^*(x, y)) dy \right|^2 \\
 & \leq \left(\int_{y \geq 0, -T/2+3\eta/2 \leq t-y \leq -T/2+3\eta} |e^{-\frac{\gamma}{2}(t+ir-y)^2}|^2 dy \right) \\
 & \quad \times \left(\int_0^{t+T/2} |\theta''(t-y)u^*(x, y) - 2\theta'(t-y)\partial_y u^*(x, y)|^2 dy \right) \\
 & \leq \left(\int_{y \geq 0, -T/2+3\eta/2 \leq t-y \leq -T/2+3\eta} e^{-\gamma((t-y)^2-r^2)} dy \right) C \|u^*(x, \cdot)\|_{H^1(0,3T)}^2 \\
 & \leq C e^{-m\gamma T} \|u^*(x, \cdot)\|_{H^1(0,3T)}^2, \quad \forall (x, t, r) \in \tilde{Q}.
 \end{aligned}$$

Indeed, $(t-y)^2 \geq (-\frac{T}{2} + 3\eta)^2$ and $r^2 \leq 9\eta^2$; therefore, $(t-y)^2 - r^2 \geq \frac{T}{4}(T - 6\eta)$. Thus, if we suppose $T > 1$, since $\eta < 1/6$, then there exists $m > 0$ independent of T such that $(t-y)^2 - r^2 \geq mT$. We can do the same work for the derivatives of F_γ . Likewise, we have

$$\|u_\gamma\|_{L^2(\tilde{Q})}^2 = \int_{\Omega} \int_{-T/2}^{5T/2} \int_{-3\eta}^{3\eta} \left| \sqrt{\frac{\gamma}{2\pi}} \int_0^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} \theta(t-y)u^*(x, y) dy \right|^2 dr dt dx$$

with, $\forall (x, t, r) \in \tilde{Q}$,

$$\begin{aligned}
 & \left| \int_0^{+\infty} e^{-\frac{\gamma}{2}(t+ir-y)^2} \theta(t-y)u^*(x, y) dy \right|^2 \\
 & \leq \left(\int_0^{t+T/2} |e^{-\frac{\gamma}{2}(t+ir-y)^2}|^2 dy \right) \left(\int_0^{t+T/2} |\theta(t-y)u^*(x, y)|^2 dy \right) \\
 & \leq \left(\int_0^{3T} e^{-\gamma((t-y)^2-r^2)} dy \right) C \|u^*(x, \cdot)\|_{L^2(0,3T)}^2 \leq C e^{C\gamma} \|u^*(x, \cdot)\|_{L^2(0,3T)}^2.
 \end{aligned}$$

And we use the same procedure for the derivatives of u_γ to obtain the desired result.

4.2. Prove a Carleman estimate

We write a Carleman estimate for the system of elliptic integro-differential equations (24), applying the same methods we used in the hyperbolic case. That is, we combine the decoupling of the equations proposed by [14] with the change of variables of [10], and we use a classical Carleman estimate for a scalar elliptic equation (lemma 2). The difference is that here we have three variables (x, t, r) , so the weight function has to be modified. Let us first introduce some notations:

- Domains: for $T > 0$ and $\eta > 0$, we note

$$\begin{aligned}
 \tilde{Q} &= \Omega \times \left(-\frac{T}{2}, \frac{5T}{2}\right) \times (-3\eta, 3\eta), \\
 \tilde{\Sigma} &= \partial\Omega \times \left(-\frac{T}{2}, \frac{5T}{2}\right) \times (-3\eta, 3\eta).
 \end{aligned}$$

- Norms

$$\begin{aligned}
 \|\star\|_{H^{k,\sigma}(\tilde{Q})}^2 &= \sum_{|\alpha| \leq k} \sigma^{2(k-|\alpha|)} \|\partial^\alpha \star\|_{L^2(\tilde{Q})}^2, \\
 \|\star\|_{H^{k,\sigma}(\tilde{\Sigma})}^2 &= \sum_{|\alpha| \leq k} \sigma^{2(k-|\alpha|)} \|\partial^\alpha \star\|_{L^2(\tilde{\Sigma})}^2.
 \end{aligned}$$

Definition 4 (A Carleman weight function). Let $x_0 \in \mathbb{R}^3 \setminus \overline{\Omega}$ and $\xi > 0$. We introduce a function ψ in the following way:

$$\psi(x, t, r) = |x - x_0|^2 + \left(t + \frac{T}{2}\right)^2 + r^2, \quad \forall (x, t, r) \in \tilde{Q},$$

and we set

$$\varphi(x, t, r) = e^{-\xi \psi(x, t, r)}, \quad \forall (x, t, r) \in \tilde{Q}. \quad (28)$$

Because we introduced this new weight function, we must now check that we still have a result, similar to one of lemma 1, to bound the integral terms in the proof of the Carleman estimate. It is given by the following lemma.

Lemma 6. Let φ be defined by (28), $\sigma > 0$ and $u \in L^2(\tilde{Q})$. Then, there exists a constant $C > 0$, independent of σ , such that

$$\int_{\tilde{Q}} \left(\int_{-T/2}^t |u(x, s, r)| \, ds \right)^2 e^{2\sigma \varphi(x, t, r)} \, dx \, dt \, dr \leq \frac{C}{\sigma} \int_{\tilde{Q}} |u(x, t, r)|^2 e^{2\sigma \varphi(x, t, r)} \, dx \, dt \, dr.$$

Proof. Using the Cauchy–Schwarz inequality, we can write

$$\begin{aligned} & \int_{\tilde{Q}} \left(\int_{-T/2}^t |u(x, s, r)| \, ds \right)^2 e^{2\sigma \varphi(x, t, r)} \, dx \, dt \, dr \\ & \leq \int_{\tilde{Q}} \left(\int_{-T/2}^t |u(x, s, r)|^2 \, ds \right) \left(t + \frac{T}{2} \right) e^{2\sigma \varphi(x, t, r)} \, dx \, dt \, dr. \end{aligned}$$

We note then that, $\forall (x, t, r) \in \tilde{Q}$,

$$\left(t + \frac{T}{2} \right) e^{2\sigma \varphi(x, t, r)} \leq -\frac{e^{\xi \psi(x, t, r)}}{4\xi\sigma} \partial_t (e^{2\sigma \varphi(x, t, r)}) \leq -\frac{C}{\sigma} \partial_t (e^{2\sigma \varphi(x, t, r)}).$$

Therefore,

$$\begin{aligned} & \int_{\tilde{Q}} \left(\int_{-T/2}^t |u(x, s, r)| \, ds \right)^2 e^{2\sigma \varphi(x, t, r)} \, dx \, dt \, dr \\ & \leq \int_{\Omega} \int_{-3\eta}^{3\eta} \int_{-T/2}^{5T/2} -\frac{C}{\sigma} \left(\int_{-T/2}^t |u(x, s, r)|^2 \, ds \right) \partial_t (e^{2\sigma \varphi(x, t, r)}) \, dt \, dx \, dr \\ & \leq \frac{C}{\sigma} \int_{\Omega} \int_{-3\eta}^{3\eta} \left[\int_{-T/2}^{5T/2} |u(x, t, r)|^2 e^{2\sigma \varphi(x, t, r)} \, dt \, dx \, dr \right. \\ & \quad \left. - e^{2\sigma \varphi(x, 5T/2, r)} \left(\int_{-T/2}^{5T/2} |u(x, s, r)|^2 \, ds \right) \, dx \, dr \right]. \end{aligned}$$

Thus, we deduce the result. \square

We are now ready to state the Carleman estimate for the operator \mathcal{Q} .

Theorem 4 (Carleman estimate). Let \mathcal{Q} be the operator defined by (24). Let K be a compact set in $\overline{\Omega} \times (-\frac{T}{2}, \frac{5T}{2}) \times (-3\eta, 3\eta)$. We assume that

$$(H1) \quad (\lambda, \mu) \in C^2(\overline{\Omega})^2 \text{ and } (\tilde{\lambda}, \tilde{\mu}) \in C^2(\overline{\Omega} \times (0, +\infty))^2.$$

Then, there exists $\xi_0 > 0$ such that, for all $\xi \geq \xi_0$, there exists $\sigma_0 > 0$ such that, for all $\sigma \geq \sigma_0$ and for all $u \in C_0^\infty(K)^3$, we have the following estimate:

$$\frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{Q})}^2 \leq C(\|(Qu) e^{\sigma\varphi}\|_{H^1(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{\Sigma})}^2),$$

where φ is defined by (28), and $C > 0$ depends on the $C^2(\overline{Q})$ -norm of the coefficient $\tilde{\mu}$ but is independent of σ .

Proof. Let K be a compact set in $\overline{\Omega} \times (-\frac{T}{2}, \frac{5T}{2}) \times (-3\eta, 3\eta)$, let $u \in C_0^\infty(K)^3$ be the solution of

$$Qu(x, t, r) = S(x, t, r), \quad \forall (x, t, r) \in \tilde{Q}, \quad (29)$$

and let us assume (H1). We must first decouple the equations. To this end, we take the divergence of (29) and we introduce the scalar $v = \nabla \cdot u$ and the vector $w = \nabla \wedge u$. We obtain the following system of seven scalar equations only coupled at the first order:

$$\begin{aligned} -2\partial_r^2 u(x, t, r) - \partial_t^2 u(x, t, r) - \mu(x)\Delta u(x, t, r) + \int_{-T/2}^t \tilde{\mu}(x, t-s)\Delta u(x, s, r) ds \\ = S(x, t, r) + A_1(u, v)(x, t, r), \quad \forall (x, t, r) \in \tilde{Q}, \\ -2\partial_r^2 v(x, t, r) - \partial_t^2 v(x, t, r) - (\lambda + 2\mu)(x)\Delta v(x, t, r) \\ + \int_{-T/2}^t (\tilde{\lambda} + 2\tilde{\mu})(x, t-s)\Delta v(x, s, r) ds \\ = (\nabla \cdot S)(x, t, r) + A_2(u, v, w)(x, t, r), \quad \forall (x, t, r) \in \tilde{Q}, \\ -2\partial_r^2 w(x, t, r) - \partial_t^2 w(x, t, r) - \mu(x)\Delta w(x, t, r) + \int_{-T/2}^t \tilde{\mu}(x, t-s)\Delta w(x, s, r) ds \\ = (\nabla \wedge S)(x, t, r) + A_3(u, v, w)(x, t, r), \quad \forall (x, t, r) \in \tilde{Q}, \end{aligned} \quad (30)$$

where A_1 , A_2 and A_3 are the first-order integro-differential operators. We only treat the first equation of (30); the other equations can be treated accordingly. We introduce the following change of variable:

$$\tilde{u}(x, t, r) = \mu(x)u(x, t, r) - \int_{-T/2}^t \tilde{\mu}(x, t-s)r u(x, s, r) ds, \quad \forall (x, t, r) \in \tilde{Q}.$$

Thus, \tilde{u} satisfies the classical elliptic equation

$$\begin{aligned} -2\partial_r^2 \tilde{u}(x, t, r) - \partial_t^2 \tilde{u}(x, t, r) - \mu(x)\Delta \tilde{u}(x, t, r) \\ = \mu(x)(S(x, t, r) + A_1(u, v)(x, t, r)) + L_1(u)(x, t, r), \end{aligned}$$

where L_1 is a first-order integro-differential operator with bounded coefficients. We now need a Carleman estimate for a scalar elliptic equation. It is given by the following lemma.

Lemma 7 ([28] and [29]). *Let Q be an open domain in \mathbb{R}^n , K be a compact set in \overline{Q} and ψ be a $C^\infty(Q)$ function satisfying $\nabla \psi(x) \neq 0$, $\forall x \in K$. Let*

$$\varphi(x) = e^{-\xi\psi(x)}, \quad \forall x \in Q,$$

where $\xi > 0$ is sufficiently large. We consider the scalar second-order elliptic operator

$$\mathcal{R}(x)\star = a(x) : \nabla^2 \star + b(x) \cdot \nabla \star + c\star, \quad \forall x \in Q,$$

where all the coefficients are $C^2(\bar{Q})$. Then, there exist $\sigma_0 > 0$ and $C > 0$ such that, for all $\sigma \geq \sigma_0$ and for all $u \in C_0^\infty(K)$, the following Carleman estimates hold true:

$$\begin{aligned} \frac{C}{\sigma} \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(Q)}^2 &\leq \|(\mathcal{R}u) e^{\sigma\varphi}\|_{L^2(Q)}^2 + \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\partial Q)}^2, \\ C\sigma \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(Q)}^2 &\leq \|(\mathcal{R}u) e^{\sigma\varphi}\|_{L^2(Q)}^2 + \sigma \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(\partial Q)}^2. \end{aligned}$$

As the weight function φ defined in (28) satisfies the hypothesis of lemma 7, we can apply the first inequality to \tilde{u} :

$$\frac{1}{\sigma} \|\tilde{u} e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{Q})}^2 \leq C \left(\|(-2\partial_r^2 \tilde{u} - \partial_t^2 \tilde{u} - \mu \Delta \tilde{u}) e^{\sigma\varphi}\|_{L^2(\tilde{Q})}^2 + \|\tilde{u} e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{\Sigma})}^2 \right).$$

We take into account the fact that the coefficients λ , $\tilde{\lambda}$, μ , $\tilde{\mu}$ are bounded in Q , that lemma 6 holds and that all the terms of the operators A_1 and L_1 are of order less than or equal to 1. Then, we obtain

$$\frac{1}{\sigma} \|\tilde{u} e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{Q})}^2 \leq C \left(\|S e^{\sigma\varphi}\|_{L^2(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \|v e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{\Sigma})}^2 \right).$$

We use again the change of variable

$$u(x, t, r) = \frac{1}{\mu(x)} \tilde{u}(x, t, r) + \int_{-T/2}^t \frac{\tilde{\mu}(x, t-s)}{\mu(x)} u(x, s, r) ds, \quad \forall (x, t, r) \in \tilde{Q}.$$

Hence,

$$\begin{aligned} \int_{\tilde{Q}} |u(x, t, r)|^2 e^{2\sigma\varphi(x,t,r)} dx dt dr &\leq C \int_{\tilde{Q}} |\tilde{u}(x, t, r)|^2 e^{2\sigma\varphi(x,t,r)} dx dt dr \\ &\quad + \frac{C}{\sigma} \int_{\tilde{Q}} |u(x, t, r)|^2 e^{2\sigma\varphi(x,t,r)} dx dt dr. \end{aligned}$$

For σ sufficiently large, the second term is absorbed. We do the same for the derivatives of u , which leads to

$$\frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{Q})}^2 \leq C \frac{1}{\sigma} \|\tilde{u} e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{Q})}^2.$$

Finally,

$$\frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{Q})}^2 \leq C \left(\|S e^{\sigma\varphi}\|_{L^2(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \|v e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{\Sigma})}^2 \right).$$

Thanks to the Carleman weights, we absorb the second term of the right-hand side on the left-hand side. We do the same work for v and w but using the second Carleman inequality of lemma 7 in order to obtain

$$\begin{aligned} \sigma \|v e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 &\leq C \left(\|(\nabla \cdot S) e^{\sigma\varphi}\|_{L^2(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \|w e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \sigma \|v e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{\Sigma})}^2 \right) \\ \sigma \|w e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 &\leq C \left(\|(\nabla \wedge S) e^{\sigma\varphi}\|_{L^2(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \|v e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \sigma \|w e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{\Sigma})}^2 \right). \end{aligned}$$

Hence,

$$\sigma \|v e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 \leq C \left(\|(\nabla S) e^{\sigma\varphi}\|_{L^2(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \|w e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{\Sigma})}^2 + \sigma \|v e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{\Sigma})}^2 \right).$$

Finally,

$$\begin{aligned} \frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{Q})}^2 &\leq C \left(\|Qu e^{\sigma\varphi}\|_{L^2(\tilde{Q})}^2 + \frac{1}{\sigma} \|(\nabla Qu) e^{\sigma\varphi}\|_{L^2(\tilde{Q})}^2 \right. \\ &\quad \left. + \frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H^{1,\sigma}(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{\Sigma})}^2 \right). \end{aligned}$$




Figure 10. Definition of $x^{(0)}$ and η .

And thanks to the Carleman weights, we absorb the third term of the right-hand side in the left-hand side to obtain the result

$$\frac{1}{\sigma} \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{Q})}^2 \leq C (\|Qu e^{\sigma\varphi}\|_{H^1(\tilde{Q})}^2 + \|u e^{\sigma\varphi}\|_{H^{2,\sigma}(\tilde{\Sigma})}^2)$$

and this concludes the proof of theorem 4.

4.3. Obtain local estimations

We are now going to apply the Carleman estimate we proved in theorem 4 to the function u_γ . To ensure that its support is in a compact set of $\overline{\Omega} \times (-\frac{T}{2}, \frac{5T}{2}) \times (-3\eta, 3\eta)$, we multiply it by a cut-off function. A good choice of this cut-off function leads to two local estimations. Summing these estimations, we obtain the desired interpolation result. This work is inspired from [23] which uses some results of [24] but note that we have here an additional variable.

4.4. First estimation

Let $\Gamma \subset \partial\Omega$ be arbitrarily small. We begin by estimating u_γ in a ball, which is close to Γ . Let us choose $0 < \eta < \varepsilon$ and $x^{(0)} \in \mathbb{R}^3 \setminus \overline{\Omega}$ such that

$$\overline{B(x^{(0)}, \eta)} \cap \overline{\Omega} = \emptyset, \quad B(x^{(0)}, 2\eta) \cap \Omega \neq \emptyset, \quad B(x^{(0)}, 4\eta) \cap \partial\Omega \subset \Gamma.$$

Figure 10 illustrates this construction. We define then

$$\psi^{(0)}(x, t, r) = |x - x^{(0)}|^2 + \frac{\eta^2}{T^2} \left(t + \frac{T}{2}\right)^2 + r^2, \quad \forall (x, t, r) \in \tilde{Q},$$

and we set

$$\varphi^{(0)}(x, t, r) = e^{-\frac{\xi}{\eta^2} \psi^{(0)}(x, t, r)}, \quad \forall (x, t, r) \in \tilde{Q}.$$

As in figure 11, we introduce the cut-off function $\chi_4 \in C_0^\infty(\mathbb{R})$ such that

$$\chi_4(z) = \begin{cases} 0, & \text{if } z < \frac{1}{2}, z > 8, \\ 1, & \text{if } \frac{3}{4} < z < 7, \end{cases}$$

and we set

$$u_\gamma^*(x, t, r) = \chi_4\left(\frac{\psi^{(0)}}{\eta^2}\right) u_\gamma(x, t, r), \quad \forall (x, t, r) \in \tilde{Q}.$$




Figure 11. Definition of the cut-off function χ_4 .

Here, u_γ^* has a compact support in $\bar{\Omega} \times (-\frac{T}{2}, \frac{5T}{2}) \times (-3\eta, 3\eta)$ since

$$\psi^{(0)}(x, t, \pm 3\eta) > 10\eta^2 \implies u_\gamma^*(x, t, r) = 0 \quad \text{if } |r| \geq 3\eta,$$

$$\psi^{(0)}\left(x, \frac{5T}{2}, r\right) > 10\eta^2 \implies u_\gamma^*(x, t, r) = 0 \quad \text{if } t \geq \frac{5T}{2},$$

and

$$u_\gamma(x, t, r) = 0 \quad \text{if } t < -\frac{T}{2} + \frac{3\eta}{2}.$$

Therefore, we can apply the Carleman estimate we showed in theorem 4 to this function:

$$\frac{1}{\sigma} \|u_\gamma^* e^{\sigma\varphi^{(0)}}\|_{H^{2,\sigma}(\tilde{Q})}^2 \leq C (\|(\mathcal{Q}u_\gamma^*) e^{\sigma\varphi^{(0)}}\|_{H^1(\tilde{Q})}^2 + \|u_\gamma^* e^{\sigma\varphi^{(0)}}\|_{H^{2,\sigma}(\tilde{\Sigma})}^2).$$

The last norm is actually in $\tilde{\Gamma} = \Gamma \times (-\frac{T}{2}, \frac{5T}{2}) \times (-3\eta, 3\eta) \subset \tilde{\Sigma}$. Indeed, we supposed $B(x^{(0)}, 4\eta) \cap \partial\Omega \subset \Gamma$; therefore, if $x \in \partial\Omega \setminus \Gamma$, then $u_\gamma^*(x, t, r) = 0$. In addition, $\forall(x, t, r) \in \tilde{Q}$,

$$\begin{aligned} \mathcal{Q}u_\gamma^*(x, t, r) &= \chi_4\left(\frac{\psi^{(0)}}{\eta^2}\right) \mathcal{Q}u_\gamma(x, t, r) + \left[\mathcal{Q}, \chi_4\left(\frac{\psi^{(0)}}{\eta^2}\right)\right] u_\gamma(x, t, r) \\ &= \chi_4\left(\frac{\psi^{(0)}}{\eta^2}\right) (F_\gamma(x, t, r) + G_\gamma(x, t, r)) + \left[\mathcal{Q}, \chi_4\left(\frac{\psi^{(0)}}{\eta^2}\right)\right] u_\gamma(x, t, r) \\ &= \chi_4\left(\frac{\psi^{(0)}}{\eta^2}\right) F_\gamma(x, t, r) + \left[\mathcal{Q}, \chi_4\left(\frac{\psi^{(0)}}{\eta^2}\right)\right] u_\gamma(x, t, r), \end{aligned}$$

since we saw in (25) that $G_\gamma(x, t, r) = 0$, $\forall x \in \Omega \setminus \overline{\Omega(3\varepsilon)}$. And here, $\chi_4\left(\frac{\psi^{(0)}}{\eta^2}\right)$ is different from zero if $|x - x^{(0)}|^2 \leq 8\eta^2$, thus, in particular, if $|x - x^{(0)}| \leq 3\eta < 3\varepsilon$. Then, we can write

$$\begin{aligned} \frac{1}{\sigma} \|u_\gamma^* e^{\sigma\varphi^{(0)}}\|_{H^{2,\sigma}(\tilde{Q})}^2 &\leq C (\|(\mathcal{Q}u_\gamma^*) e^{\sigma\varphi^{(0)}}\|_{H^1(\tilde{Q})}^2 + \|u_\gamma^* e^{\sigma\varphi^{(0)}}\|_{H^{2,\sigma}(\tilde{\Gamma})}^2) \\ &\implies \frac{1}{\sigma} e^{2\sigma e^{-6\varepsilon}} \|u_\gamma\|_{H^{2,\sigma}(\{\eta^2 \leq \psi^{(0)} \leq 6\eta^2\} \cap \tilde{Q})}^2 \\ &\leq C (\sigma^2 e^{2\sigma e^{-7\varepsilon}} \|u_\gamma\|_{H^2(\tilde{Q})}^2 + \sigma^2 e^{2\sigma e^{-\frac{\varepsilon}{2}}} \|F_\gamma\|_{H^1(\tilde{Q})}^2 + \sigma^4 e^{2\sigma e^{-\frac{\varepsilon}{2}}} \|u_\gamma\|_{H^2(\tilde{\Gamma})}^2). \end{aligned}$$




Figure 12. Definition of $x^{(1)}$ and ρ .

As illustrated in figure 12, we choose then ρ and $x^{(1)}$ such that

$$\text{dist}(x^{(1)}, \partial\Omega) \geq 4\rho, \quad \tilde{B}_1 = B(x^{(1)}, \rho) \times \left(-\frac{T}{2}, \frac{T}{2}\right) \times (-\rho, \rho) \subset \{\eta^2 \leq \psi^{(0)} \leq 6\eta^2\}.$$

This choice is valid since, in \tilde{B}_1 , we have

$$\begin{aligned} \psi^{(0)}(x, t, r) &\leq |x - x^{(1)}|^2 + |x^{(1)} - x^{(0)}|^2 + \frac{\eta^2}{T^2} \left(t + \frac{T}{2}\right)^2 + r^2 \\ &\leq \rho^2 + |x^{(1)} - x^{(0)}|^2 + \eta^2 + \rho^2 \end{aligned}$$

and since $B(x^{(0)}, 2\eta) \cap \Omega \neq \emptyset$, we can choose $x^{(1)}$ such that $|x^{(1)} - x^{(0)}|^2 < 4\eta^2$ and $4\rho \leq \eta$ so that $\psi^{(0)}(x, t, r) \leq 6\eta^2$. Then, if σ is sufficiently large, we have

$$\|u_\gamma\|_{H^2(\tilde{B}_1)}^2 \leq C e^{-C_1\sigma} \|u_\gamma\|_{H^2(\tilde{Q})}^2 + C e^{C_2\sigma} (\|F_\gamma\|_{H^1(\tilde{Q})}^2 + \|u_\gamma\|_{H^2(\tilde{\Gamma})}^2).$$

We minimize this with respect to σ (cf [24]) to obtain, with $v_0 = \frac{C_2}{C_1+C_2}$,

$$\|u_\gamma\|_{H^2(\tilde{B}_1)} \leq C (\|u_\gamma\|_{H^2(\tilde{Q})})^{1-v_0} (\|F_\gamma\|_{H^1(\tilde{Q})} + \|u_\gamma\|_{H^2(\tilde{\Gamma})})^{v_0}. \quad (31)$$

4.5. Second estimation

Now, we extend the estimation in $B(x^{(1)}, \rho)$ into $\Omega(\varepsilon) \setminus \Omega(2\varepsilon)$. Let $B(x^{(j)}, \rho)$, $2 \leq j \leq N$, be a cover of $\Omega(\varepsilon) \setminus \Omega(2\varepsilon)$. We suppose that $x^{(j)}$ is such that for $2 \leq j \leq N$,

$$\text{dist}(x^{(j)}, \partial\Omega) \geq 4\rho \quad \text{and} \quad B(x^{(j+1)}, \rho) \subset B(x^{(j)}, 2\rho).$$

We introduce

$$\tilde{B}_j = B(x^{(j)}, \rho) \times \left(-\frac{T}{2}, \frac{T}{2}\right) \times (-\rho, \rho).$$

This construction is illustrated in figure 13. We define

$$\psi^{(j)}(x, t, r) = |x - x^{(j)}|^2 + \frac{\rho^2}{T^2} \left(t + \frac{T}{2}\right)^2 + r^2, \quad \forall (x, t, r) \in \tilde{Q},$$

and

$$\varphi^{(j)}(x, t, r) = e^{-\frac{\xi}{\rho^2} \psi^{(j)}(x, t, r)}, \quad \forall (x, t, r) \in \tilde{Q}.$$




Figure 13. Definition of the balls $B(x^{(j)}, \rho)$.

We set

$$u_\gamma^*(x, t, r) = \chi_4 \left(\frac{\psi^{(j)}}{\rho^2} \right) u_\gamma(x, t, r), \quad \forall (x, t, r) \in \tilde{Q},$$

to which we apply theorem 4, without the boundary term, since the support of u_γ^* is an interior domain:

$$\frac{1}{\sigma} \|u_\gamma^* e^{\sigma \varphi^{(j)}}\|_{H^{2,\sigma}(\tilde{Q})}^2 \leq C \|(\mathcal{Q}u_\gamma^*) e^{\sigma \varphi^{(j)}}\|_{H^1(\tilde{Q})}^2.$$

In addition, $\forall (x, t, r) \in \tilde{Q}$,

$$\mathcal{Q}u_\gamma^*(x, t, r) = \chi_4 \left(\frac{\psi^{(j)}}{\rho^2} \right) F_\gamma(x, t, r) + \left[\mathcal{Q}, \chi_4 \left(\frac{\psi^{(j)}}{\rho^2} \right) \right] u_\gamma(x, t, r).$$

Thus, we have

$$\begin{aligned} \frac{C}{\sigma} e^{2\sigma e^{-6\xi}} \|u_\gamma\|_{H^{2,\sigma}(\{\rho^2 \leq \psi^{(j)} \leq 6\rho^2\} \cap \tilde{Q})}^2 &\leq \sigma^2 e^{2\sigma e^{-\frac{\xi}{2}}} \|u_\gamma\|_{H^2(\psi^{(j)} \leq \rho^2)}^2 \\ &+ \sigma^2 e^{2\sigma e^{-7\xi}} \|u_\gamma\|_{H^2(\psi^{(j)} \leq 8\rho^2)}^2 + \sigma^2 e^{2\sigma e^{-\frac{\xi}{2}}} \|F_\gamma\|_{H^1(\tilde{Q})}^2, \end{aligned}$$

and, if we choose σ large, we can write

$$\begin{aligned} e^{2\sigma e^{-6\xi}} \|u_\gamma\|_{H^{2,\sigma}(\{\rho^2 \leq \psi^{(j)} \leq 6\rho^2\} \cap \tilde{Q})}^2 &\leq C e^{2\sigma e^{-\frac{\xi}{2}}} \|u_\gamma\|_{H^2(\psi^{(j)} \leq \rho^2)}^2 \\ &+ C e^{2\sigma e^{-\frac{13\xi}{2}}} \|u_\gamma\|_{H^2(\psi^{(j)} \leq 8\rho^2)}^2 + C e^{2\sigma e^{-\frac{\xi}{2}}} \|F_\gamma\|_{H^1(\tilde{Q})}^2. \end{aligned}$$

This allows us to conclude that

$$\|u_\gamma\|_{H^2(\tilde{B}_{j+1})}^2 \leq C e^{C_3\sigma} (\|u_\gamma\|_{H^2(\tilde{B}_j)}^2 + \|F_\gamma\|_{H^1(\tilde{Q})}^2) + C e^{-C_4\sigma} \|u_\gamma\|_{H^2(\tilde{Q})}^2,$$

since $\tilde{B}_{j+1} \subset \{\psi^{(j)} \leq 6\rho^2\}$ and $\{\psi^{(j)} \leq \rho^2\} \subset \tilde{B}_j$. We minimize this with respect to σ to obtain, with $v_1 = \frac{C_4}{C_3+C_4}$,

$$\|u_\gamma\|_{H^2(\tilde{B}_{j+1})} \leq C (\|u_\gamma\|_{H^2(\tilde{Q})})^{1-v_1} (\|F_\gamma\|_{H^1(\tilde{Q})} + \|u_\gamma\|_{H^2(\tilde{B}_j)})^{v_1}.$$

We use then the recurrence result of the following lemma.

Lemma 8 (Lemma 4 in [29]). *Let $\alpha_j > 0$ satisfying, for all $j \geq 0$,*

$$\alpha_j \leq B^{1-v}(\alpha_{j-1} + A)^v \quad \text{and} \quad \alpha_j \leq B$$

where $A > 0$, $B > 0$ and $v \in]0, 1[$. Then, for all $\mu \in]0, v^N[$, we have

$$\alpha_N \leq 2^{1/(1-v)} B^{1-\mu} (\alpha_0 + A)^\mu.$$

Here,

$$\alpha_j = \|u_\gamma\|_{H^2(\tilde{B}_{j+1})}, \quad A = \|F_\gamma\|_{H^1(\tilde{Q})}, \quad B = \|u_\gamma\|_{H^2(\tilde{Q})}^2.$$

Therefore, we obtain

$$\|u_\gamma\|_{H^2(\tilde{B}_n)} \leq C(\|u_\gamma\|_{H^2(\tilde{Q})})^{1-\nu} (\|F_\gamma\|_{H^1(\tilde{Q})} + \|u_\gamma\|_{H^2(\tilde{B}_1)})^\nu.$$

We apply the Young inequality and we obtain

$$\|u_\gamma\|_{H^2(\tilde{B}_n)} \leq \epsilon^q \|u_\gamma\|_{H^2(\tilde{Q})} + \epsilon^{-q'} (\|F_\gamma\|_{H^1(\tilde{Q})} + \|u_\gamma\|_{H^2(\tilde{B}_1)}),$$

where $q = \frac{1}{1-\nu}$ and $q' = \frac{1}{\nu}$. We use the estimations (26) and (27) on F_γ and u_γ to write

$$\|u_\gamma\|_{H^2(\tilde{B}_n)} \leq \epsilon^q e^{M\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))} + \epsilon^{-q'} (e^{-m\gamma T} \|u^*\|_{H^2(\Omega \times (0,3T))} + \|u_\gamma\|_{H^2(\tilde{B}_1)}).$$

We choose then $\epsilon = e^{-2M\gamma/q}$, so that

$$\|u_\gamma\|_{H^2(\tilde{B}_n)} \leq e^{-M\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))} + e^{-(mT-2M\frac{q'}{q})\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))} + e^{2M\gamma\frac{q'}{q}} \|u_\gamma\|_{H^2(\tilde{B}_1)}.$$

If we fix $T > T_n$ with

$$mT_n - 2M\frac{q'}{q} = M,$$

we obtain, with $\kappa = 2M\gamma\frac{q'}{q}$,

$$\|u_\gamma\|_{H^2(\tilde{B}_n)} \leq e^{-M\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))} + e^{\kappa\gamma} \|u_\gamma\|_{H^2(\tilde{B}_1)}.$$

We can do the same work on the first estimation (31), the one in \tilde{B}_1 . We apply the Young inequality and the estimations (26) and (27) to obtain

$$\|u_\gamma\|_{H^2(\tilde{B}_1)} \leq \epsilon^{q_0} e^{M\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))} + \epsilon^{-q'_0} (e^{-m\gamma T} \|u^*\|_{H^2(\Omega \times (0,3T))} + \|u_\gamma\|_{H^2(\tilde{\Gamma})})$$

where $q_0 = \frac{1}{1-\nu_0}$ and $q'_0 = \frac{1}{\nu_0}$. We choose then $\epsilon = e^{-(2M+\kappa)\gamma/q_0}$, so that

$$\begin{aligned} \|u_\gamma\|_{H^2(\tilde{B}_1)} &\leq e^{-(M+\kappa)\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))} \\ &\quad + e^{-(mT-(2M+\kappa)\frac{q'_0}{q_0})\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))} + e^{(2M+\kappa)\gamma\frac{q'_0}{q_0}} \|u_\gamma\|_{H^2(\tilde{\Gamma})}. \end{aligned}$$

If we fix $T > T_0$ with

$$mT_0 - (2M + \kappa)\frac{q'_0}{q_0} = M + \kappa,$$

we obtain

$$\|u_\gamma\|_{H^2(\tilde{B}_1)} \leq e^{-(M+\kappa)\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))} + e^{C\gamma} \|u_\gamma\|_{H^2(\tilde{\Gamma})}.$$

And then we regroup the two estimations in

$$\|u_\gamma\|_{H^2(\tilde{B}_n)} \leq e^{-M\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))} + e^{C\gamma} \|u^*\|_{H^2(\Gamma \times (0,3T))}^2.$$

Finally, we take $T > \max_n T_n$ and we sum the inequalities to obtain

$$\|u_\gamma\|_{H^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (-\frac{T}{2}, \frac{T}{2}) \times (-\rho, \rho))}^2 \leq e^{-C\gamma} \|u^*\|_{H^2(\Omega \times (0,3T))}^2 + e^{C\gamma} \|u^*\|_{H^2(\Gamma \times (0,3T))}^2.$$

We can now go back to the variable u in the estimations.

4.6. Return to the variable u

We introduce the new variable

$$\begin{aligned} w_\gamma(x, t) &= u_\gamma(x, t, r = 0) = \sqrt{\frac{\gamma}{2\pi}} \int_0^\infty e^{-\frac{\gamma}{2}(t-y)^2} \theta(t-y) u^*(x, y) dy \\ &= (K_\gamma * u^*)(x, t), \quad \forall (x, t) \in \Omega \times \mathbb{R}, \end{aligned}$$

with

$$K_\gamma(t) = \sqrt{\frac{\gamma}{2\pi}} e^{-\frac{\gamma}{2}t^2} \theta(t), \quad \forall t \in \mathbb{R}.$$

We see that w_γ converges to u^* when γ tends to infinity. We have

$$\begin{aligned} \|w_\gamma\|_{H^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2} - \rho))}^2 &\leq C \|u_\gamma\|_{H^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (-\frac{T}{2}, \frac{T}{2}) \times (-\rho, \rho))}^2 \\ &\leq e^{-C\gamma} \|u^*\|_{H^2(\Omega \times (0, 3T))}^2 + e^{C\gamma} \|u^*\|_{H^2(\Gamma \times (0, 3T))}^2. \end{aligned}$$

Indeed, the Cauchy formula states that, for $0 < d < \rho$, we have

$$w_\gamma(x, a) = \frac{1}{2i\pi} \int_{|w-a|=d} \frac{w_\gamma(x, w)}{w-a} dw,$$

which, using the polar coordinates, implies that

$$|w_\gamma(x, a)|^2 \leq C \int_0^{2\pi} |w_\gamma(x, a + d e^{i\theta})|^2 d\theta.$$

We integrate the above then, for $0 < d < \rho$, into

$$|w_\gamma(x, a)|^2 \leq \frac{C}{\rho} \int_0^\rho \int_0^{2\pi} |w_\gamma(x, a + d e^{i\theta})|^2 d\theta d\rho,$$

so that we can write

$$\begin{aligned} |w_\gamma(x, a)|^2 &\leq C \int_{-\rho}^\rho \int_{|t-a| \leq \rho} |w_\gamma(x, t + ir)|^2 dt dr \\ &= C \int_{-\rho}^\rho \int_{|t-a| \leq \rho} |u_\gamma(x, t, r)|^2 dt dr, \end{aligned}$$

according to the definition of w_γ . Then, we integrate for $x \in \Omega(\varepsilon) \setminus \Omega(2\varepsilon)$:

$$\begin{aligned} \|w_\gamma(\cdot, a)\|_{L^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon))}^2 &\leq C \int_{-\rho}^\rho \int_{|t-a| \leq \rho} \|u_\gamma(\cdot, t, r)\|_{L^2(\Omega(\varepsilon, 2\varepsilon))}^2 dt dr \\ &\leq C \|u_\gamma\|_{L^2(\Omega(\varepsilon, 2\varepsilon) \times (-\frac{T}{2}, \frac{T}{2}) \times (-\rho, \rho))}^2. \end{aligned}$$

We integrate for $a \in (0, \frac{T}{2} - \rho)$:

$$\|w_\gamma\|_{L^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2} - \rho))}^2 \leq C \|u_\gamma\|_{L^2(\Omega(\varepsilon, 2\varepsilon) \times (-\frac{T}{2}, \frac{T}{2}) \times (-\rho, \rho))}^2$$

and we do the same for the derivatives of w_γ . We can now come back to u^* (and u) using the classical Fourier transform (noted with a big hat symbol) because we note that

$$w_\gamma(x, t) = (K_\gamma * u^*)(x, t) \implies \widehat{w_\gamma}(x, \tau) = \widehat{K_\gamma}(\tau) \widehat{u^*}(x, \tau),$$

where

$$\begin{aligned} \widehat{K_\gamma}(\tau) &= \sqrt{\frac{\gamma}{2\pi}} \int_{-\infty}^{+\infty} e^{-i\tau t} e^{-\frac{\gamma}{2}t^2} \theta(t) dt \\ &= \sqrt{\frac{\gamma}{2\pi}} \int_{-\infty}^{+\infty} e^{-i\tau t} e^{-\frac{\gamma}{2}t^2} dt + \sqrt{\frac{\gamma}{2\pi}} \int_{-\infty}^{+\infty} e^{-i\tau t} e^{-\frac{\gamma}{2}t^2} (\theta(t) - 1) dt. \end{aligned}$$

We have

$$\begin{aligned}\sqrt{\frac{\gamma}{2\pi}} \int_{-\infty}^{+\infty} e^{-i\tau t} e^{-\frac{\gamma}{2}t^2} dt &= \sqrt{\frac{\gamma}{2\pi}} \int_{-\infty}^{+\infty} e^{-(\sqrt{\frac{\gamma}{2}}t + \sqrt{\frac{1}{2\gamma}}i\tau)^2} e^{-\frac{\tau^2}{2\gamma}} dt \\ &= \frac{1}{\sqrt{\pi}} e^{-\frac{\tau^2}{2\gamma}} \int_{-\infty}^{+\infty} e^{-z^2} dz = e^{-\frac{\tau^2}{2\gamma}}.\end{aligned}$$

Hence,

$$\widehat{K}_\gamma(\tau) = e^{-\frac{\tau^2}{2\gamma}} + \sqrt{\frac{\gamma}{2\pi}} \int_{-\infty}^{-\frac{T}{2}+3\eta} e^{-i\tau t} e^{-\frac{\gamma}{2}t^2} (\theta(t) - 1) dt.$$

Therefore, we have

$$\begin{aligned}|(1 - \widehat{K}_\gamma)(\tau)| &\leq |1 - e^{-\frac{\tau^2}{2\gamma}}| + \sqrt{\frac{\gamma}{2\pi}} \int_{-\infty}^{-\frac{T}{2}+3\eta} |e^{-i\tau t} e^{-\frac{\gamma}{2}t^2}| |\theta(t) - 1| dt \\ &\leq \frac{\tau^2}{2\gamma} + \sqrt{\frac{\gamma}{2\pi}} \int_{-\infty}^{-\frac{T}{2}+3\eta} e^{-\frac{\gamma}{2}t^2} dt,\end{aligned}$$

where we used for the first term the fact that the function $\frac{1-e^{-t^2}}{t^2}$ is bounded in \mathbb{R} . We win one order of convergence since we have $\frac{1}{\gamma}$ and no $\frac{1}{\sqrt{\gamma}}$ as in [14]. Then, we can introduce in the second term the change of variable $z = e^{-\frac{\gamma}{2}t^2}$, that is $t = \sqrt{-\frac{2}{\gamma} \ln(z)}$, and $dz = -\gamma t e^{-\frac{\gamma}{2}t^2} dt$, and so we have

$$\begin{aligned}\sqrt{\frac{\gamma}{2\pi}} \int_{-\infty}^{-\frac{T}{2}+3\eta} e^{-\frac{\gamma}{2}t^2} dt &\leq \frac{1}{2\sqrt{\pi}} \int_0^{e^{-\frac{\gamma}{2}(-\frac{T}{2}+3\eta)^2}} \frac{dz}{\sqrt{-\ln(z)}} \\ &\leq \frac{1}{2\sqrt{\pi}} \frac{e^{-\frac{\gamma}{2}(-\frac{T}{2}+3\eta)^2}}{\sqrt{\frac{\gamma}{2}(-\frac{T}{2}+3\eta)^2}} \leq \frac{C}{\gamma}.\end{aligned}$$

Then,

$$\begin{aligned}\|u^* - w_\gamma\|_{L^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2}-\rho))}^2 &\leq \|u^* - w_\gamma\|_{L^2(\Omega \times \mathbb{R})}^2 = \|\widehat{u^* - w_\gamma}\|_{L^2(\Omega \times \mathbb{R})}^2 \\ &\leq \int_{\Omega} \int_{-\infty}^{+\infty} \left| \frac{\tau^2 + C}{\gamma} \right|^2 |\widehat{u^*}(x, \tau)|^2 d\tau \\ &\leq \frac{C}{\gamma^2} \int_{\Omega} \int_{-\infty}^{+\infty} (|\widehat{u^*}(x, \tau)|^2 + |\partial_\tau^2 \widehat{u^*}(x, \tau)|^2) d\tau \leq \frac{C}{\gamma^2} \|\widehat{u^*}\|_{H^2(\Omega \times \mathbb{R})}^2.\end{aligned}$$

Finally, if we suppose that u^* is prolonged by zero outside $(0, 3T)$, we have

$$\|u^* - w_\gamma\|_{L^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2}-\rho))} \leq \frac{C}{\gamma} \|u^*\|_{H^2(\Omega \times (0, 3T))}.$$

Thus,

$$\begin{aligned}\|u^*\|_{H^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2}-\rho))} &\leq C(\|u^* - w_\gamma\|_{H^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2}-\rho))} + \|w_\gamma\|_{H^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2}-\rho))}) \\ &\leq \frac{C}{\gamma} \|u^*\|_{H^4(\Omega \times (0, 3T))} + C\|w_\gamma\|_{H^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2}-\rho))}.\end{aligned}$$

Coming back to u and taking into account that $u^*(x, t) = \chi_3(x)u(x, t)$, so that $u = u^*$ in $\Omega(\varepsilon) \setminus \Omega(2\varepsilon)$, we have

$$\|u\|_{H^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2}-\rho))} \leq \frac{C}{\gamma} \|u\|_{H^4(\Omega \times (0, 3T))} + e^{C\gamma} \|u\|_{H^2(\Gamma \times (0, 3T))}.$$

We use the fact that $\|u\|_{H^4(\Omega \times (0,3T))}$ is bounded and that

$$\|u\|_{H^2(\Gamma \times (0,3T))}^2 = \sum_{|\alpha|=1}^2 \|\partial_x^\alpha u\|_{L^2(\Gamma \times (0,3T))}^2,$$

since \hat{u} and its time derivatives vanish on Γ to write

$$\|u\|_{H^2(\Omega(\varepsilon) \setminus \Omega(2\varepsilon) \times (0, \frac{T}{2} - \rho))}^2 \leq \frac{C_5}{\gamma^2} + e^{C_6 \gamma} \sum_{|\alpha|=1}^2 \|\partial_x^\alpha u\|_{L^2(\Gamma \times (0,3T))}^2.$$

Then, we choose γ such that the first term dominates the second. We can for example choose

$$\gamma = \frac{1}{2C_6} \log \left(2 + \frac{C_5}{\sum_{|\alpha|=1}^2 \|\partial_x^\alpha u\|_{L^2(\Gamma \times (0,3T))}^2} \right).$$

We achieve the proof of theorem 3.

5. Numerical results

To illustrate the theoretical result (12), we present here one numerical example in 2D, even if the result is also valid in 3D. The numerical resolution that we propose leads to recover the values of the unknown coefficient p at each vertex of the discretized domain. Since the linear viscoelastic system (1) is a simplified model for the mechanical behavior of brain structures [30], we could use the method to retrieve the localization of a brain tumor. However, this example can be considered as an academic test case, sufficient to emphasize the motivation of the study, to illustrate the stability result and to propose an efficient numerical method.

5.1. Direct problem

We consider now the system (1)–(3), in two dimensions, with the following coefficients:

- $\mu(x) = \lambda(x) = 1200$, $h(t) = e^{-t/\tau}$ with $\tau = 1$, $\tilde{\lambda}(x, t) = 400 h(t)$,
- $\bar{p}(x) = \begin{cases} 400, & \text{in the healthy tissue,} \\ >400, & \text{in the tumor (cf figure 14),} \end{cases}$
- $f(x, t) = 0$, $\bar{u}_1(x) = 0$ and \bar{u}_0 is the solution of the stationary problem associated with (1):

$$\begin{cases} -\nabla \cdot (\mu(x)(\nabla \bar{u}_0(x) + \nabla \bar{u}_0(x)^T) + \lambda(x)(\nabla \cdot \bar{u}_0(x))I) = 1, & \forall x \in \Omega, \\ u_0(x) = 0, & \forall x \in \partial\Omega. \end{cases}$$

When dealing with a real experiment, it is difficult to ensure that the initial data \bar{u}_0 satisfy hypothesis (H5). In practice, the recuperation is carry out with $\bar{u}_0 \neq 0$. Moreover, we assume that \bar{p} is known in the boundary of Ω . This hypothesis, less restrictive than hypothesis (H3), is more likely to be acknowledged because experimental measurements of the coefficient on the boundary may be possible.

We can solve numerically the direct problem (1)–(3) by discretizing the equations

- in space using \mathbb{P}^1 Lagrange finite elements in the mesh shown in figure 14,
- in time using a θ -scheme with $\theta = 0.5$ (implicit centered scheme) and $\delta t = 1$,
- by using the trapezium formula for the integral term.




Figure 14. Unknown coefficient \bar{p} (on the left), computational mesh and initial data \bar{u}_0 (in the center), observation zone ω (on the right).

The time of observation is taken equal to $T = 50$ whereas the observation zone ω is the one shown in figure 14. A uniform relative error of 2% corresponding to the experimental error is added to the solution, i.e. the observed displacement measured experimentally u_{obs} is obtained by the formula

$$u_{\text{obs}}(x, t) = (1 + r \cdot 0.02)\bar{u}(x, t), \quad \forall (x, t) \in \omega \times (0, T),$$

where r is a constant in $[-1, 1]$, randomly chosen (according to a uniform distribution law) at each discretization point in the domain $\omega \times (0; T)$.

5.2. Inverse problem

We are looking for the minimizer of the non-quadratic functional

$$J(p) = \frac{1}{2} \int_0^T \int_{\omega} (|u - u_{\text{obs}}|^2 + |\nabla(u - u_{\text{obs}})|^2) \, dx \, dt,$$

with $u = \mathcal{M}(p)$, \mathcal{M} being the nonlinear operator from P to U associated with the system (1)–(3). In general, $u_{\text{obs}} \notin \mathcal{M}(P)$. Note that adding a Tikhonov regularizing term to J does not numerically help in this case since we will filter high frequencies when choosing the parameter space for p (see section 5.4). We solve the minimization problem by a Broyden–Fletcher–Goldfarb–Shanno (BFGS) algorithm [31]. Thus, we calculate

$$\begin{aligned} \nabla J(p, \delta p) &= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} (J(p + \varepsilon \delta p) - J(p)) \\ &= \int_0^T \int_{\omega} M_p \delta p \cdot ((u - u_{\text{obs}}) - \Delta(u - u_{\text{obs}})) \, dx \, dt, \end{aligned}$$

with M_p being the linearized operator of \mathcal{M} around p ; i.e. $M_p \delta p = \delta u$ satisfies

$$\begin{cases} \mathcal{P} \delta u(x, t) = - \int_0^t \nabla \cdot (\delta p(x) h(t-s)(\nabla u(x, s) + \nabla u(x, s)^T)) \, ds, \\ \delta u(x, 0) = 0, & \forall x \in \Omega, \\ \partial_t \delta u(x, 0) = 0, & \forall x \in \Omega, \\ \delta u(x, t) = 0, & \forall (x, t) \in \partial\Omega \times (0, T). \end{cases} \quad (32)$$

We introduce the adjoint operator \mathcal{P}^* of (2)

$$\begin{aligned} \mathcal{P}^* v(x, t) &= \partial_t^2 v(x, t) - \nabla \cdot (\mu(x)(\nabla v(x, t) + \nabla v(x, t)^T) + \lambda(x)(\nabla \cdot v(x, t))I) \\ &\quad + \int_t^T \nabla \cdot (p(x) h(s-t)(\nabla v(x, s) + \nabla v(x, s)^T) + \tilde{\lambda}(x, s-t)(\nabla \cdot v)(x, s)I) \, ds, \end{aligned}$$

and we define δu^* , the solution of

$$\begin{cases} \mathcal{P}^* \delta u^*(x, t) = \begin{cases} (u - u_{\text{obs}})(x, t) - \Delta(u - u_{\text{obs}})(x, t), & \forall x \in \omega, \\ 0, & \forall x \in (\Omega \setminus \omega), \end{cases} \\ \delta u^*(x, T) = 0, & \forall x \in \Omega, \\ \partial_t \delta u^*(x, T) = 0, & \forall x \in \Omega, \\ \delta u^*(x, t) = 0, & \forall (x, t) \in \partial\Omega \times (0, T). \end{cases}$$

Therefore, we can write

$$\begin{aligned} \nabla J(p, \delta p) &= \int_0^T \int_{\Omega} \delta u(x, t) \cdot \mathcal{P}^* \delta u^*(x, t) \, dx \, dt = \int_0^T \int_{\Omega} \mathcal{P} \delta u(x, t) \cdot \delta u^*(x, t) \, dx \, dt \\ &= \int_{\Omega} \delta p(x) \left(\int_0^T \int_0^t h(t-s) (\nabla u(x, s) + \nabla u(x, s)^T) : \nabla \delta u^*(x, t) \, ds \, dt \right) dx. \end{aligned}$$

5.3. Regularization method

We denote by p^* the numerical parameter and by \bar{p} the exact parameter, and we suppose p^* close to \bar{p} . We can write

$$\|p^* - \bar{p}\|_P = \|M_{\bar{p}}^{-1}(M_{\bar{p}} p^* - \bar{u})\|_P \leq \|M_{\bar{p}}^{-1}\| \|M_{\bar{p}} p^* - \bar{u}\|_U.$$

If $M_{\bar{p}}$, linear from P (of infinite dimension) to U , is compact for $\|\cdot\|_P$ and $\|\cdot\|_U$, then its inverse $M_{\bar{p}}^{-1}$ is not bounded. So, even if we find p^* such that $M_{\bar{p}} p^*$ is close to \bar{u} , we cannot guarantee that p^* will be close to \bar{p} . Thus, as a regularization method [26], we introduce the operator

$$R_K = \Pi_K M_{\bar{p}}^{-1},$$

where Π_K is the projection of P to a finite-dimensional space P_K . This operator R_K is bounded in U and verifies

$$R_K \longrightarrow M_{\bar{p}}^{-1} \quad \text{if } K \rightarrow +\infty,$$

and we now look for $p^* \in P_K$.

5.4. Mesh and basis adaptation

We choose to look for the unknown coefficient p^* in the space P_K of the K first eigenfunctions of the mesh, that is

$$P_K = \left\{ p \in P, p = \bar{p}|_{\partial\Omega} + \sum_{i=1}^K p_i \varphi_i \right\},$$

where $\bar{p}|_{\partial\Omega}$ is a raising of the trace of the exact value of p (which is known as we assumed \bar{p} known in ω) and

$$\begin{cases} -\Delta \varphi_i(x) = \sigma_i \varphi_i(x), & \forall x \in \Omega, \\ \varphi_i(x) = 0, & \forall x \in \partial\Omega. \end{cases}$$

We propose an adaptive method to solve the problem accurately. After computing a first solution p_0^* on the initial mesh, we use it to refine the mesh and to adapt the spectral basis. Indeed, we consider next as basis functions the solutions of the following problem:

$$\begin{cases} -\nabla \cdot (a(x) \nabla \varphi_i(x)) = \sigma_i \varphi_i(x), & \forall x \in \Omega, \\ \varphi_i(x) = 0, & \forall x \in \partial\Omega, \end{cases}$$

with

$$a(x) = \frac{1}{\nabla p_0^*(x)}.$$

The initial and adapted meshes and basis functions are shown in figures 15 and 16.




Figure 15. Initial mesh (left), initial basis functions n°1, n°2 and n°5 (right).




Figure 16. Adapted mesh (left), adapted basis functions n°1, n°2 and n°5 (right).




Figure 17. Relative error for the coefficient p in L^2 -norm with respect to K .

5.5. The results

We solve the inverse problem by the method introduced in sections 5.2–5.4. In figure 17, we plot the relative error between p^* and \bar{p} in L^2 -norm with respect to the number K of eigenfunctions in the basis and for different steps of the iterative process. In figure 18, we show the exact coefficient and the numerical result we obtain at different steps.




Figure 18. Exact coefficient \bar{p} (on the left), recovered coefficient p^* at step 0 with $K_0 = 50$ and $K_0 = 100$ (in the center), recovered coefficient p^* at step 2 with $K_0 = K_1 = K_2 = 50$ (on the right).

5.6. Perspectives

Note that we only try to recover the coefficient \bar{p} , assuming that all the other coefficients are known. Indeed, the aim of this example is to illustrate the main result of the paper, which is the theoretical stability for recovering the spatial part of only one viscoelastic coefficient from only one measurement. The other objective of this example is to show the interesting choice we can do for the spectral basis in an adaptive way. To recover simultaneously all the coefficients, there are various methods (see [32]). One method consists in adding to the minimal criteria more parameters, but this could be unstable if only a single observation is available, so we would need more independent observations. Another method consists in using an iterative procedure fixing all coefficients except one at each step. In both cases, we should start in a neighborhood of the real coefficients and play with noise levels. All this deep analysis of the numerical simulations is of course very relevant but it is out of the scope of the present publication. We are preparing a forthcoming publication that will include these numerical issues and more details about the choice of the spectral adaptive basis.

Acknowledgments

This work has been partially supported by the MathAmSud CIP-EDP project, by doctoral fellowships from CNRS and CONICYT for the first author and by the FONDECYT-CONICYT 1061263 grant for the second author.

References

- [1] Bellout H and Nečas J 1994 Existence of global weak solutions for a class of quasilinear hyperbolic integro-differential equations describing visco-elastic materials *Math. Ann.* **299** 275–91
- [2] Grasselli M 1994 Determining the relaxation tensor in linear viscoelasticity of integral type *Japan J. Ind. Appl. Math.* **11** 131–53
- [3] Janno J and von Wolfersdorf L 1997 Inverse problems for identification of memory kernels in viscoelasticity *Math. Methods Appl. Sci.* **20** 291–314

- [4] von Wolfersdorf L 1993 On identification of memory kernels in linear viscoelasticity *Math. Nachr.* **161** 203–17
- [5] Cavaterra C and Grasselli M 1997 On an inverse problem for a model of linear viscoelastic Kirchhoff plate *J. Integral Equ. Appl.* **9** 179–218
- [6] Lorenzi A 1999 A multidimensional identification problem related to a hyperbolic integrodifferential equation *Z. Anal. Anwendungen* **18** 407–35
- [7] Lorenzi A and Romanov V G 2006 Stability estimates for an inverse problem related to viscoelastic media *J. Inverse Ill-Posed Problems* **18** 407–35
- [8] Bukhgeim A L and Klibanov M V 1981 Global uniqueness of class of multidimensional inverse problems *Sov. Math. Dokl.* **24** 244–7
- [9] Carleman T 1939 Sur un problème d'unicité pour les systèmes d'équations aux dérivées partielles à deux variables indépendantes *Ark. Mat. Astr. Fys.* **2B** 1–9
- [10] Lorenzi A, Messina F and Romanov V G 2007 Recovering a Lamé kernel in a viscoelastic system *Appl. Anal.* **86** 1375–95
- [11] Lavrent'ev M M 1967 *Some Ill-Posed Problems of Mathematics Physics* (Berlin: Springer)
- [12] de Buhan M and Osses A 2009 Un résultat de stabilité pour la récupération d'un paramètre du système de la viscoélasticité 3D *C. R. Acad. Sci., Paris I* **347** 1373–8
- [13] Bellassoued M 2004 Global logarithmic stability in inverse hyperbolic problem by arbitrary boundary observation *Inverse Problems* **20** 1033–52
- [14] Bellassoued M, Imanuvilov O and Yamamoto M 2008 Inverse problem of determining the density and the Lamé coefficients by boundary data *SIAM J. Math. Anal.* **40** 238–65
- [15] Hörmander L 1963 *Linear Partial Differential Operators* (Berlin: Springer)
- [16] Isakov V 1998 *Inverse Problems for Partial Differential Equations* (Berlin: Springer)
- [17] Tataru D 1996 Carleman estimates and unique continuation for solutions to boundary value problems *J. Math. Pures Appl.* **75** 367–408
- [18] Imanuvilov O and Yamamoto M 2001 Global uniqueness and stability in determining coefficients of wave equations *Commun. Partial Differ. Equ.* **26** 1409–25
- [19] Cavaterra C, Lorenzi A and Yamamoto M 2006 A stability result via Carleman estimates for an inverse source problem related to a hyperbolic integro-differential equation *Comput. Appl. Math.* **25** 229–50
- [20] Klibanov M V and Timonov A 2004 *Carleman Estimates for Coefficient Inverse Problems and Numerical Applications* (Utrecht: VSP)
- [21] Imanuvilov O and Yamamoto M 2005 Carleman estimates for the three-dimensional nonstationary Lamé system and applications to an inverse problem *Control Theory of Partial Differential Equations (Lecture Notes in Pure and Applied Mathematics vol 242)* 337–74
- [22] Imanuvilov O Y 2002 On Carleman estimate for hyperbolic equations *Asymptotic Anal.* **32** 185–220
- [23] Bellassoued M 2001 Unicité et contrôle pour le système de Lamé *ESAIM: COCV* **6** 561–92
- [24] Robbiano L 1995 Fonction de coût et contrôle des solutions des équations hyperboliques *Asymptotic Anal.* **10** 95–115
- [25] Bros J and Iagolnitzer D 1975 Tuboïdes et structure analytique des distributions. II. Support essentiel et structure analytique des distributions *Séminaire Goulaouic–Lions–Schwartz 1974–1975: Equations aux dérivées partielles linéaires et non linéaires (Centre Math., Ecole Polytech.)* vol 18
- [26] Kirsch A 1991 *An Introduction to the Mathematical Theory of Inverse Problems (Applied Mathematical Sciences vol 120)* (Berlin: Springer)
- [27] Dautray R and Lions J L 1987 *Analyse mathématique et calcul numérique pour les sciences et les techniques* vol 3 (Paris: Masson)
- [28] Imanuvilov O and Puel J P 2003 Global Carleman estimates for weak solutions of elliptic nonhomogeneous Dirichlet problems *Int. Math. Res. Not.* **16** 883–913
- [29] Lebeau G and Robbiano L 1995 Contrôle exact de l'équation de la chaleur *Commun. Partial Differ. Equ.* **20** 335–56
- [30] Sarron J C, Blondeau C, Guillaume A and Osmont D 2000 Identification of linear viscoelastic constitutive models *J. Biomech.* **33** 685–93
- [31] Modercai A 2003 *Nonlinear Programming: Analysis and Methods* (New York: Dover)
- [32] Pagnacco E, Moreau A and Lemoss D 2007 Inverse strategies for the identification of elastic and viscoelastic material parameters using full-field measurements *Mater. Sci. Eng.* **452–453** 737–45