

Cubic differentials and finite volume convex projective surfaces*

Yves Benoist and Dominique Hulin

Abstract

We prove that there exists a natural bijection between the set of finite volume oriented convex projective surfaces with non-abelian fundamental group, and the set of finite volume hyperbolic Riemann surfaces endowed with a holomorphic cubic differential with poles of order at most 2 at the cusps.

1 Introduction

A projective structure \mathcal{A} on an n -dimensional manifold M is the data of a maximal atlas with values in the real projective space $\mathbb{R}P^n$ and with transition functions in the group $\mathrm{PGL}_{n+1}\mathbb{R}$ of projective transformations. The charts glue together to define the developing map $\mathrm{dev} : \widetilde{M} \rightarrow \mathbb{R}P^n$, where \widetilde{M} is the universal cover of M . The developing map is well defined up to $\mathrm{PGL}_{n+1}\mathbb{R}$ and satisfies, for any loop c in the fundamental group $\pi_1(M)$,

$$\mathrm{dev} \circ c = \mathrm{hol}(c) \circ \mathrm{dev}$$

where $\mathrm{hol} : \pi_1(M) \rightarrow \mathrm{PGL}_{n+1}\mathbb{R}$ is the holonomy representation.

The projective structure is said to be properly convex when the developing map is a diffeomorphism onto a properly convex domain $\Omega \subset \mathbb{R}P^n$, namely a domain that reads as a bounded convex open set in a suitable affine chart. The Hilbert distance on the properly convex subset $\Omega \subset \mathbb{R}P^n$ is defined by

$$d_\Omega(x, y) = |\log[x, y, a, b]|,$$

where $[x, y, a, b]$ denotes the cross ratio of the quadruple (x, y, a, b) , and a, b are the points of intersection of the line (xy) and the boundary $\partial\Omega$ of Ω . The Hilbert distance derives from a Finsler metric, which gives rise to a measure on Ω (section 2.2). This measure, being invariant under the holonomy group $\mathrm{hol}(\pi_1(M))$, induces a measure μ_F on M .

In this paper, we will mainly focus on projective surfaces S with finite Finsler volume, namely such that $\mu_F(S) < \infty$. When S is an oriented compact surface with negative Euler characteristic, François Labourie [10] and John Loftin [14] (see also CP Wang [22]) have parameterized the set of properly convex projective structures on S by the data of a complex structure J on S together with a holomorphic cubic differential U on (S, J) . The aim of this paper is to extend this result

*Geom. Topol. vol 17 (2013) pp. 595-620

to non compact projective surfaces with finite Finsler volume, hence positively answering the conjecture by Loftin in [13].

To an oriented surface S equipped with a properly convex projective structure \mathcal{A} , we associate the following geometric objects. Denote by $\text{dev} : \tilde{S} \rightarrow \Omega$ the developing map. Cheng–Yau’s solution of the Monge–Ampère equation on Ω produces an imbedding $\tilde{S} \hookrightarrow \mathbb{R}^3$ as an affine sphere (Theorem 2.5). Let h be the induced “affine” metric on S (Definition 2.2), J be the conformal class of h – which is a complex structure on S – and U be the Pick form (Definition 4.9) which is a holomorphic cubic differential on (S, J) .

Theorem 1.1 *Let S be an oriented surface with non-abelian fundamental group. The map $\mathcal{A} \rightarrow (J, U)$ is a bijection between :*

1. *the set of properly convex projective structures \mathcal{A} on S with finite Finsler volume*
2. *the set of hyperbolic Riemann surface structures J on S with finite volume together with a holomorphic cubic differential U on (S, J) with poles of order at most 2 at the cusps*

Remark 1.2 – *Theorem 1.1 was conjectured by Loftin in [13], under the condition that S has finite affine volume, instead of finite Finsler volume. In Proposition 2.6, we will prove these conditions to be equivalent by using Benzécri’s cocompactness theorem 2.7.*

– *Note that Marquis proved in [18] that, when $\pi_1(S)$ is not finitely generated, S does not admit any properly convex projective structure with finite Finsler volume.*

– *This bijection is a homeomorphism for the natural topologies on the two sets. Indeed, our construction ensures that the map $\mathcal{A} \rightarrow (J, U)$ is a continuous bijective map between two sets which are homeomorphic to a real vector space (see Marquis [17]).*

Let us review the main ingredients of the proof. In order to prove that the map $\mathcal{A} \rightarrow (J, U)$ is well defined we must first show that, when (S, \mathcal{A}) is assumed to have finite Finsler volume, the corresponding hyperbolic Riemann surface (S, J) also has finite volume. Then we must control the Pick form at the cusps. To this end we first remark that, since the affine metric h is complete (Corollary 3.5) and negatively pinched at infinity (Proposition 3.1), the generalized maximum principle of Yau ensures that h is conformally quasi-isometric to the hyperbolic metric h_0 on (S, J) (Proposition 5.1). Hence (S, h_0) has finite volume, thus the ends of (S, J) are parabolic (Corollary 5.4). The cocompactness theorem of Benzécri is then used again to provide estimates for the Pick form U , ensuring that the measure $|U|^{2/3}$ on S has finite mass : this is the reason why U will have poles of order no more than 2 at the cusps (Corollary 5.8).

The paper is organized as follows. In section 2, we mainly recall standard material from affine geometry and Monge–Ampère equations, and prove the equivalence of the finite Finsler volume condition and the finite affine volume condition. In section 3 we assume that S is non compact, and study the curvature of the affine metric at infinity. The Pick form of an affine sphere in \mathbb{R}^3 is defined in section 4, and the estimates for the Pick form that ensure that the map $\mathcal{A} \rightarrow (J, U)$ in our main theorem 1.1 is well defined are given in section 5. In the final section 6, we prove that this map is a bijection.

A right inverse map $(J, U) \rightarrow \mathcal{A}$ as in Theorem 1.1 was constructed by Loftin in [13]. This construction was the motivation for Loftin to state his conjecture. (See also [15], where Loftin extends this construction to allow cubic differentials U with poles of order at most 3.)

Acknowledgements. This work was completed while both authors were visiting the MSRI in Berkeley. We thank this institution for its support and hospitality. We also thank Alessio Figalli and Ludovic Marquis for interesting discussions on this topic.

2 Affine spheres and the real Monge–Ampère equation

In this section, we recall the definition of an affine sphere (references are for example Loftin [16] or Nomizu and Sasaki [19]) and the fundamental existence and uniqueness theorem by Cheng and Yau. We then derive from Benzécri’s cocompactness theorem the equivalence of the finite Finsler volume condition and of the finite affine volume condition (Proposition 2.6).

2.1 Affine spheres

Let $M \subset \mathbb{R}^{n+1}$ be a hypersurface and $E = M \times \mathbb{R}^{n+1}$ be the trivial vector bundle of rank $n + 1$ over M . The standard affine (flat) connection on \mathbb{R}^{n+1} induces a flat connection ∇ on E . Each choice of a transverse vector field $\xi : M \rightarrow \mathbb{R}^{n+1}$ yields a decomposition $E = TM \oplus L$, where L stands for the trivial line bundle over M spanned by ξ . It also yields a decomposition

$$\begin{cases} \nabla_X Y = D_X Y + h(X, Y) \xi \in TM \oplus L \\ \nabla_X \xi = -S(X) + \tau(X) \xi \in TM \oplus L \end{cases} \quad (2.1)$$

where X and Y are tangent vector fields. Observe that D is a torsion-free connection on TM , h is a symmetric 2-form on TM , S is an endomorphism of TM and τ is a 1-form on TM .

Recall that an hypersurface $M \subset \mathbb{R}^{n+1}$ is locally strictly convex when it can locally be written in an affine coordinate system as the graph of a function with positive definite hessian.

Proposition 2.1 *Assume that the hypersurface $M \subset \mathbb{R}^{n+1}$ is locally strictly convex. Then, there exists a unique transverse vector field ξ for which :*

1. $\tau = 0$
2. h is positive definite
3. $|\det(Y_1, \dots, Y_n, \xi)| = 1$ for any h -orthonormal frame (Y_1, \dots, Y_n) of TM

Definition 2.2 *This vector field ξ is the “affine normal” of M . The corresponding connection D on TM is the “Blaschke connection”, the metric h is the “affine metric” on M and the endomorphism $S : TM \rightarrow TM$ is the “affine shape operator”. They are invariant under the action of the group $SI_{n+1}^{\pm} \mathbb{R}$ of real matrices with determinant ± 1 .*

Proposition 2.1 will be a consequence of the following.

Lemma 2.3 *Let ξ and ξ' be two transverse vector fields on M , respectively associated with (h, S, τ) and (h', S', τ') .*

- If $\xi' = \alpha \xi$, where α is a non vanishing function on M , then $\tau'(X) = \tau(X) + (X \cdot \alpha)/\alpha$ and $h' = h/\alpha$.
- If $\xi' = \xi + \eta$ with η a tangential vector field on M , then $\tau'(X) = \tau(X) + h(X, \eta)$ and $h' = h$.

Proof Straightforward. \square

Proof of Proposition 2.1. Since M is assumed to be strictly locally convex, h is either positive or negative definite, and Lemma 2.3 ensures that the conformal class of h does not depend on the choice of ξ . Now, starting with any transverse vector field ξ , the affine normal will be $\alpha \xi + \eta$ where α is the unique non vanishing function on M such that $\alpha \xi$ satisfies the positivity and normalization conditions 2 and 3, and η is the unique tangent vector field for which the 1-form $\tau_{\alpha \xi}$ corresponding to $\alpha \xi$ reads as $\tau_{\alpha \xi} = -h(\cdot, \eta)$. \square

Definition 2.4 A locally strictly convex hypersurface $M \subset \mathbb{R}^{n+1}$ with affine normal ξ is an affine sphere with affine curvature -1 when the affine shape operator S satisfies $S = -\text{Id}$. In this case, the point $O = m - \xi(m)$ does not depend on $m \in M$: it is called the center of the affine sphere.

The following existence and uniqueness theorem was conjectured by Calabi in [3], and proved by Cheng and Yau in [4]. See also Gigena [6].

Theorem 2.5 (Cheng–Yau) *Let $\mathcal{C} \subset \mathbb{R}^{n+1}$ be an open convex cone that contains no line. There exists a unique embedded affine sphere $\mathcal{H} \subset \mathbb{R}^{n+1}$ with center the origin and affine curvature -1 which is asymptotic to the boundary of \mathcal{C} .*

In any chart where \mathcal{C} reads (with respect to an adapted unimodular frame) as

$$\mathcal{C} = \{t(1, x) \mid x \in \Omega, t > 0\}$$

where $\Omega \subset \mathbb{R}^n$ is a bounded open convex set, \mathcal{H} is the radial graph

$$\mathcal{H} = \left\{ \frac{-1}{u(x)}(1, x) \mid x \in \Omega \right\}$$

where $u := u_\Omega \in C^0(\bar{\Omega}) \cap C^\infty(\Omega)$ is the unique convex solution of the real Monge–Ampère equation

$$\begin{cases} \det D^2 u = (-1/u)^{n+2} & \text{in } \Omega \\ u|_{\partial\Omega} = 0. \end{cases} \quad (2.2)$$

Theorem 2.5 will allow us to systematically identify any properly convex open subset $\Omega \subset \mathbb{RP}^n$ with the corresponding affine sphere \mathcal{H} with affine curvature -1 . The affine metric on \mathcal{H} induces a Riemannian metric on Ω : we still denote it by h , and call it the affine metric on Ω . The corresponding measure μ_h is called the affine measure on Ω .

2.2 From properly convex projective manifolds to affine spheres

Assume now that M is an n -dimensional manifold equipped with a properly convex projective structure. Any of the developing maps identifies its universal cover \widetilde{M} with a properly convex open set $\Omega \subset \mathbb{RP}^n$. Since Ω is invariant under the action of the group $\text{hol}(\pi_1(M)) \subset \text{PGL}_{n+1}\mathbb{R}$, the uniqueness part in Theorem 2.5 ensures that the affine metric on Ω goes to the quotient to give a metric h on M , that we will also call the “affine metric” on M .

On the other hand, the Hilbert distance on Ω derives from the Finsler metric defined for any point $x \in \Omega$ and any vector $X \in T_x\Omega$ as

$$\|X\|_{F,\Omega} = \left(\frac{1}{\|x-a\|} + \frac{1}{\|x-b\|} \right) \|X\|,$$

where a and b are the points of intersection of $\partial\Omega$ with the line defined by $(x; X)$, and $\|\cdot\|$ is any Euclidean norm on an affine chart with $\bar{\Omega} \subset \mathbb{R}^n$. To this Finsler metric, we associate the Borel measure μ_F on $\Omega \subset \mathbb{R}^n$ with density $(1/m(B_{F,\Omega}(x,1)))$ with respect to the Lebesgue measure m on \mathbb{R}^n , where $B_{F,\Omega}(x,1) = \{X \in T_x\Omega, \|X\|_{F,\Omega} < 1\}$ is the unit ball at the point x for the Finsler metric (see eg Marquis [18]). The Finsler metric and the corresponding measure μ_F also go to the quotient to M .

Observe that both the affine and Finsler metrics on M do not depend on the choice of the developing map.

Proposition 2.6 *Let M be a properly convex projective manifold. Then M has finite Finsler volume if and only if it has finite affine volume.*

The rest of the section is devoted to the proof of this proposition. The proof will rely on the continuity of the map $(x, \Omega) \rightarrow u_\Omega(x)$ and on the following cocompactness result due to Benzécri in [2], which will be used again later in this paper to provide estimates for the Pick measure (see Lemma 5.7).

Theorem 2.7 (Benzécri) *Let \mathcal{E} be the set of pairs (x, Ω) , where $\Omega \subset \mathbb{RP}^n$ is a properly convex open subset of \mathbb{RP}^n and x is a point in Ω , equipped with the Hausdorff topology. The natural action of $\mathrm{PGL}_{n+1}\mathbb{R}$ on \mathcal{E} is cocompact.*

As a consequence, the ratio of any two continuous $\mathrm{PGL}_{n+1}\mathbb{R}$ invariant positive functions on \mathcal{E} will be bounded.

Let \mathcal{F} denote the set of pairs (x, Ω) , where $\Omega \subset \mathbb{R}^n$ is a bounded convex domain and x is a point in Ω .

Proposition 2.8 *The value $u_\Omega(x)$ of the solution of the Monge–Ampère equation (2.2) at point x depends continuously on $(x, \Omega) \in \mathcal{F}$.*

Proof Let $(x_0, \Omega_0) \in \mathcal{F}$. When the convex set Ω is close to Ω_0 in the Hausdorff topology,

$$(1 - \varepsilon)\Omega_0 \subset \Omega \subset (1 + \varepsilon)\Omega_0$$

holds for dilations with center x_0 and $\varepsilon > 0$ small. The proposition follows readily from the continuity of $u_{\Omega_0} : \Omega_0 \rightarrow \mathbb{R}$, the easy fact that

$$u_{t\Omega_0}(tx) = t^{n/(n+1)} u_{\Omega_0}(x)$$

holds for all $x \in \Omega_0$ and $t > 0$, and the following lemma. □

Lemma 2.9 *Let $\Omega_1 \subset \Omega_2$ be two proper convex open subsets of \mathbb{R}^n . Then $u_{\Omega_2} \leq u_{\Omega_1}$ on Ω_1 .*

Proof of Lemma 2.9 This assertion is a consequence of the maximum principle. Observe first that, since the function u_{Ω_2} is convex, $u_{\Omega_2} \leq u_{\Omega_1}$ holds on $\partial\Omega_1$. Assuming that the function $u_{\Omega_1} - u_{\Omega_2}$ reaches a negative minimum at an interior point $y \in \Omega_1$, the hessian $D_y^2(u_{\Omega_1} - u_{\Omega_2})$ is positive semi-definite, hence

$$(-1/u_{\Omega_2})^{n+2} = \det D_y^2(u_{\Omega_2}) \leq \det D_y^2(u_{\Omega_1}) = (-1/u_{\Omega_1})^{n+2},$$

thus $|u_{\Omega_1}(y)|^{n+2} \leq |u_{\Omega_2}(y)|^{n+2}$: a contradiction. \square

Proof of Proposition 2.6 Let μ_F and μ_h denote respectively the Finsler and affine measures on a proper convex open subset of $\Omega \subset \mathbb{R}^n$, and let λ be their ratio. It suffices to prove that the function $\lambda : \mathcal{E} \rightarrow]0, \infty[$ is bounded, as well as bounded away from zero. This follows immediately from Benzécri's Theorem 2.7 since :

– λ is invariant under $\text{PGL}_{n+1}\mathbb{R}$, as both volume elements are ;

– λ is continuous. Indeed, in a neighborhood of (x_0, Ω_0) in \mathcal{E} , we may work in an affine chart where Ω_0 reads as a bounded set $\Omega_0 \subset \mathbb{R}^n$. It follows immediately from the definition that the density of the Finsler measure with respect to the Lebesgue measure dx on \mathbb{R}^n depends continuously on (x, Ω) . It remains to check that the density of the affine measure also depends continuously on (x, Ω) : this follows from Proposition 2.8 and the following formula

$$\mu_h = (-u_\Omega)^{-n-1} dx \tag{2.3}$$

that we now prove. Let \mathcal{H} be the affine sphere with affine curvature -1 and center the origin which is asymptotic to the cone \mathcal{C} generated by Ω (Theorem 2.5). Since the affine normal at a point $y \in \mathcal{H}$ is $\xi(y) = y$, the Lebesgue measure dv of \mathbb{R}^{n+1} reads on the cone, in coordinates $v = sy$ with $(s, y) \in]0, \infty[\times \mathcal{H}$, as $dv = s^n ds \mu_h$. On the other hand, in coordinates $v = t(1, x)$ with $(t, x) \in]0, \infty[\times \Omega$, one has $dv = t^n dt dx$. The fact that $s = -u_\Omega(x)t$ yields (2.3). \square

3 Estimates for the affine metric

The aim of this section is to prove the following.

Proposition 3.1 *Let S be a non compact properly convex projective surface with finite volume. Then, the curvature of the affine metric on S tends to a negative constant at infinity.*

Proposition 3.1 will rely on a priori interior estimates which were used by Cheng and Yau to solve the Monge–Ampère equation (see [4]). We recall the by now classical estimates for the Hölder norms of the solutions. For $x \in \mathbb{R}^n$ and $r > 0$, let $B(x, r)$ denote the Euclidean open ball with center x and radius r .

Proposition 3.2 *Let $r > 1$, $\delta > 0$ and $0 < c_0 \leq c_1$.*

Assume $\Omega \subset \mathbb{R}^n$ is a convex open subset such that $B(0,1) \subset \Omega \subset B(0,r)$, and let $v \in C^0(\overline{\Omega}) \cap C^\infty(\Omega)$ be a solution of the Monge–Ampère equation

$$\begin{cases} \det D^2 v = f & \text{in } \Omega \\ v|_{\partial\Omega} = 0, \end{cases} \quad (3.1)$$

where $f \in C^\infty(\overline{\Omega})$ satisfies $c_0 \leq f \leq c_1$. Then, for any $k \in \mathbb{N}$, $k \neq 1$, and $\alpha \in]0, 1[$, one has

$$\|v\|_{C^{k+2,\alpha}(\Omega_\delta)} \leq C,$$

where $\Omega_\delta = \{x \in \Omega / d(x, \partial\Omega) \geq \delta\}$, and C is a constant that depends only on $r, \delta, c_0, c_1, k, \alpha$, and $\|f\|_{C^{k,\alpha}(\Omega)}$.

For $k = 0$ this bound is Caffarelli’s estimate (see Trudinger–Wang [21, Theorem 3.2 ii]). For $k \geq 2$, these bounds result from the uniform ellipticity of the Monge–Ampère operator due to Pogorelov, and from the Schauder’s estimates (see [21, Theorem 3.1]).

Corollary 3.3 *For any $k \in \mathbb{N}$, the k -jet $J_k(u_\Omega)(x)$ of the solution of the Monge–Ampère equation (2.2) on Ω at point x depends continuously on $(x, \Omega) \in \mathcal{F}$.*

Proof of Corollary 3.3. We may assume $k \geq 1$. Let (x_j, Ω_j) converge to (x, Ω) in \mathcal{F} , and $K \subset \Omega$ be a compact neighborhood of x . We may assume that $K \subset \Omega_j$ for every j . Proposition 2.8 ensures that u_{Ω_j} converges uniformly to u_Ω on K and, by convexity of the u_{Ω_j} ’s, one has a uniform bound for the first derivative of every u_{Ω_j} on K . When applied to the function $v := u_{\Omega_j} + \varepsilon$ on the domain $\Omega_{j,\varepsilon} := \{u_{\Omega_j} < -\varepsilon\}$, where $\varepsilon > 0$ is small enough, Proposition 3.2 gives a uniform bound for the jet of order $k + 1$ of every u_{Ω_j} on K . After extracting a subsequence, the Ascoli–Arzela theorem ensures that u_{Ω_j} , as well as its derivatives of order at most k , converges uniformly on K . \square

As a first consequence, we obtain a uniform control of the affine metric in terms of the Finsler metric.

Proposition 3.4 *There exists a constant $c > 0$ such that, for any $(x, \Omega) \in \mathcal{E}$ and $X \in T_x\Omega$:*

$$(1/c) \|X\|_{F,\Omega} \leq \|X\|_{h,\Omega} \leq c \|X\|_{F,\Omega}.$$

Proof One easily verifies that the affine metric on $T_x\Omega$ can be expressed in terms of the 2-jet of u_Ω at point x . The proposition follows again from Benzécri’s Theorem 2.7 since both metrics are $\text{PGL}_{n+1}\mathbb{R}$ invariant and depend continuously on (x, Ω) : this is obvious for the Finsler metric from its definition, and is a consequence of Corollary 3.3 for the affine metric. \square

This observation allows us to recover the following, due to Calabi–Nirenberg [5, Corollary 2].

Corollary 3.5 *The affine metric on any properly convex domain $\Omega \subset \mathbb{RP}^n$ is complete.*

Proof Consequence of Proposition 3.4, and of the completeness of the Finsler metric on Ω . \square

From now on, we specialize to surfaces. Before going into the proof of Proposition 3.1, we state a preliminary result. A projective disk is defined as the orientable component of the complement of a proper conic in \mathbb{RP}^2 . In a suitable affine chart, it reads as the interior of an ellipse.

Proposition 3.6 *Let $\Gamma \subset \text{PGl}_3\mathbb{R}$ be a discrete subgroup preserving a properly convex domain Ω of \mathbb{RP}^2 .*

1. *There exists a convex fundamental domain $K \subset \Omega$ for the action of Γ on Ω , such that the map $K \rightarrow \Gamma \backslash \Omega$ is proper.*

Assume moreover that the quotient $S = \Gamma \backslash \Omega$ has finite volume. Then :

2. *The closure \overline{K} of K in \mathbb{RP}^2 is a finite-sided polyhedron. Let p be a point in $\overline{K} \cap \partial\Omega$. It corresponds to a cusp of S . The holonomy γ of the cusp is regular unipotent, namely is generated by a matrix in the conjugacy class of $\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$.*

3. *There exist two projective disks D_1 and D_2 with*

$$D_1 \subset \Omega \subset D_2,$$

and such that ∂D_1 and ∂D_2 osculate each other at the point $p \in \overline{K} \cap \partial\Omega$, namely the disks have a contact of order at least 2 at this point.

Remark 3.7 *As a consequence, the curve $\partial\Omega$ not only admits a tangent at point p , but also osculating conics.*

Assertion 1 is due to Lee [11], and assertion 2 to Marquis [18].




Figure 1: a. The pencil of conics – b. The disks D_1 and D_2

Proof We only have to prove assertion 3. We develop a strategy initiated by Benzécri in [2] (see Goldman [8, Proposition 6.14] and Marquis [18, Proposition 5.21]). We choose homogeneous

coordinates $[x, y, z]$ with $p = [1, 0, 0]$, and where the holonomy of the cusp is generated by $\gamma = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$. The holonomy preserves each quadratic form in the pencil generated by z^2 and $y^2 - (y + 2x)z$. The corresponding pencil of conics has base point p and reads in the chart $x = 1$ (see Figure 1a) as

$$\begin{cases} z^2 = 0 \\ (y - z/2)^2 + (\mu - 1/4)z^2 - 2z = 0 \quad (\mu \in \mathbb{R}). \end{cases}$$

All these conics admit the line $z = 0$ as a tangent at the point p and, except for the singular one, they do osculate each other at this point.

Now pick an affine chart containing Ω as a bounded subset and new affine coordinates (y, z) on this chart for which $p = (0, 0)$ and the degenerate conic of the pencil still reads as $z^2 = 0$. In this chart the bounded domain Ω , being invariant under γ , lies in the half-plane (say $z > 0$) containing the ellipses in the pencil.

- We first build the exterior disk D_2 . Choose $m \in \Omega$ and let Δ be the sector in the half-plane $\{z > 0\}$ delimited by the line (pm) and its image $\gamma(pm) = p\gamma(m)$. One of the conics in the pencil (actually, one of the hyperbolas) contains Δ in its interior D_2 . This obviously gives $\Omega \cap \Delta \subset D_2$ hence, by invariance under γ ,

$$\gamma^k(\Omega \cap \Delta) = \Omega \cap \gamma^k(\Delta) \subset D_2 \quad \forall k \in \mathbb{Z},$$

and finally $\Omega \subset D_2$.

- We now construct the interior disk. The domain Ω , being convex, contains the triangle with vertices $p, m, \gamma(m)$. Since the ellipses in the pencil become tiny when the parameter μ goes to infinity, there exists one of them, say D_1 , for which $D_1 \cap \Delta$ is included in this triangle. We conclude as above that $D_1 \subset \Omega$. \square

Observe that the affine metric on a projective disk has constant negative curvature : this follows from the fact that $\text{SO}(2, 1) \subset \text{Sl}_3\mathbb{R}$ acts transitively on each sheet of the hyperboloid. This observation will be crucial in the following.

Proof of Proposition 3.1 We use the notations and conclusion of the previous proposition. Let us focus on the cusp $p \in \overline{K} \cap \Omega$. Choose a second point $q \neq p$ on the boundary of the exterior disk D_2 . Let r be the point of intersection of the tangent lines to D_2 at points p and q , and $(\psi_t)_{t \in \mathbb{R}}$ be the 1-parameter group of projective transformations that read as $\psi_t = \begin{pmatrix} e^{-t} & 0 & 0 \\ 0 & e^t & 0 \\ 0 & 0 & 1 \end{pmatrix}$

with respect to the frame (p, q, r) .

In the corresponding homogeneous coordinates $[x, y, z]$, this 1-parameter group preserves both quadratic forms z^2 and xy . Hence, each orbit of (ψ_t) is an arc delimited by p and q either on a conic through p and q and with tangent (pr) at p and (qr) at q , or on the line (pq) . Observe that, except for ∂D_2 itself, none of these conics osculate ∂D_2 at the point p (Figure 2a).




Figure 2: The orbits of (ψ_t)

Since ∂D_1 does osculate ∂D_2 at p , this implies that the orbit of any point m in D_2 meets D_1 or, more specifically, that

$$\forall m \in D_2, \exists T \in \mathbb{R} / \forall t \leq T, \psi_t(m) \in D_1.$$

We infer that $\psi_t(D_1)$ converges to D_2 in the Hausdorff topology when t goes to $+\infty$. Since $D_1 \subset \Omega \subset D_2$, $\psi_t(\Omega)$ also converges to D_2 . Thus, by Corollary 3.3, the curvature of the affine metric $h_{\psi_t(\Omega)}$ converges uniformly to the curvature of h_{D_2} , that is to a negative constant, on compact subsets of D_2 when $t \rightarrow +\infty$.

Let \mathcal{T} be a triangular neighborhood of p in \overline{K} with interior $\overset{\circ}{\mathcal{T}} \subset D_1$. There exists a compact subset $C \subset D_2$ such that each orbit $(\psi_t(m))_{t \in \mathbb{R}}$ of a point $m \in \mathcal{T} \cap \Omega$ intersects C (Figure 2b). The farther $m \in \mathcal{T}$ is in the cusp (that is, the closer to p), the greater the values of t for which $\psi_t(m) \in C$. Since the curvature of h_Ω at the point m equals the curvature of $h_{\psi_t(\Omega)}$ at the point $\psi_t(m)$, this concludes the proof. \square

4 The Pick tensor

The aim of this section, which basically follows the presentation by Labourie in [10], is to define the Pick form U of an affine 2-dimensional sphere (Definition 4.9) – which is the second component of the map $\mathcal{A} \rightarrow (J, U)$ in Theorem 1.1.

4.1 From affine spheres to properly convex projective manifolds

To an n -dimensional properly convex projective manifold M with developing map $\text{dev} : \widetilde{M} \rightarrow \Omega \subset \mathbb{RP}^n$, Theorem 2.5 associates an embedding of the universal cover $\widetilde{M} \hookrightarrow \mathbb{R}^{n+1}$ as the affine sphere with affine curvature -1 asymptotic to the boundary of the cone over Ω . Moreover, the affine metric on \widetilde{M} is complete (Corollary 3.5).

On the other hand, all complete affine spheres of \mathbb{R}^{n+1} have been described by Cheng–Yau and An-Min Li in [5], [12] as follows, thus complementing Theorem 2.5 and proving the whole Calabi’s conjecture stated in [3].

Theorem 4.1 (Cheng and Yau, An-Min Li) *Let $\mathcal{H} \subset \mathbb{R}^{n+1}$ be an immersed hypersurface, which is an affine sphere with constant curvature -1 and center the origin. Assume that the corresponding affine metric on \mathcal{H} is complete. Then \mathcal{H} is embedded, and is asymptotic to the boundary of a convex cone $\mathcal{C} \subset \mathbb{R}^{n+1}$ which contains no line.*

Thus, the data of a properly convex projective structure on the manifold M is equivalent to the data of an immersion $\widetilde{M} \hookrightarrow \mathbb{R}^{n+1}$ of its universal cover as a complete affine sphere with affine curvature -1 and center the origin, which is equivariant under a representation $\pi_1(M) \rightarrow \mathrm{SI}_{n+1}^{\pm} \mathbb{R}$.

When $M \subset \mathbb{R}^{n+1}$ is an affine sphere with affine curvature -1 and ξ is the affine normal, recall that the affine flat connection ∇ on \mathbb{R}^{n+1} induces a torsion-free connection D on TM as well as a positive definite 2-form h on TM which satisfy Equation (4.1) below. Indeed, in this case, Equation (2.1) holds with $S = -\mathrm{Id}$ and $\tau = 0$ (see Proposition 2.1(1) and Definition 2.4). In the following subsection, we will study in an intrinsic way such pairs (D, h) on M , where D is a torsion-free connection and h is a Riemannian metric.

4.2 The intrinsic geometry of an affine sphere

Let M be an n -dimensional manifold. The letters X, Y, Z, W will denote tangent vector fields on M .

Let D be a torsion-free connection on TM , h be a positive definite symmetric 2-form on TM and ξ be a non vanishing section of the trivial line bundle L over M . To these data, we associate a torsion-free connection ∇ on $TM \oplus L$ by letting

$$\begin{cases} \nabla_X Y = D_X Y + h(X, Y) \xi \\ \nabla_X \xi = X. \end{cases} \quad (4.1)$$

In the next three lemmas, we will explore the conditions under which the connection ∇ will be flat and volume preserving.

Lemma 4.2 *The curvature R^∇ of the connection ∇ is given by*

$$\begin{cases} R^\nabla(X, Y)Z = R(X, Y)Z - h(Y, Z)X + h(X, Z)Y + d^D h(X, Y, Z) \xi \\ R^\nabla(X, Y) \xi = 0, \end{cases} \quad (4.2)$$

where R is the curvature of D and $d^D h(X, Y, Z) := (D_X h)(Y, Z) - (D_Y h)(X, Z)$.

Proof Use the definition $R^\nabla(X, Y)Z := \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z$. \square

Lemma 4.3 *Write $D = D^h + A$, where D^h is the Levi-Civita connection of the Riemannian metric h and A is a section of the bundle $T^*M \otimes \mathrm{End} TM$. Then :*

1. $A(X)Y = A(Y)X$

2. The connection ∇ defined by Equation (4.1) is flat if and only if :
- each endomorphism $A(X)$ is symmetric with respect to h
 - $d^{D^h} A = 0$, where $d^{D^h} A(X, Y) := (D_X^h A)(Y) - (D_Y^h A)(X)$
 - $R^h(X, Y)Z + [A(X), A(Y)]Z = h(X, Z)Y - h(Y, Z)X$, where R^h is the curvature of D^h

Proof 1. Both connections D and D^h are torsion-free.

2. The normal part of R^∇ vanishes if and only if $d^D h = 0$ (Lemma 4.2). Since $D^h h = 0$,

$$(D_X h)(Y, Z) = -h(A(X)Y, Z) - h(Y, A(X)Z)$$

holds thus (1.) yields

$$d^D h(X, Y, Z) = h(X, A(Z)Y) - h(Y, A(Z)X),$$

which proves (a).

The tangential part of R^∇ vanishes if and only if $R(X, Y)Z = h(Y, Z)X - h(X, Z)Y$. A simple computation gives

$$R(X, Y)Z = R^h(X, Y)Z + (d^{D^h} A(X, Y))Z + [A(X), A(Y)]Z \quad (4.3)$$

hence the result since, for fixed X and Y , the endomorphism $Z \rightarrow h(X, Z)Y - h(Y, Z)X$ as well as those defined by each term in the right-hand side of Equation (4.3) are skew symmetric with respect to h , except for $Z \rightarrow (d^{D^h} A(X, Y))Z$ which is symmetric. \square

Lemma 4.4 Assume that M is oriented. Define a volume form ω on $TM \oplus L$ by imposing that $\omega(X_1, \dots, X_n, \xi) = 1$, whenever (X_1, \dots, X_n) is a positive h -orthonormal frame of TM . Then

$$\nabla \omega = 0 \quad \text{iff} \quad D\omega_h = 0 \quad \text{iff} \quad \forall X, \text{Tr} A(X) = 0,$$

where ω_h denotes the volume form of h and Tr is the trace operator.

Remark 4.5 This lemma is motivated by condition (3) in Proposition 2.1, where a constant volume form on \mathbb{R}^{n+1} is used to normalize the affine normal vector.

Proof Observe that, since $\nabla_X Y - D_X Y$ is colinear to ξ and $\nabla_X \xi$ is a tangent vector field, one has

$$\begin{aligned} D_X \omega_h(X_1, \dots, X_n) &= X \cdot \omega(X_1, \dots, X_n, \xi) - \sum_i^n \omega(X_1, \dots, \nabla_X X_i, \dots, X_n, \xi) \\ &= \nabla_X \omega(X_1, \dots, X_n, \xi). \end{aligned}$$

This proves the first equivalence. On the other hand, since $D = D^h + A$ and $D^h \omega_h = 0$, one has

$$D_X \omega_h(X_1, \dots, X_n) = - \sum_{i=1}^n \omega_h(X_1, \dots, A(X)X_i, \dots, X_n) = -\text{Tr} A(X) \omega_h(X_1, \dots, X_n),$$

which concludes the proof. \square

Corollary 4.6 Define a tensor C on TM by letting

$$C(X, Y, Z) := h(A(X)Y, Z).$$

The 3-tensor C is symmetric if and only if each endomorphism $A(X)$ is symmetric with respect to h . In that case, one also has $(D_X h)(Y, Z) = -2C(X, Y, Z)$.

Proof The first assertion immediately follows from Lemma 4.3 (1), and the second from the fact that $D^h h = 0$. \square

Remark-Definition 4.7 Assume that $M \subset \mathbb{R}^{n+1}$ is an affine sphere, and that D and h are respectively the Blaschke connection and the affine metric on M . Then the corresponding 3-tensor C is symmetric. It is called the “Pick tensor” of M . The Pick tensor C vanishes if and only if the affine sphere M is part of an hyperquadric (see Cheng–Yau [5, Theorem 1]).

4.3 The Pick form

When specializing the previous section to surfaces, we have the following.

Lemma 4.8 Assume that the manifold M is 2-dimensional and oriented. Let J denote the underlying complex structure to h and suppose that the 3-tensor C is symmetric. Then, one has the equivalence : each endomorphism $A(X)$ is trace-free and $d^{D^h} A = 0$ if and only if C is the real part of a holomorphic cubic differential U on (M, J) .

Proof Pick local isothermal coordinates (x, y) where the affine metric reads as $h = e^{2w}(dx^2 + dy^2)$, and let $z = x + iy$. Observe that both endomorphisms $A(\partial_x)$ and $A(\partial_y)$ are trace-free if and only if the symmetric tensor C reads in this chart as

$$C = P dx^3 - 3P dx dy^2 - 3Q dx^2 dy + Q dy^3$$

where P and Q are real-valued functions or, in other words, if

$$C = \operatorname{Re}((P + iQ)(dx + idy)^3) = \operatorname{Re}(f(z) dz^3)$$

where $f = P + iQ$ is complex-valued. Assume that these conditions are satisfied. It remains to check under which condition $d^{D^h} A = 0$ holds. Observe first that this is equivalent to requiring that $d^{D^h} C = 0$, where $d^{D^h} C(X, Y, Z, W) := (D_X^h C)(Y, Z, W) - (D_Y^h C)(X, Z, W)$.

Let $\partial_z = (1/2)(\partial_x - i\partial_y)$ and $\partial_{\bar{z}} = (1/2)(\partial_x + i\partial_y)$. Since $h(\partial_z, \partial_z) = h(\partial_{\bar{z}}, \partial_{\bar{z}}) = 0$ while $h(\partial_z, \partial_{\bar{z}}) = (1/2)e^{2w}$, and since the connection D^h is torsion-free, we infer that $D_{\partial_z}^h \partial_z$ and $D_{\partial_{\bar{z}}}^h \partial_{\bar{z}}$ are respectively colinear to ∂_z and $\partial_{\bar{z}}$, and that $D_{\partial_{\bar{z}}}^h \partial_z = D_{\partial_z}^h \partial_{\bar{z}} = 0$.

The definition of C ensures that, for any vector field Y , one has $C(Y, \partial_z, \partial_{\bar{z}}) = 0$. It then follows from the expression of the Levi–Civita connection D^h with respect to the frame $(\partial_{\bar{z}}, \partial_{\bar{z}})$ that $(D_X^h C)(Y, \partial_z, \partial_{\bar{z}}) = 0$ holds for any vector fields X and Y . Since the tensor $d^{D^h} C$ is skew-symmetric with respect to the first couple of variables, and symmetric with respect to the second, it follows that $d^{D^h} C = 0$ if and only if

$$d^{D^h} C(\partial_{\bar{z}}, \partial_z, \partial_z, \partial_z) = 0.$$

The easy fact that $d^{D^h} C(\partial_{\bar{z}}, \partial_z, \partial_z, \partial_z) = \partial f / \partial \bar{z}$ now ends the proof. \square

Remark-Definition 4.9 *It follows from Lemma 4.8 that, when $\mathcal{H} \subset \mathbb{R}^3$ is a 2-dimensional affine sphere, the Pick tensor C reads as $C = \operatorname{Re}U$ where the “Pick form” U is a holomorphic cubic differential on (\mathcal{H}, J) . We will also denote by U the Pick form on any properly convex domain of \mathbb{RP}^2 , and on any properly convex projective surface.*

5 The conformal structure of (S, h) and the Pick form

Let S be an oriented surface with non-abelian fundamental group, equipped with a finite volume properly convex projective structure. We first explore the conformal structure of the affine metric on S , and show that it has only parabolic ends (Corollary 5.4). Then, we prove the estimates for the Pick form at the cusps (Corollary 5.8) that will ensure that the map $\mathcal{A} \rightarrow (J, U)$ of our main theorem 1.1 is well-defined.

Proposition 5.1 *Let S be an oriented finite volume properly convex projective surface with non-abelian fundamental group, h be the affine metric on S and write $h = e^{2v}h_0$ (with $v \in C^\infty(S)$) where h_0 is the hyperbolic metric in the conformal class of h . Then h and h_0 are “conformally quasi-isometric”, namely the conformal factor v is bounded.*

Proof Since S has non-abelian fundamental group, the uniformization theorem ensures that there exists an hyperbolic metric h_0 in the conformal class of h . The affine metric is complete (Corollary 3.5) and its curvature is negatively pinched at infinity (Proposition 3.1). This makes Proposition 5.1 a special case of the following version of the Ahlfors–Schwarz lemma [1]. \square

Lemma 5.2 *Let $h = e^{2v}h_0$ be two conformal metrics on a non compact surface S , and with respective curvatures K and K_0 . Assume that both metrics are complete and that*

$$-a \leq K, K_0 \leq -b < 0$$

holds at infinity for some positive constants a and b . Then h and h_0 are conformally quasi-isometric.

This lemma will be, as in Troyanov [20], a consequence of the generalized maximum principle of Yau [23].

Theorem 5.3 (Yau) *Let w be a smooth function on a complete Riemannian manifold (M, h_0) whose Ricci curvature is bounded below. If w is bounded above and does not reach its maximum, there exists a sequence of points $p_n \in M$ going to infinity, and such that $w(p_n) \rightarrow \sup_M w$, $|\nabla_0 w(p_n)| \rightarrow 0$, and with $\lim \Delta_0 w(p_n) \in [0, \infty]$.*

Here ∇_0 and Δ_0 respectively denote the gradient and the Laplace operator with respect to the metric h_0 , with the sign convention $\Delta_0 := -\operatorname{Tr}D^{h_0}\nabla_0$.

Proof of Lemma 5.2 By symmetry, it is sufficient to prove that v is bounded above. If this is not the case, the function $w := 1/(1 + e^{-v})$ does not reach its maximum, and there exists a sequence $p_n \in S$ as in Theorem 5.3, with $w(p_n) \rightarrow 1$. A simple computation yields

$$\Delta_0 w = \Delta_0 v \frac{e^{-v}}{(1 + e^{-v})^2} - |\nabla_0 w|^2 (e^v - e^{-v}).$$

Plugging the equation for curvatures

$$\Delta_0 v = K e^{2v} - K_0 \tag{5.1}$$

into this expression gives

$$\Delta_0 w = K \frac{e^v}{(1 + e^{-v})^2} - K_0 \frac{e^{-v}}{(1 + e^{-v})^2} - |\nabla_0 w|^2 (e^v - e^{-v}),$$

hence a contradiction when evaluated at the points p_n , since the first term goes to $-\infty$, the second one goes to zero and the third one is non positive when $n \rightarrow \infty$. \square

Corollary 5.4 *Let S be an oriented finite volume properly convex projective surface with non-abelian fundamental group. Let J be the underlying complex structure to the affine metric h on S . Then S has finite topological type and the Riemann surface (S, J) has only parabolic ends. In other words, there exist a compact Riemann surface \bar{S} and a finite set $\{p_1, \dots, p_k\} \subset \bar{S}$ such that (S, J) identifies with $\bar{S} \setminus \{p_1, \dots, p_k\}$.*

The fact that S has finite topological type is due to Marquis, and was used in the proof of Proposition 3.1.

Proof Since h has finite volume, Proposition 5.1 ensures that the hyperbolic metric h_0 on (S, J) also has finite volume, hence all its ends are parabolic. \square

Remark 5.5 *In a complex chart $\{z \in \mathbb{C}, 0 < |z| \leq 1/2\}$ around a puncture, both metrics h and h_0 will be conformally quasi-isometric to the Beltrami metric $g = \frac{|dz|^2}{|z|^2 |\log|z||^2}$. This is again a consequence of Lemma 5.2.*

To conclude this section, it only remains to prove that the Pick form U on S (Definition 4.9) is a meromorphic form on the compactification \bar{S} of S , with a pole of order at most 2 at each puncture p .

Definition 5.6 *Let Ω be a properly convex open subset of \mathbb{RP}^2 and U be the corresponding Pick form on Ω . We define the Pick measure $\mu_P := |U|^{2/3}$ on Ω by*

$$\mu_P = |f(z)|^{2/3} |dz|^2$$

in any complex chart where U reads as $U(z) = f(z) dz^3$.

Lemma 5.7 *There exists a constant $C > 0$ such that, for any properly convex open subset Ω of \mathbb{RP}^2 , the ratio $\Lambda \in [0, \infty[$ of the Pick measure μ_P by the affine measure μ_h is uniformly bounded above by C .*

Proof Since the Pick measure can be expressed in terms of the 3-jet of u_Ω , Corollary 3.3 and Equation (2.3) ensure that the ratio Λ depends continuously on $(x, \Omega) \in \mathcal{E}$. This ratio being invariant under the action of $\mathrm{PGL}_3\mathbb{R}$, we conclude by Benzécri's cocompactness theorem 2.7 that Λ is bounded. \square

Corollary 5.8 *Let S be an oriented finite volume properly convex projective surface with non-abelian fundamental group. Then the Pick form U is a meromorphic cubic differential on the compactification \bar{S} of (S, J) , and with poles of order at most 2 at each puncture p_i ($1 \leq i \leq k$).*

Proof Work in a complex chart $D^* = \{z \in \mathbb{C}, 0 < |z| \leq 1/2\}$ around a cusp of (S, J) and let

$$U(z) = f(z) dz^3,$$

where f is a holomorphic function on the punctured disk D^* . Since the affine metric h is conformally quasi-isometric to the Beltrami metric in the cusp (Remark 5.5), the boundedness of Λ (Lemma 5.7) implies that there exists a constant $c > 0$ such that

$$|f(z)| \leq \frac{c}{|z|^3 |\log |z||^3} \quad \text{holds for } 0 < |z| \leq 1/2.$$

Hence f is meromorphic on the disk $D = \{|z| \leq 1/2\}$, with a pole of order at most 2 at the origin. \square

We just proved that the map $\mathcal{A} \rightarrow (J, U)$ is well-defined. This concludes the first part of the proof of Theorem 1.1.

6 Wang's equation

In this final section, we wrap up the proof of our main theorem 1.1 by constructing the inverse map $(J, U) \rightarrow \mathcal{A}$ as in Loftin's thesis [13].

6.1 Retrieving the projective structure

In this subsection, we continue the discussion in section 4, and characterize those pairs (D, h) of a torsion-free connection D and a positive symmetric 2-form h on the tangent bundle of a manifold M which come from an immersion $M \hookrightarrow \mathbb{R}^{n+1}$ as an affine sphere.

Proposition 6.1 *Let M be an n -dimensional manifold endowed with a torsion-free connection D and h be a positive definite symmetric 2-form on M . Write $D = D^h + A$, where D^h denotes the Levi-Civita connection of h and A is a section of $T^*M \otimes \text{End} TM$.*

Assume the manifold M to be simply-connected. Then D is the Blaschke connection and h is the affine metric corresponding to an immersion $M \hookrightarrow \mathbb{R}^{n+1}$ as an affine sphere with constant curvature -1 if and only if the following hold :

- a. *each endomorphism $A(X)$ is trace-free and symmetric with respect to h*
- b. *$d^{D^h} A = 0$*
- c. *$R^h(X, Y)Z + [A(X), A(Y)]Z = h(X, Z)Y - h(Y, Z)X$, where R^h is the curvature of D^h*

Proof Lemmas 4.3 and 4.4 ensure that conditions a,b and c are necessary.

Assume now that these conditions are satisfied and choose a non vanishing section ξ of the trivial line bundle over M . Then, the connection ∇ on $TM \oplus L$ defined by (4.1) is flat, hence it identifies the bundle $TM \oplus L$ with the trivial bundle $\mathbb{R}^{n+1} \times M \rightarrow M$ and the section

$$m \in M \rightarrow \xi(m) \in T_m M \oplus L_m \simeq \mathbb{R}^{n+1}$$

provides an immersion of M in \mathbb{R}^{n+1} such that the connection ∇ derives from the affine flat connection on \mathbb{R}^{n+1} . Moreover since each endomorphism $A(X)$ is supposed to be trace-free, the form ω in Lemma 4.4, being parallel with respect to ∇ , is the restriction to M of a multiple of the constant volume form \det on \mathbb{R}^{n+1} . Hence, after renormalizing the determinant, ξ identifies with the affine normal of this immersion (Proposition 2.1). Equation (4.1) now tells us that the affine shape operator of this immersion is $S = -\text{Id}$, that is, M is immersed in \mathbb{R}^{n+1} as an affine sphere with affine curvature -1 . \square

6.2 Stating Wang's equation

We now specialize to surfaces and introduce Wang's equation, which relates the Pick form U of an affine sphere (Definition 4.9) and the conformal factor between the affine metric h and its uniformization h_0 .

Let S be an oriented surface with non-abelian fundamental group. The construction of the reciprocal map $(J, U) \rightarrow \mathcal{A}$ in theorem 1.1 goes as follows. Given a pair (J, U) , where J is a complex structure on S and U is a holomorphic cubic differential on (S, J) , we want to retrieve the properly convex projective structure \mathcal{A} . As explained in subsection 4.1, this boils down to constructing a suitable embedding $\tilde{S} \subset \mathbb{R}^3$ of the universal cover of S as an affine sphere.

We first define a symmetric 3-tensor on S by letting $C = \text{Re}U$. Let h_0 be the hyperbolic metric on (S, J) . To each conformal metric $h = e^{2v}h_0$ on S we associate the section A of the bundle $T^*S \otimes \text{End}TS$ for which $C(X, Y, Z) := h(A(X)Y, Z)$ (see Corollary 4.6). We then define a torsion-free connection D on TS by $D := D^h + A$, where D^h denotes the Levi-Civita connection of h . These data (D, h) lift to the universal cover \tilde{S} , and Proposition 6.1 gives necessary and sufficient conditions for this pair to correspond to an immersion of \tilde{S} in \mathbb{R}^3 as an affine sphere.

Remark 6.2 *When dealing with surfaces, we have proved in Lemma 4.8 that the above conditions a and b in Proposition 6.1 amount to saying that the tensor $C(X, Y, Z) = h(A(X)Y, Z)$ is the real part of a holomorphic cubic differential U on (S, J) . Thus, it only remains to examine condition c, which we will do in the next corollary.*

Let A_0 be the section of $T^*S \otimes \text{End}TS$ corresponding to the hyperbolic metric h_0 , namely such that $C(X, Y, Z) := h_0(A_0(X)Y, Z)$, and let $\Delta_0 := -\text{Tr}D^{h_0}\nabla_0$ denotes the Laplace operator on (S, h_0) .

Corollary 6.3 *Let the function $k : S \rightarrow \mathbb{R}$ be defined as*

$$k := -h_0([A_0(X_1^0), A_0(X_2^0)]X_2^0, X_1^0), \quad (6.1)$$

where (X_1^0, X_2^0) is any orthonormal frame for the hyperbolic metric h_0 .

The pair (D, h) derives from an immersion of \tilde{S} in \mathbb{R}^3 as an affine sphere with affine curvature -1 if and only if the conformal factor v is solution of Wang's equation

$$\Delta_0 v = -e^{2v} + 1 + k e^{-4v}. \quad (6.2)$$

Proof Condition c is equivalent to

$$h(R^h(X_1, X_2)X_2, X_1) + h([A(X_1), A(X_2)] X_2, X_1) = -1, \quad (6.3)$$

where (X_1, X_2) is any orthonormal basis for the metric h . It is elementary to check that

$$-h([A(X_1), A(X_2)] X_2, X_1) = e^{-6v} k.$$

Taking into account the equation for curvature $\Delta_0 v = K_h e^{2v} + 1$ – where K_h is the curvature of the affine metric h – reduces this condition to Wang’s equation (6.2). \square

6.3 Solving Wang’s equation

In this section, we finally end the proof of Theorem 1.1 by proving that Wang’s equation admits a unique bounded solution.

Let S be an oriented surface equipped with a pair (J, U) , where J is a complex structure and U is a holomorphic cubic differential as in Theorem 1.1. If these data derive (by the construction of sections 2–4) from a properly convex structure on S with finite volume, we have seen that the corresponding affine metric h is complete and must read as $h = e^{2v} h_0$, where h_0 is the hyperbolic metric on (S, J) and the function v is a *bounded solution* of Wang’s equation (Proposition 5.1 and Corollary 6.3).

Lemma 6.4 *Let (S, J) be a hyperbolic Riemann surface with finite volume, and U be a holomorphic cubic differential with poles of order at most 2 at the cusps. Then, the function k defined by Equation (6.1) is non negative and bounded.*

Proof It is elementary to check that k reads as $k = 2e^{-6\varphi} |f|^2$ in isothermal coordinates where $h_0 = e^{2\varphi}(dx^2 + dy^2)$ and $U = f(z) dz^3$. The hypothesis on U and Remark 5.5 ensure that k is bounded on S . \square

Proposition 6.5 (Loftin [13]) *Let (S, h_0) be a Riemannian surface, and $k : S \rightarrow \mathbb{R}^+$ be a smooth non negative bounded function. Then, Wang’s equation (6.2)*

$$\Delta_0 v = -e^{2v} + 1 + k e^{-4v}$$

admits bounded solutions, which are C^∞ . When S is compact, or the metric h_0 is complete and with curvature bounded from below, such a bounded solution is unique.

End of proof of Theorem 1.1 Let v be the unique bounded solution of Wang’s equation (6.2). The conformal metric $h = e^{2v} h_0$ on S is complete. The affine metric h and the Pick form U lift to \tilde{S} , and Corollary 6.3 and Theorem 4.1 ensure that these data correspond to an embedding of $\tilde{S} \hookrightarrow \mathbb{R}^3$ which is asymptotic to a convex cone $\mathcal{C} \subset \mathbb{R}^3$. Such a bounded solution being unique, this embedding is equivariant under a representation $\pi_1(S) \rightarrow \mathrm{Sl}_3^\pm \mathbb{R}$, hence provides the desired properly convex projective structure on S . \square

For the sake of completeness we now give a proof of Proposition 6.5, which relies on the classical so-called method of upper and lower solutions.

Proof of Proposition 6.5 Define $f : S \times \mathbb{R} \rightarrow \mathbb{R}$ by $f(x, t) := -e^{2t} + 1 + k(x)e^{-4t}$. Since k is positive and bounded, there exist constant functions $v_- \leq v_+$ which provide respectively lower and upper solutions for Equation (6.2), namely such that $\Delta_0 v_- \leq f(x, v_-)$ and $\Delta_0 v_+ \geq f(x, v_+)$. Moreover, there exists a constant $c > 0$ such that, for any $v_- \leq s < t \leq v_+$ and $x \in S$,

$$f(x, s) - c(t - s) \leq f(x, t) < f(x, s) \quad (6.4)$$

holds.

• Assume first that the surface S is non compact, and choose an exhaustion $\Sigma_k \subset \mathring{\Sigma}_{k+1}$, $\cup \Sigma_k = S$ of S by compact surfaces with smooth boundaries.

Lemma 6.6 *Let Σ be a compact surface with smooth boundary. Let ψ, v_+ be two functions in $C^\infty(\Sigma)$ and $c > 0$ be a constant. Then, there exists a solution $w \in C^\infty(\Sigma)$ of the linear Dirichlet problem*

$$\begin{cases} \Delta_0 w = -cw + \psi & \text{on } \mathring{\Sigma} \\ w = v_+ & \text{on } \partial\Sigma. \end{cases}$$

Proof Let \mathring{H}_1^2 be the Hilbert completion of the space $C_0^\infty(\mathring{\Sigma})$ of smooth functions with compact support in $\mathring{\Sigma}$ for the inner product $(f, g)_{L_2} + c(\nabla_0 f, \nabla_0 g)_{L_2}$. The Riesz representation theorem in \mathring{H}_1^2 yields a solution $\hat{w} \in \mathring{H}_1^2$ for the equation $\Delta_0 \hat{w} = -c\hat{w} + \hat{\psi}$, where $\hat{\psi} := \psi - cv_+ - \Delta_0 v_+$ lies in $C^\infty(\Sigma)$. Standard elliptic regularity and Sobolev embedding (see eg Gilbarg–Trudinger [7, 8.13 and 7.26]) ensure that $\hat{w} \in C^\infty(\Sigma)$. The function $w := \hat{w} + v_+$ is the desired solution. \square

Corollary 6.7 *Let Σ be a compact surface with smooth boundary, $f \in C^\infty(\mathbb{R} \times \Sigma)$ and $c > 0$ be a constant such that (6.4) holds. Let ψ, v_+ and v_- be functions in $C^\infty(\Sigma)$, and assume that $v_- \leq v_+$, and that $\Delta_0 v_+ \geq f(x, v_+)$ holds while $\Delta_0(x, v_-) \leq f(x, v_-)$. Then, the non-linear Dirichlet problem*

$$\begin{cases} \Delta_0 v = f(x, v) & \text{on } \mathring{\Sigma} \\ v = v_+ & \text{on } \partial\Sigma \end{cases} \quad (6.5)$$

admits a solution $v \in C^\infty(\Sigma)$ with $v_- \leq v \leq v_+$.

Proof We use an iterative scheme. Begin with $w_0 \equiv v_+$, and let $w_j \in C^\infty(\Sigma)$ ($j \geq 1$) be the solution of the linear Dirichlet problem

$$\begin{cases} \Delta_0 w_j = -cw_j + f(x, w_{j-1}) + cw_{j-1} & \text{on } \mathring{\Sigma} \\ w_j = v_+ & \text{on } \partial\Sigma \end{cases}$$

provided by Lemma 6.6. The maximum principle and property (6.4) ensure recursively that (w_j) is a decreasing sequence, that the w_j are upper solutions for Equation (6.5), and that

$v_- \leq w_j \leq v_+$ holds for $j \geq 1$. Global elliptic estimates and Kondrakov embedding (Gilbarg–Trudinger [7, 9.14 and 7.26]) show that the sequence w_j is bounded in $C^{1,\alpha}$ ($0 < \alpha < 1$). Hence a subsequence of (w_j) converges uniformly on Σ to a weak solution v of Equation (6.5). Elliptic regularity ensures that v is smooth. \square

End of proof of Proposition 6.5. For each compact surface $\Sigma_k \subset S$, let $v_k \in C^\infty(\Sigma_k)$ with $v_- \leq v_k \leq v_+$ be the solution of the Dirichlet problem (6.5) on Σ_k given by Corollary 6.7. The maximum principle ensures that the sequence (v_k) is decreasing, namely that $v_{k+1} \leq v_k$ holds on Σ_k . Since the sequence is bounded below, it converges to a weak bounded solution v of Wang’s equation. Elliptic regularity again shows that v is smooth.

Uniqueness of a bounded solution for Equation (6.2) is an immediate consequence of the generalized maximum principle of Yau (Theorem 5.3).

- When the surface S is compact, a bounded solution is provided by the same recursive scheme – without boundary conditions – and uniqueness of a bounded solution follows from the classical maximum principle. \square

Remark 6.8 Denote by \mathcal{G} the set of properly convex projective structures with finite volume on S , quotiented by the natural action of the group $\text{Diff}_0(S)$ of diffeomorphisms isotopic to the identity map (see Goldman [9]). One of the interesting features of Theorem 1.1 is that, by applying Riemann-Roch, it allows to identify \mathcal{G} with $\mathbb{C}^{8(g-1)+3r}$, where g is the genus of the compactification \bar{S} of S , and r is the number of punctures. In particular, it endows \mathcal{G} with a canonical complex structure.

Y. Benoist and D. Hulin
 Université Paris-Sud
 Laboratoire de Mathématiques – UMR 8628
 Orsay 91405, France

yves.benoist@math.u-psud.fr
 dominique.hulin@math.u-psud.fr

References

- [1] Lars Ahlfors. An extension of Schwarz's lemma. *Trans. AMS*, 43:359–364, 1938.
- [2] Jean-Paul Benzécri. Sur les variétés localement affines et localement projectives. *Bull. Soc. Math. France*, 88:229–332, 1960.
- [3] Eugenio Calabi. Complete affine hyperspheres. I. In *Symposia Mathematica, Vol. X*, pages 19–38. 1972.
- [4] Shiu Yuen Cheng and Shing Tung Yau. On the regularity of the Monge–Ampère equation $\det(\partial^2 u / \partial x_i \partial x_j) = F(x, u)$. *Comm. Pure Appl. Math.*, 30:41–68, 1977.
- [5] Shiu Yuen Cheng and Shing Tung Yau. Complete affine hypersurfaces I. The completeness of affine metrics. *Comm. Pure Appl. Math.*, 39:839–866, 1986.
- [6] Salvador Gigena. On a conjecture by E. Calabi. *Geom. Dedicata*, 11:387–396, 1981.
- [7] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, 2001.
- [8] William Goldman. Projective geometry on manifolds. Lecture notes, <http://www2.math.umd.edu/~wmg/pgom.pdf>, 1988.
- [9] William Goldman. Convex real projective structures on compact surfaces. *J. Differential Geom.*, 31:791–845, 1990.
- [10] François Labourie. Flat projective structures on surfaces and cubic holomorphic differentials. *Pure Appl. Math. Q.*, 3:1057–1099, 2007.
- [11] Jaejeong Lee. Convex fundamental domains for properly convex real projective structures. <http://users.math.yale.edu/~jl232/>.
- [12] An Min Li. Calabi conjecture on hyperbolic affine hyperspheres. II. *Math. Ann.*, 293:485–493, 1992.
- [13] John Loftin. Applications of affine differential geometry to \mathbb{RP}^2 surfaces. Ph.D. Thesis, <http://andromeda.rutgers.edu/~loftin>.
- [14] John Loftin. Affine spheres and convex \mathbb{RP}^n manifolds. *Amer. J. Math.*, 123:255–274, 2001.
- [15] John Loftin. The compactification of the moduli space of convex \mathbb{RP}^2 surfaces. I. *J. Differential Geom.*, 68:223–276, 2004.
- [16] John Loftin. Survey on affine spheres. In *Handbook of geometric analysis*, volume 13 of *Adv. Lect. Math.*, pages 161–191. 2010.
- [17] Ludovic Marquis. Espaces des modules des surfaces projectives proprement convexes de volume fini. *Geometry and Topology*, 14:2103–2149, 2010.
- [18] Ludovic Marquis. Surface projective convexe de volume fini. *Ann. Inst. Fourier (Grenoble)*, 62(1):325–392, 2012.
- [19] Nomizu and Sasaki. *Affine differential geometry*. Cambridge University Press, 1994.
- [20] Marc Troyanov. The Schwarz lemma for nonpositively curved Riemannian surfaces. *Manuscripta Math.*, 72:251–256, 1991.
- [21] Neil Trudinger and Xu-Jia Wang. The Monge–Ampère equation and its geometric applications. In *Handbook of geometric analysis*, Adv. Lect. Math. (7), pages 467–524. 2008.

- [22] Chang Ping Wang. Some examples of complete hyperbolic affine 2-spheres in \mathbb{R}^3 . L.N.M. 1481, pages 271–280. Springer, 1991.
- [23] Shing Tung Yau. Harmonic functions on complete Riemannian manifolds. *Comm. Pure Appl. Math.*, 28:201–228, 1975.