

FURSTENBERG MAPS FOR CAT(0) TARGETS OF FINITE TELESCOPIC DIMENSION

URI BADER, BRUNO DUCHESNE AND JEAN LÉCUREUX

ABSTRACT. We consider actions of locally compact groups G on certain CAT(0) spaces X by isometries. The CAT(0) spaces we consider have finite dimension at large scale. In case B is a G -boundary, that is a measurable G -space with amenability and ergodicity properties, we prove the existence of equivariant maps from B to the visual boundary ∂X .

1. INTRODUCTION

Furstenberg maps, or boundary maps, first appeared in H. Furstenberg's work [Fur63, Fur73]. These maps proved to be powerful tools for rigidity results. Indeed, the existence of such Furstenberg maps is used in particular in order to prove commensurator rigidity [AB94] or superrigidity phenomena [Mar91].

Our main topic of investigation in this paper is the existence of Furstenberg maps in the context of actions of groups on CAT(0) spaces. Recall that a CAT(0) space is a complete metric space that is non-positively curved in a way defined via the Bruhat-Tits inequality [BH99, p. 163].

We do not restrict ourselves to locally compact CAT(0) spaces but we replace this hypothesis by a condition at large scale. A CAT(0) space has finite telescopic dimension if any of its asymptotic cones has finite geometric dimension. Recall that a CAT(0) space has finite geometric dimension if there is a finite upper bound on the topological dimensions of its compact subspaces. Geometric dimension was introduced by B. Kleiner [Kle99] and telescopic dimension by P.-E. Caprace and A. Lytchack [CL10].

For example, CAT(0) cube complexes with an upper bound on the dimensions of cubes, Euclidean buildings — not necessarily locally compact nor with discrete affine Weyl groups — and infinite dimensional symmetric spaces with non-positive operator curvature and finite rank [Duc12] have finite telescopic dimension.

Our main result is the proof of existence of Furstenberg maps for such spaces.

Theorem 1.1. *Let X be a CAT(0) space of finite telescopic dimension and let G be a locally compact second countable group acting continuously by isometries on X without invariant flats. If (B, ν) is a G -boundary then there exists a measurable G -map from B to ∂X .*

Roughly speaking a G -boundary is a measurable space with an amenable G -action and strong ergodic properties (see §4 for a precise definition). In case G is amenable, B can be chosen to be a point and our theorem reduces to the following statement [CL10]: If G acts continuously by isometries on a CAT(0) space of finite telescopic dimension without invariant flat subspace then there is a fixed point at infinity, that is there is a (trivial) map from B to ∂X . Our tools to pass from that last statement to Theorem 1.1 are *measurable fields of CAT(0) spaces* over B , see §3.

In this theorem, the G -boundary can be taken to be any Poisson boundary of G (for an admissible measure). However, the theorem can be proved for an *a priori* larger class of spaces B , with suitable ergodic properties. Our work relies heavily on a very strong ergodic property of the boundary, namely *relative metric ergodicity*. This property was introduced by the first author together with A. Furman in [BF13], where (generalizing results of [BM02] and [Kai03]) it is proved that the Poisson boundary has indeed this strong ergodic property. For the sake of completeness, we also include a proof in this paper.

Date: April 11, 2014.

We warmly thank Alex Furman who kindly permit us to include the material of §4.

We also prove a similar result for *boundary pairs*, see Theorem 5.2. For example, Poisson boundaries associated to forward and backward random walks form a boundary pair.

Another possible approach to prove Theorem 1.1, at least for the Poisson boundary associated to some random walk on G , would be to try to understand the behavior of the random walk itself. More precisely, if Z_n is the n -th step of the random walk in G , and $o \in X$, one could hope that $Z_n.o$ converges to a point in the visual boundary. This would give a measurable, G -equivariant map from B to ∂X .

It is known that this approach works in some cases by the work of A. Karlsson and G. Margulis [KM99, Theorem 2.1]. However, there is a crucial assumption in this theorem, namely, that the drift is positive. Our result doesn't rely on this hypothesis.

In general, it is a natural question to relate the map obtained by Theorem 1.1 and the random walk in the space X . For example, if the drift is positive, and with finite first moment, we know that the random walk converges to a boundary point, and one can wonder whether the boundary map can be far away (in the sense of the Tits metric) from this limit. In case X is Gromov hyperbolic, the two essentially coincide. This is not the case for a symmetric space for example, as there are several different boundary maps.

In the same vein, one can ask whether the boundary map can be a (measurable) isomorphism, thus providing a geometric identification of the Poisson-Furstenberg boundary. The main tools for this possible application would be V. Kaimanovich's ray and strip criteria [Kai00, Theorems 5.5 & 6.4].

Finally, it also turns out that the methods on which our paper relies are also useful to answer a natural question raised by an implicit argument in [CL10]. Namely, in Appendix A, we prove the following result of non-emptiness for boundaries of CAT(0) spaces of finite telescopic dimension. Quite surprisingly, its proof relies on the existence of (G, μ) -boundaries for countable groups G and the dichotomy given between existence of Furstenberg maps and existence of Euclidean subfields given by Theorem 5.1.

Theorem 1.2. *Let X be a CAT(0) space of finite telescopic dimension. If $\text{Isom}(X)$ has no fixed point then the visual boundary ∂X is not empty.*

A CAT(0) space X is called *boundary minimal* [CM09, §1.B] if there is no non-empty closed convex subspace $Y \subset X$ such that $\partial Y = \partial X$. For proper CAT(0) spaces with boundary of finite dimension, minimality of X (that is no non-trivial closed convex subset is invariant under $\text{Isom}(X)$) implies boundary minimality [CM09, Proposition 1.5]. The same holds for CAT(0) spaces of finite telescopic dimension.

Corollary 1.3. *Let X be a minimal CAT(0) space of finite telescopic dimension not reduced to a point. Then X is boundary minimal.*

2. CAT(0) GEOMETRY

2.1. CAT(0) spaces of finite telescopic dimension. All along this text, we will deal with CAT(0) spaces of finite telescopic dimension as introduced in [CL10]. For reader's convenience, we recall facts about these spaces which will be useful for us.

We will start by defining dimension in a metric way, by Jung's Theorem [Jun01]. It is shown in the aforementioned paper that for any bounded subspace Y of a Euclidean space of dimension n

$$(1) \quad \text{rad}(Y) \leq \sqrt{\frac{n}{2(n+1)}} \text{diam}(Y)$$

and there is equality if and only if the closure of Y contains a regular n -simplex of diameter $\text{diam}(Y)$.

Using this inequality, P.-E. Caprace and A. Lytchack showed [CL10, Theorem 1.3] that a CAT(0) space Y has *geometric dimension* (as introduced by B. Kleiner [Kle99]) at most n if and only if for any bounded subset $Y \subseteq X$, Inequality (1) holds.

Telescopic dimension is a notion at large scale. More precisely, a CAT(0) space X has telescopic dimension at most n if and only if any asymptotic cone of X has geometric dimension at most n . It can be expressed quantitatively: a CAT(0) space X has telescopic dimension at most n if and only if for any $\delta > 0$, there is $D > 0$ such that for any bounded subset $Y \subseteq X$ of diameter larger than D we have

$$(2) \quad \text{rad}(Y) \leq \left(\delta + \sqrt{\frac{n}{2(n+1)}} \right) \text{diam}(Y).$$

Note that a locally compact CAT(0) space may have infinite telescopic dimension.

One main feature of CAT(0) spaces of finite telescopic dimension is the following: if the intersection of a filtering family $\{X_\alpha\}$ of closed convex subsets is empty then the intersection of boundaries $\cap_\alpha \partial X_\alpha$ is not empty and there is a canonical point ξ in this intersection.

This canonical point is given by the fact that the boundary of a CAT(0) space of telescopic dimension at most n has geometric dimension at most $n-1$ and the fact that a CAT(1) space Ξ of finite geometric dimension and radius at most $\pi/2$ has such a canonical point. This point is defined as the unique circumcenter of the set of circumcenters of Ξ .

This property of filtering families of closed convex subspaces can be thought as a compactness property for $\bar{X} = X \cup \partial X$. Let us define the topology \mathcal{T}_c on \bar{X} as the weakest topology such that \bar{C} is \mathcal{T}_c -closed for any closed — for the usual topology — convex subset $C \subseteq X$ [Mon06, §3.7&§3.8]. Then for a CAT(0) space X of finite telescopic dimension \bar{X} is compact — not necessarily Hausdorff.

2.2. Geometry of flats in CAT(0) spaces. In the heart of the proof of Theorem 1.1 we will deal with a Euclidean subfield of a CAT(0) field. We gather in this subsection useful facts about the geometry of Euclidean subspaces (also called flats subspaces) in CAT(0) spaces.

Let (X, d) be a CAT(0) space. If C is a closed convex subset of X , we denote by d_C the distance function to C , that is $d_C(x) = \inf_{c \in C} d(c, x) = d(x, \pi_C(x))$ where $\pi_C(x)$ is the projection of x on C .

Let Y be a bounded subset of a metric space (Z, d) . The *circumradius* (or simply *radius*) $\text{rad}(Y)$ of Y is the non-negative number $\inf_{z \in Z} \sup_{y \in Y} d(y, z)$, its *intrinsic circumradius* is $\inf_{z \in Y} \sup_{y \in Y} d(y, z)$ and a *circumcenter* of Y is a point in Z minimizing $\sup_{y \in Y} d(y, \cdot)$.

More generally, we define a *center* of Y as a point of Z which is fixed by any isometry of Z stabilizing Y . The Bruhat-Tits fixed point lemma asserts that, in a CAT(0) space, a bounded set has a unique circumcenter, which is therefore a center.

If S_1 and S_2 are two subsets of Euclidean spheres, we denote by $S_1 * S_2$ their spherical join [BH99, Definition I.5.13]. This is the spherical analogue of Euclidean products. Such Euclidean products appear, for example, in the de Rham decomposition [BH99, Theorem II.6.15] of the CAT(0) space X : the space X is isometric to a product $H \times Y$ where H is a Hilbert space and Y that does not split with a Euclidean factor. Isometries of X preserve this decomposition acting diagonally on $H \times Y$ and the choice of a base point in X allows us to identify H and Y with closed convex subspaces of X containing this point.

The following lemma details the possibilities for a convex function on a Euclidean space. It will be applied in two situations: the restriction of a Busemann function to a Euclidean subspace of X and to the restriction to a Euclidean subspace of the distance function to another Euclidean subspace.

Proposition 2.1. *Let E be a Euclidean space, f be a convex function on E and $m = \inf\{f(x) \mid x \in E\}$. Then :*

- (i) *If m is not attained, then the intersection $\bigcap_{\varepsilon > 0} \partial(f^{-1}((m, m + \varepsilon)))$ is not empty and has a center.*

If m is a minimum, let $E_m = f^{-1}(\{m\})$. Let $E_m = F_m \times T$ be its de Rham decomposition, where F_m is a maximal flat contained in E_m . Then exactly one of the following possibilities happen.

- (ii) *E_m is bounded and thus has a center (i.e. F is a point and T is compact),*
- (iii) *T is bounded and $\partial E_m = \partial F_m$ is a sphere,*
- (iv) *T is unbounded and its boundary ∂T has radius less than $\pi/2$ and thus has a center.*

Proof. If m is not a minimum then the net of closed convex subsets $\left(f^{-1}((m, m + \varepsilon))\right)_{\varepsilon > 0}$ has empty intersection and the result follows from \mathcal{T}_c -compactness.

The other cases coincide with the join decomposition $\partial E_m = \partial F * \partial T_m$, using Lemma 2.2 below where ∂E_m will correspond to C , ∂F_m to S_1 and ∂T to C_2 . \square

Lemma 2.2. *Let (S, d) be a Euclidean sphere. Let C be a non-empty closed convex subset of S and let S_0 be the minimal subsphere of S containing C . Then there is a unique decomposition of S_0 as a spherical join $S_0 = S_1 * S_2$ where S_1 and S_2 are reduced to a point or are subspheres of S_0 such that*

$$C = S_1 * C_2$$

where C_2 is a closed convex subset of S_2 with intrinsic radius $< \frac{\pi}{2}$.

In particular, any closed convex subset that is not a subsphere has a center (the one of C_2). Moreover, it coincides with the unique circumcenter of the set of circumcenters of C .

Proof. First observe that intersections of subspheres are empty or subspheres themselves. This yields the existence of S_0 . In the same way, convex hulls of subspheres are subspheres and thus there exists a maximal subsphere S_1 contained in C . Let S_2 be the set of points of S_0 at distance $\pi/2$ from every points in S_1 . Then $S_0 = S_1 * S_2$. Any point of C can be written (x_1, x_2, α) with $x_i \in S_i$ and $\alpha \in [0, \pi/2]$. Since C is convex and $S_1 \subseteq C$, there exists C_2 , which is $S_2 \cap C$, such that $C = S_1 * C_2$. Observe that C_2 does not contain any sphere by maximality of S_1 . Now C has diameter $< \pi$ — otherwise, it contains at least a sphere of dimension 0 (that is two antipodal points) — thus has also intrinsic circumradius $< \pi/2$.

Observe that C is not a sphere if and only if C_2 is not empty. In the case where S_1 and C_2 are not empty then any point of C_2 is a circumcenter since in that case, the intrinsic circumradius is $\pi/2$. The fact that any convex subset of circumradius $< \pi/2$ has a unique circumcenter implies the last sentence of the lemma. \square

There is a particular situation for the relative position of two Euclidean subspaces E, F in X : when the restriction on F of the distance to E is constant and vice-versa. In that situation E and F are said to be *parallel*. The Sandwich Lemma [BH99, Exercise II.2.12(2)] implies that their convex hull splits isometrically as $\mathbb{R}^n \times [0, d]$. In particular, E and F are isometric and thus have same dimension n .

Lemma 2.3. *Let E be a Euclidean subspace of X and let Y be the union of subspaces parallel to E . Then Y is a closed convex subspace of X . A point $y \in X$ belongs to Y if and only if for any $x_1, \dots, x_n \in E$, $\text{Conv}(y, x_1, \dots, x_n)$ is isometric to a convex subset of a Euclidean space.*

Let p be the restriction to Y of the projection to E . Fix some $x \in E$ and let Z be $p^{-1}(\{x\})$. Then Z is a closed convex subspace of X and Y decomposes isometrically as $E \times Z$.

Proof. The lemma is proved when E is a line in [BH99, Proposition II.2.14]. Let n be the dimension of E . We proceed by induction on n . Assume this is true for $n - 1$ and choose an orthogonal splitting $F \times L$ of E where F has dimension $n - 1$, L is a line and $F \cap L = \{x\}$. The induction assumption for F implies the convex hull of subspaces parallel to F splits isometrically as $F \times Z_F$.

Now let us apply the case $n = 1$ for the union of lines parallel to L in Z_0 which splits as $L \times Z_0 \subseteq Z_F$. It is clear that $E \times Z_0 \subset Y$.

If E' is a Euclidean subspace parallel to E then the restriction p' of p to E' is an isometry. Set $F' = p'^{-1}(F)$ and L' to be the orthogonal line to F' containing $x' = p'^{-1}(x)$. For any $y \in L'$, $F \times \{y\}$ is parallel to F and thus $L' \subset Z_F$. Since L' is parallel to L , $x' \in Z_0$. Finally $Z_0 = Z$ and Y splits isometrically as $E \times Z$.

Assume that for any $x_1, \dots, x_n \in E$, $\text{Conv}(y, x_1, \dots, x_n)$ is isometric to a convex subset of a Euclidean space. Then $\text{Conv}(\{y\} \cup E)$, which is the union of such spaces, satisfies the following property: for any $z_1, \dots, z_n \in \text{Conv}(\{y\} \cup E)$, $\text{Conv}(z_1, \dots, z_n)$ is isometric to a convex subset of a Euclidean space. This property is a characterization of CAT(0) spaces that are isometric to a convex subset of a Hilbert space. In particular, in our case $\text{Conv}(\{y\} \cup E)$ is necessarily isometric to $\mathbb{R}^d \times d(y, E)$ where $d = \dim(E)$. \square

Lemma 2.4. *Let E be a Euclidean subspace of X and $\xi \in \partial X$ such that the Busemann function β_ξ associated to ξ (with respect to some fixed base point) is constant on E . If $x \in E$ and ρ is the ray from x to ξ then the convex hull of $E \cup \rho$ is isometric to $E \times [0, +\infty)$.*

In particular, with the notations of Lemma 2.3, $\xi \in \partial Z$.

Proof. Choose $x \in E$ and y on the geodesic ray from x to ξ . We claim that $\overline{\angle}_x(y, z) = \pi/2$ for any point $z \neq x$ in E . Assume $\overline{\angle}_x(y, z) < \pi/2$. Then there is $x' \in (x, z)$ such that $d(y, x') < d(y, x)$. This implies $\beta_\xi(x') < \beta_\xi(x)$, which is a contradiction. Arguing the same way with the symmetric of z with respect to x , we get the claim. In particular, for any $x, x' \in E$, $\angle_x(x', \xi) + \angle_{x'}(x, \xi) = \pi$ and [BH99, Proposition II.9.3] implies that the convex hull of x, x' and ξ is isometric to $[0, d(x, x')] \times [0, \infty)$.

This shows that the projection of E on any closed horoball is a flat parallel to E and is the lemma is now a consequence of Lemma 2.3. \square

3. MEASURABLE FIELDS OF COMPLETE SEPARABLE METRIC SPACES

3.1. Metric fields. An important — although slightly technical — tool in our proof will be the notion of fields of metric spaces. Roughly speaking a measurable field of metric spaces over a measurable space A is a way to attach measurably a metric space to any point in A . Thanks to [Val77] one can think to measurable fields of metric spaces over A in the following way: to each point of A one associates a closed subspace of a fixed metric space, namely the Urysohn space.

Let (A, η) be a Lebesgue space, that is a standard Borel space with a Borel probability measure [Kec95, §12]. All our definitions will only depend on the class of the measure η .

Definition 3.1. Let $\{X_a\}_{a \in A}$ be a collection of complete separable metric spaces. The distance on X_a is denoted d_a or simply d if there is no ambiguity. Measurability conditions are defined thanks to the notion of fundamental families. A *fundamental family* $\mathcal{F} = \{x^n\}$ is a countable set of elements of $\prod_{a \in A} X_a$ with the following properties

- $\forall n, m, a \mapsto d_a(x_a^n, x_a^m)$ is measurable,
- for almost every $a \in A$, $\{x_a^n\}$ is dense in X_a .

A *measurable field \mathbf{X} of complete separable metric spaces* — or simply a *metric field* for short — is then the collection of data: (A, η) , $\{X_a\}_{a \in A}$ and $\{x^n\}$.

A *section* of \mathbf{X} is an element $x \in \prod_a X_a$ such that for all $y \in \mathcal{F}$, $a \mapsto d_a(x_a, y_a)$ is measurable.

Two sections are identified if they agree almost everywhere. The set of all sections is the *measurable structure \mathcal{M} of \mathbf{X}* . If x, y are two sections, the equality

$$d_a(x_a, y_a) = \sup_{z \in \mathcal{F}} |d_a(x_a, z_a) - d_a(z_a, y_a)|$$

shows that $a \mapsto d_a(x_a, y_a)$ is also measurable.

Let G be a second countable locally compact group. The Lebesgue space (A, η) is a G -space if G acts by measure class preserving automorphisms on A and the map $(g, a) \mapsto ga$ is measurable.

Definition 3.2. Let (A, η) be a G -space. A *cocycle* for G on \mathbf{X} is a collection $\{\alpha(g, a)\}_{g \in G, a \in A}$ such that

- for all g and almost every a , $\alpha(g, a) \in \text{Isom}(X_a, X_{ga})$;
- for all g, g' and almost every a , $\alpha(gg', a) = \alpha(g, g'a)\alpha(g', a)$;
- for all $x, y \in \mathcal{F}$, the map $(g, a) \mapsto d_a(x_a, \alpha(g, g^{-1}a)y_{g^{-1}a})$ is measurable.

In that case we say that G *acts* on \mathbf{X} via the cocycle α or that there is an *action* of G on \mathbf{X} . A section x is invariant if for all g and almost all a , $\alpha(g, g^{-1}a)x_{g^{-1}a} = x_a$.

3.2. Fields of CAT(0) spaces. A special case of metric fields is the case of CAT(0) fields. The theory of measurable fields of complete separable metric spaces and more specifically of CAT(0) spaces appeared in [AH14, Duc13] and references therein.

Definition 3.3. Let \mathbf{X} be a metric field and $\kappa \in \mathbb{R}$. We say that \mathbf{X} is a $\text{CAT}(\kappa)$ field if for almost every a , X_a is a $\text{CAT}(\kappa)$ space.

A *subfield* \mathbf{Y} of a $\text{CAT}(0)$ field \mathbf{X} is a collection $\{Y_a\}_{a \in A}$ of non-empty closed convex subsets such that for every section x of \mathbf{X} , the function $a \mapsto d(x_a, Y_a)$ is measurable. We identify subfields \mathbf{Y} and \mathbf{Y}' if $Y_a = Y'_a$ for almost every a .

Similarly we speak of *Euclidean fields* and subfields of such fields. If a group G acts on \mathbf{X} , a subfield \mathbf{Y} is *invariant* if for all g and almost all a , $\alpha(g, g^{-1}a)Y_{g^{-1}a} = Y_a$.

In $\text{CAT}(1)$ spaces, subsets of circumradius less than $\pi/2$ are strictly convex and admits a unique circumcenter as well, see [BH99, Proposition II.2.7] or [LS97, Proposition 3.1] for quantitative statements. Those circumcenters can be defined canonically by means of the metric structure and arguing as in [Duc13, Lemma 8.7] we obtain the following.

Lemma 3.4. *Let \mathbf{X} be a $\text{CAT}(1)$ field. If \mathbf{C} is a subfield of \mathbf{X} with fibers of radius less than $\pi/2$ then the family of circumcenters of \mathbf{C} is a section of \mathbf{X} .*

Lemma 3.5. *Let \mathbf{X} be a metric field and x be a section of it. For $a \in A$, let B_a^r be the closed ball of radius r around x_a then $\mathbf{B}^r = (B_a^r)$ is a metric field. Moreover if G acts on \mathbf{X} and x is an invariant section then G acts on \mathbf{B}^r as well.*

Proof. This statement is almost a straightforward verification of the definitions. The only non completely obvious fact is maybe the construction of a fundamental family. Fix a fundamental family x^n of \mathbf{X} . For $n \in \mathbb{N}$, $a \in A$, set inductively k_a^n to be $\min\{k > k_a^{n-1}; x_a^k \in \overline{B}(x_a, r)\}$. Let us denote by y_a^n the point $x_a^{k_a^n}$. The family (y^n) is a fundamental family of \mathbf{B}^r . \square

Building on Lemma 2.3 we obtain the following measurable decomposition of the union of flats parallel to a Euclidean subfield.

Lemma 3.6. *Let \mathbf{X} be a $\text{CAT}(0)$ field and \mathbf{E} be a Euclidean subfield. For $a \in A$ let Y_a be the union of flats parallel to E_a . Then $\mathbf{Y} = (Y_a)$ is a subfield of \mathbf{X} which splits as a product of $\text{CAT}(0)$ fields $\mathbf{Y} = \mathbf{E} \times \mathbf{Z}$. Moreover if G acts on \mathbf{X} and \mathbf{E} is invariant then \mathbf{Y} is invariant and G acts diagonally on $\mathbf{E} \times \mathbf{Z}$ with an invariant section in \mathbf{Z} .*

Proof. Fix $a \in A$. Thanks to Lemma 2.3, the condition $y \in Y_a$ can checked only using distances $d(y, x_a^n)$ where (x^n) is a fundamental family of the subfield \mathbf{E} . One can then readily check that \mathbf{Y} is a subfield of \mathbf{X} . Fixing a section x of \mathbf{E} one can recover \mathbf{Z} as $p_a^{-1}\{x\}$ where p_a is the projection on E_a . \square

Lemma 3.7. *Let \mathbf{E} be a Euclidean field. The map $a \mapsto \dim(E_a)$ is measurable.*

Proof. Fix a fundamental family $\{x^n\}$ for \mathbf{E} . Thanks to Jung's Inequality (1) the dimension d of E_a can be obtained via the quantity $\sqrt{\frac{d}{2(d+1)}}$ which is the minimal non-negative number K such that $\text{rad}(\{x_a^{n_1}, \dots, x_a^{n_k}\}) \leq K \text{diam}(\{x_a^{n_1}, \dots, x_a^{n_k}\})$ where $n_1, \dots, n_k \in \mathbb{N}$. \square

Any CAT(0) space X has a visual boundary ∂X (which may be empty). Making this construction pointwise, we can consider, at least in a set-theoretic way, the boundary field of a CAT(0) field \mathbf{X} .

However, the measurable structure is not so clear. We would need a separable metric on each fiber. One way to endow the boundary of a CAT(0) space with a metric is to consider the angle metric, or the Tits metric which is the length metric associated to the previous one. These are invariant metrics but then ∂X is not separable in general — for example, the boundary of the hyperbolic plane, endowed with the Tits metric, is an uncountable discrete space.

Actually there is no way to construct a boundary field $\partial\mathbf{X}$ for a CAT(0) field \mathbf{X} such that a group acting on \mathbf{X} also acts (isometrically) on $\partial\mathbf{X}$. Otherwise, any Furstenberg map given by Theorem 1.1 would be constant because of double metric ergodicity.

One way to avoid these problems, is to let down the desired invariance of the metric. Recall that a separable CAT(0) space X embeds continuously (not isometrically) to a subset of the Fréchet space $\mathcal{C}(X)$ of continuous functions on X endowed with the distance $d(f,g) = \sum_{n \in \mathbb{N}} 2^{-n} \frac{|f(x_n) - g(x_n)|}{1 + |f(x_n) - g(x_n)|}$ where $\{x_n\}$ is a dense subset of X . This metric topology coincides with the topology of pointwise convergence. More precisely, if x_0 is a base point in X , the embedding is given by $\iota: y \mapsto d(\cdot, y) - d(y, x_0)$. The closure of $\iota(X)$ is a compact metric space which allows us to define a *bordification field* \mathbf{K} for a CAT(0) field \mathbf{X} (see [Duc13, §9.2]). Sections of \mathbf{K} corresponds to some collections of Busemann functions. We define them as follows.

Definition 3.8. Let \mathbf{X} be a CAT(0) field. We define its *boundary field* $\partial\mathbf{X}$ to be the set of sections f of its bordification field \mathbf{K} such that for a.e. $a \in A$ there is $\xi_a \in \partial X_a$ with $f_a = \beta_{\xi_a}(\cdot, x_0)$ where β_{ξ_a} is the Busemann function associated to $\xi_a \in \partial X_a$.

By an abuse of notations we will say that $\xi = (\xi_a)$ is a section of the boundary field. Observe in that case for all x, y sections of \mathbf{X} , the function $a \mapsto \beta_{\xi_a}(x_a, y_a)$ is measurable. Even if we will not need it we observe that one can decide if a section of \mathbf{K} corresponds to a section of the boundary field.

Lemma 3.9. Let f be a section of \mathbf{K} . Let A' be the set of elements $a \in A$ such that there is $\xi_a \in \partial X_a$ with $f_a = \beta_{\xi_a}(\cdot, x_0)$. Then A' is a measurable subset of A .

Proof. We use the fact that f_a coincides with a Busemann function if it is a limit of points in $\iota(X_a)$ for the topology of uniform convergence on bounded subsets [BH99, §II.8]. Fix x^n a fundamental family of \mathbf{X} and $x^n(r)$ fundamental families of the fields \mathbf{B}^r of closed balls around x^0 . Now, $a \in A'$ if and only for any $r > 0$

$$\inf_{\{n \in \mathbb{N}; d(x_a^n, x_a^0) > r\}} \sup_{m \in \mathbb{N}} |f_a(x_a^m(r)) - \iota(x_a^n)(x_a^m(r))| = 0.$$

□

The following lemma shows that our notion of boundary field gives something natural in case of a constant field.

Lemma 3.10. Let (X, d) be a complete separable metric space and let us denote by \mathbf{X} the measurable field over (A, η) with constant fibers equal to X . Sections of \mathbf{X} and measurable maps $A \rightarrow X$ are in bijective correspondence. Moreover if X is CAT(0) space then sections of $\partial\mathbf{X}$ and measurable maps $A \rightarrow \partial X$ are in bijective correspondence.

Proof. Recall that a fundamental family of \mathbf{X} is given by constant maps $a \mapsto x_n$ where (x_n) is a countable dense subset of X . Now a map $f: A \rightarrow X$ is measurable if and only if for any x , $a \mapsto d(x, f(a))$ is measurable if and only if for any n , $a \mapsto d(x_n, f(a))$ is measurable.

Now consider the case where X is a CAT(0) space. We use the identification of the boundary of X with the set of Busemann functions vanishing at some base point x_0 . In this identification, the cone topology corresponds to the topology of uniform convergence on closed balls around x_0 . In particular, a map $f: A \rightarrow \partial X$ is measurable if and only if for any $x, y \in X$, $a \mapsto \beta_{f(a)}(x_0, y)$ is measurable if and only if for any $n, m \in \mathbb{N}$, $a \mapsto \beta_{f(a)}(x_n, x_m)$ is measurable. □

Among CAT(0) spaces, Euclidean spaces have a special feature: the angle between two points at infinity is the same from any point you look at them. This gives a distance at infinity independent of the choice of a base point and this distance is invariant under the action of the isometry group of the Euclidean space. In case of a field of Euclidean spaces we get the following fact.

Lemma 3.11. *Let \mathbf{E} be a measurable field of Euclidean spaces. The boundary field $\partial\mathbf{E}$ has a structure of a field of CAT(1) spaces for the Tits metric. If G acts isometrically on \mathbf{E} then it acts also isometrically on $\partial\mathbf{E}$.*

Proof. Each ∂E_a is a complete separable metric space (isometric to a Euclidean sphere) for the Tits metric, and the action of G preserves this distance. What is left to do is to check the measurable structure.

We choose a fundamental family (x^n) of \mathbf{E} such that there is a section x^0 such that for any n and almost all a , $d(x_a^0, x_a^n) \neq 0$. Now define $\xi_a^n \in \partial E_a$ to be the end point of $[x_a^0, x_a^n]$. We claim that (ξ^n) is a fundamental family for $\partial\mathbf{E}$. We define

$$m(a, n, k) = \min \left\{ m; d(x_a^0, x_a^m) > k, |d(x_a^0, x_a^n) + d(x_a^n, x_a^m) - d(x_a^0, x_a^m)| < 1 \right\}$$

and

$$y_a^{(n,k)} = x_a^{m(a,n,k)}.$$

This way, for any (n, k) , $y^{(n,k)}$ is a section and for almost every a , $y_a^{(n,k)} \rightarrow \xi_a^n$ as $k \rightarrow \infty$. This shows that for any $n, m, a \mapsto \angle(\xi_a^n, \xi_a^m) = \lim_{k \rightarrow \infty} \angle_{x_a^0}(y_a^{(n,k)}, y_a^{(m,k)})$ is measurable. \square

Remark 3.12. Let E be a Euclidean space and ξ, η be two points at infinity. The very special geometry of E implies the following formula between Busemann functions β_ξ, β_η and the visual angle $\angle_{x_0}(\xi, \eta)$ (which does not depend on x_0 and is also the Tits angle) :

$$\sup_{1/2 < d(x, x_0) < 1} \frac{|\beta_\xi(x, x_0) - \beta_\eta(x, x_0)|}{d(x, x_0)} = 2(1 - \cos(\angle(\xi, \eta))).$$

This formula shows that in the case of a measurable field of Euclidean spaces \mathbf{E} , the notion of section of $\partial\mathbf{E}$ defined in Definition 3.8 and the notion of section for the structure of a metric field introduced in Lemma 3.11 coincide.

Let E be a Euclidean space of dimension d_0 . It is not hard to define a distance on the set S of subspheres of dimension $0 \leq d < d_0$ in ∂E turning S into a complete separable metric space. The following lemma does the same in a measurable context.

Lemma 3.13. *Let \mathbf{E} be a Euclidean field of constant dimension d_0 . Let d be a positive integer less than d_0 . For any $a \in A$, let S_a be the set of subspheres of dimension d of ∂E_a . The collection $\mathbf{S} = (S_a)$ has a structure of a metric field. If G acts on \mathbf{E} then it acts also on \mathbf{S} (isometrically).*

Let E_a^s be the set of Euclidean subspaces F of E_a such that $\partial F = s_a$. Then $\mathbf{E}^s = (E_a^s)$ has a natural structure of Euclidean field such that any section of \mathbf{E}^s corresponds to a Euclidean subfield of \mathbf{E} .

Moreover if G acts on \mathbf{E} and s is an invariant section of \mathbf{S} then G acts on \mathbf{E}^s .

Proof. First, we claim that one can construct a fundamental family (x^n) of \mathbf{E} such that for any choice n_0, \dots, n_d , and almost all a , $x_a^{n_0}, \dots, x_a^{n_d}$ is not included in a Euclidean subspace of dimension $< d$ since this condition can be checked only with distances.

For all a , we define S_a to be the set of subspheres of ∂E_a of dimension d . For $s^1, s^2 \in S_a$, we define $d(s^1, s^2)$ to be the Hausdorff distance between two compact subspaces associated to the Tits distance on ∂E_a . Now, for a choice of $N = \{n_0, \dots, n_d\}$ we define s_a^N to be the boundary of the affine span of $x_a^{n_0}, \dots, x_a^{n_d}$. There are countably many possibilities for N and $\{(s_a^N)\}_N$ defines a fundamental family of \mathbf{S} . If G acts on \mathbf{E} then it acts on $\partial\mathbf{E}$ and since the Hausdorff distance is defined via supremum of some Tits angles, G acts (measurably) on \mathbf{S} .

Let s be a section of \mathbf{S} and for any x^n element of the fundamental family of \mathbf{E} let F_a^n be the unique Euclidean subspace of E_a containing x_a^n such that $\partial F_a^n = s_a$.

The last statement comes from the fact that in that case for any $g \in G$ and all almost $a \in A$, $\alpha(g, a)E_a^s = E_{ga}^{\alpha(g,a)s_a} = E_{ga}^{s_{ga}}$. \square

4. METRIC ERGODICITY AND ITS RELATIVE VERSION

In [BF12] U. Bader and A. Furman introduced the notion of a boundary pair, which is further developed in their yet unpublished paper [BF]. In this section we review this theory.

We fix a locally compact second countable group G for the rest of this section. All conditions of measurability in G will be relative to the Haar measure class.

Definition 4.1 ([BF13, Definition 4.1]). Let (A, η) be a G -space. The action $G \curvearrowright (A, \eta)$ is *metrically ergodic* if for any action of G by isometries on a complete separable metric space (X, d) , any G -equivariant measurable map from $A \rightarrow X$ is essentially constant.

If the diagonal action $G \curvearrowright A \times A$ is metrically ergodic, we say that $G \curvearrowright A$ is *doubly metrically ergodic*.

Remark 4.2. We will only use complete separable metric spaces. However, this is not an important restriction. About completeness, one may consider the extended action on the completion \overline{X} and observe that $\overline{X} \setminus X$ has zero measure. Moreover one may reduce to separable spaces as the following argument shows.

Let f be a measurable map $A \rightarrow X$. Thanks to [Fre81], f is actually η -measurable in N. Bourbaki's sense and thus separably valued (see [Bou04, IV §5 No 5, Theorem 4]), meaning that there is $X' \subseteq X$ closed and separable such that $f_*\eta(X \setminus X') = 0$. Now, using separability of X' , it can be easily checked that the set $S = \{x \in X \mid \forall \varepsilon > 0, f_*\eta(B(x, \varepsilon)) > 0\}$ is a closed subset of X' which is G -invariant and satisfies $f_*\eta(X \setminus S) = 0$.

Below we present a *relative* notion of metric ergodicity as well [BF12]. The definition that we give here is not exactly the one given in the paper [BF12], but a version of it modified in order to fit in the context of measurable fields of complete separable metric spaces

Definition 4.3. Let (A, η) and (B, ν) be two Lebesgue spaces. A measurable map $\pi: A \rightarrow B$ is a *factor map* if $\pi_*\eta$ and ν are in the same measure class. If A and B are G -spaces and π is G -equivariant then we say that π is a *G -factor*.

Definition 4.4 ([BF]). Let (A, η) and (B, ν) be two Lebesgue spaces and $\pi: A \rightarrow B$ be a factor map. Let \mathbf{X} be a metric field over (B, ν) . A *relative section* is a map $\varphi: A \rightarrow \sqcup_{b \in B} X_b$ such that :

- for all $a \in A$, $\varphi(a) \in X_{\pi(a)}$,
- for any section x of \mathbf{X} , $a \mapsto d(x(\pi(a)), \varphi(a))$ is measurable.

If π is a G -factor and G acts on \mathbf{X} via a cocycle α , such a relative section is said to be *invariant* if for almost every a and all $g \in G$, $\varphi(ga) = \alpha(g, \pi(a))\varphi(a)$.

Definition 4.5 ([BF]). We say that the G -map $\pi: A \rightarrow B$ is *relatively metrically ergodic* (or equivalently $G \curvearrowright A$ is *metrically ergodic relatively to π*) if any invariant relative section coincides with a section. In other words, for any G -metric field \mathbf{X} and any invariant relative section φ , there is an invariant section x of \mathbf{X} such that for almost all $a \in A$, $\varphi(a) = x(\pi(a))$.

The following lemma shows how relative metric ergodicity implies metric ergodicity. Actually metric ergodicity of $G \curvearrowright A$ is equivalent to metric ergodicity of $G \curvearrowright A$ relatively to the projection to a point.

Lemma 4.6 ([BF]). Let (A, η) , (B, μ) be two G -spaces. If $A \times B \rightarrow B$ is relatively metrically ergodic then $G \curvearrowright A$ is metrically ergodic.

Proof. Let $\phi: A \rightarrow X$ be an equivariant G -map to some complete separable metric space. Consider \mathbf{X} be the trivial field $X \times B$ over B and define $\varphi(a, b) = \phi(a)$. The map φ is an invariant relative section and thus does not depend on a , that is ϕ is essentially constant. \square

Definition 4.7 ([BF]). Let (B_-, ν_-) , (B_+, ν_+) be G -spaces, we say that (B_-, B_+) is a *G -boundary pair* if

- the actions $G \curvearrowright B_+$ and $G \curvearrowright B_-$ are amenable in Zimmer's sense [Zim84],
- both first and second projections $B_- \times B_+ \rightarrow B_\pm$ are relatively ergodic.

A G -space (B, μ) is *G -boundary* if (B, B) is a G -boundary pair.

Let μ be a probability measure on G . Recall that a (G, μ) -space is a G -space (A, η) such that $\mu * \eta = \eta$, where the convolution measure $\mu * \eta$ is the pushforward measure of $\mu \times \eta$ under the map $(g, a) \mapsto ga$. Such a measure ν is called μ -harmonic or μ -stationary in the literature. Let $i: G \rightarrow G$ be the inversion given by $i(g) = g^{-1}$ for any $g \in G$. We denote by $\check{\mu}$ the probability measure $i_*\mu$. Recall that μ is *symmetric* if $\check{\mu} = \mu$.

For the remainder of this section, we fix a spread out non-degenerate probability measure μ on G , that is μ is absolutely continuous with respect to a Haar measure and its support generates G as semigroup. Let (B, ν) be the Poisson boundary associated to μ . We also denote by $(\check{B}, \check{\nu})$ the Poisson boundary of $(G, \check{\mu})$. We refer to [Fur63, Kai03] for notions and references about Poisson boundaries and related ergodicity properties. We emphasize that (B, ν) and $(\check{B}, \check{\nu})$ are respectively a (G, μ) -space and a $(G, \check{\mu})$ -space.

Theorem 4.8 ([BF]). *The pair (\check{B}, B) is a G -boundary pair.*

This will be deduced from the following statement.

Theorem 4.9 ([BF]). *Let (A, η) be a $(G, \check{\mu})$ -space. The factor map $A \times B \rightarrow A$ is relatively ergodic.*

Corollary 4.10 ([BF]). *The diagonal action of G on $(\check{B} \times B, \check{\nu} \times \nu)$ is metrically ergodic.*

Remark 4.11. The same argument as in Corollary 4.10 shows that if (B_-, B_+) is a G -boundary pair then $G \curvearrowright B_- \times B_+$ is metrically ergodic. In particular if B is a G boundary then $G \curvearrowright B$ is doubly metrically ergodic.

Proof of Corollary 4.10. Let U be a metric separable space on which G acts continuously and by isometries. Assume $f: \check{B} \times B \rightarrow U$ is a G -equivariant measurable map. Take $A = \check{B}$. It follows from the Theorem 4.9 that f only depends on the first coordinate.

Now consider the measure $\check{\mu}$. Then B is a $(G, \check{\mu})$ -space (in other words, a (G, μ) -space), so we can apply Theorem 4.9 to $B \times \check{B}$. This implies that f does not depend on the first coordinate.

Putting together the two results, we see that f does not depend on the first, nor on the second coordinate. Hence, f is constant. \square

The following proposition is a key tool in the proof of Theorem 4.8. It combines Poincaré recurrence theorem for A and SAT property for B . Recall that SAT, which means *strongly almost transitive*, is a weak mixing property introduced by W. Jarowski [Jaw94].

Theorem 4.12 ([BF]). *Let (A, η) be a $(G, \check{\mu})$ -space and $Y \subset A \times B$ be a set of positive $\eta \times \nu$ -measure. For $a \in A$, denote by Y_a the set $\{b \in B \mid (a, b) \in Y\}$. Then, for almost every $a \in \pi_1(Y)$, and for every $\varepsilon > 0$, there is a $g \in G$ such that*

- (i) $\nu(gY_a) > 1 - \varepsilon$
- (ii) $ga \in \pi_1(Y)$ where π_1 is the first projection $A \times B \rightarrow A$.

Proof. We will use the definition of the Poisson boundary as a space of ergodic components of the space of increments of the random walk $\Omega = G \times G^{\mathbb{N}}$ by the shift S .

We can define another shift T from $A \times \Omega$ to itself, defined by $T(a, g, \omega_1, \omega_2, \dots) = (g^{-1}a, \omega_1, \omega_2, \dots)$. Since A is a $(G, \check{\mu})$ -space, we have $\check{\mu} * \eta = \eta$, and we conclude that T preserves the measure $m := \eta \times \mu \times \mu^{\mathbb{N}}$ on $A \times \Omega$.

Let us consider the fiber product $X = A \times_G \Omega$. This is defined as the quotient space of $Y \times \Omega$ by the relation $(a, hg, \omega_1, \dots) \sim (h^{-1}a, g, \omega_1, \dots)$, for all $h \in G$. It follows that the space X is isomorphic to $A \times G^{\mathbb{N}}$. The pushforward of the measure on $A \times \Omega$ to X is simply the measure $\eta \times \mu^{\mathbb{N}}$. Furthermore, the shift T preserves the equivalence relation, so it still acts on X , and also preserves the measure.

Let $Y \subset A \times B$ be such that $\eta \times \nu(Y) > 0$. We can consider the preimage of Y in $A \times \Omega$, and then push it forward to get a subset \tilde{Y} of X . First we note that, by Poincaré recurrence theorem, we have that, for almost every $x \in \tilde{Y}$, there are infinitely many $n \in \mathbb{N}$ such that $T^n x \in \tilde{Y}$.

If $a \in A$, define the set $\tilde{Y}_a = \{\omega \in G^{\mathbb{N}} \mid (a, \omega) \in \tilde{Y}\}$. By Fubini, we have $\mu^{\mathbb{N}}(\tilde{Y}_a) > 0$ for almost every $a \in \pi_1(\tilde{Y})$ (where $\pi_1 : A \times G^{\mathbb{N}} \rightarrow A$ denotes the first projection as well). In other words, for almost every $x \in \tilde{Y}$, we have $\nu(Y_{\pi_1(x)}) > 0$.

Let us fix $x \in \tilde{Y}$ in the intersection of the two conull sets defined above. Let $a = \pi_1(x)$. The set Y_a is of positive measure, so its characteristic function χ_{Y_a} is an element of $L^\infty(B)$ which is not zero. Let h be its Poisson transform. Recall that it is defined as

$$h(g) = \int_B \chi_{Y_a} \, dg_* \nu = \nu(g^{-1} Y_a).$$

This function is a non-zero bounded harmonic function on G .

By definition of the Poisson transform, we have, for almost every $\omega = (\omega_1, \omega_2, \dots) \in \tilde{Y}_a$,

$$\begin{aligned} \nu((\omega_1 \omega_2 \dots \omega_n)^{-1} \tilde{Y}_a) &= h(\omega_1 \omega_2 \dots \omega_n) \\ &\xrightarrow[n \rightarrow +\infty]{} \chi_{Y_a}(\omega) = 1. \end{aligned}$$

In particular, we might pick n large enough so that $\nu((\omega_1 \omega_2 \dots \omega_n)^{-1} \tilde{Y}_a) > 1 - \varepsilon$ and also satisfying $T^n(a, \omega) \in \tilde{Y}$. Setting $g = (\omega_1 \omega_2 \dots \omega_n)^{-1}$, we have by definition $\nu(g Y_a) > 1 - \varepsilon$. Furthermore since $T^n(a, \omega) \in Y$, we have $(ga, \omega_{n+1}, \dots) \in Y$; hence $Y_{ga} \neq \emptyset$. \square

Proof of Theorem 4.9. Let \mathbf{X} be a metric field over (A, η) on which G acts via the cocycle α . Observe that we may assume that any fiber X_a has diameter at most 1. Otherwise, we replace d_a by $\max(d_a, 1)$ and we obtain a new metric field over B on which G acts as well.

Let φ be an invariant relative section. Let us define

$$f(a) = \int_{B \times B} d_a(\varphi(a, b), \varphi(a, b')) \, d\nu(b) \, d\nu(b').$$

Our assumption implies that f is not essentially 0. In particular, there is an $r > 0$ such that $A(r) := f^{-1}([r, +\infty))$ is of positive measure.

Take a small $\delta > 0$. Let $\{x^n\}$ be a fundamental family of \mathbf{X} . Then there is $n \in \mathbb{N}$ such that $Y = \{(a, b) \in Y_r \times B \mid d(\varphi(a, b), x_a^n) \leq \delta\}$ has positive measure, say $> \varepsilon$.

By Theorem 4.12, this implies that there is a $a \in \pi_1(Y)$ and a $g \in G$ such that $ga \in \pi_1(Y)$ and $\nu(g Y_a) > 1 - \varepsilon$.

Now for $(ga, b) \in g Y_a$ and $(ga, b') \in g Y_a$ (in other words, $b, b' \in Y_{ga}$), we know that $d(\varphi(ga, b), \alpha(g, a)x_a^n) \leq \delta$ and $d(\varphi(ga, b'), \alpha(g, a)x_a^n) \leq \delta$. So these two points, $\varphi(ga, b)$ and $\varphi(ga, b')$ are in the same ball of radius δ . Therefore we have $d(\varphi(ga, b), \varphi(ga, b')) \leq 2\delta$.

Let us decompose $B \times B$ as

$$(g Y_a \times g Y_a) \cup ((B \setminus g Y_a) \times g Y_a) \cup (g Y_a \times (B \setminus g Y_a)) \cup ((B \setminus g Y_a) \times (B \setminus g Y_a)).$$

We know that $\nu(B \setminus g Y_a) < \varepsilon$. Hence, $\nu((B \times B) \setminus (g Y_a \times g Y_a)) < \varepsilon^2 + 2\varepsilon$.

Therefore, we have

$$f(ga) < \int_{g Y_a \times g Y_a} d(\varphi(ga, b), \varphi(ga, b')) \, d\nu(b) \, d\nu(b') + 2\varepsilon + \varepsilon^2$$

It follows that $f(ga) < 2\delta + 2\varepsilon + \varepsilon^2$. Since we also assumed that $ga \in \pi_1(A)$, we have that $g \in A(r)$. Therefore, $f(ga) > r$. Since our choice of δ and ε is arbitrary, we get a contradiction. \square

Proof of Theorem 4.8. The amenability of $G \curvearrowright B$ is due to Zimmer [Zim78] and relative ergodicity for $\check{B} \times B \rightarrow \check{B}$ comes from Theorem 4.9. Relative ergodicity for $\check{B} \times B \rightarrow B$ is similar using that $\check{\mu} = \mu$. \square

5. FURSTENBERG MAPS

The geometric part of Theorem 1.1 uses the following version of Adams-Ballmann theorem [AB98, CL10] as a key tool.

Theorem 5.1 (Equivariant Adams-Ballmann theorem [Duc13, Theorem 1.8]). *Let (A, η) be an ergodic G -space such that $G \curvearrowright A$ is amenable. Let \mathbf{X} be a CAT(0) field of finite telescopic dimension.*

If G acts on \mathbf{X} then there is an invariant section of the boundary field $\partial\mathbf{X}$ or there exists an invariant Euclidean subfield of \mathbf{X} .

Before beginning the proof of Theorem 1.1 we can give a hint about its structure. Thanks to Theorem 5.1 used for a constant field, the only case where there is something to prove is the case where there is an invariant Euclidean subfield and no Furstenberg map. In that case, we choose a minimal such invariant Euclidean subfield. Then, we analyze the possible relative positions of two subflats of X by using Proposition 2.1. Using relative metric ergodicity we conclude that this subfield must actually be constant equal to some fixed Euclidean subspace.

Theorem 1.1 will actually be deduced straightforwardly from this more general theorem for boundary pairs.

Theorem 5.2. *Let X be a CAT(0) space of finite telescopic dimension and let G be a locally compact second countable group acting continuously by isometries on X without invariant flats. If (B_-, ν_-) and (B_+, ν_+) form a G -boundary pair then there exist measurable G -maps $\varphi_\pm: B_\pm \rightarrow \partial X$.*

Proof. First we do some reductions to prove the theorem. Clearly, it suffices to prove the theorem in case there is no fixed point of G in ∂X since otherwise we have a trivial map $B_\pm \rightarrow \partial X$. Let us assume there is no fixed point at infinity. In that case, there is a G -invariant closed convex space with a minimal G -action. That is there is no non-trivial G -invariant closed convex subset [CL10, Proposition 1.8 (ii)]. Thus we may assume that this space is actually X itself and X is the closed convex hull of any orbit. Orbits are separable since G acts continuously and G is second countable. Thus X is separable.

Let $X = H \times Y$ be the de Rham decomposition of X where H is a Hilbert space — of finite dimension since X has finite telescopic dimension — and Y is another CAT(0) space of finite telescopic dimension. The action of G on $X = H \times Y$ is diagonal and since the action of G on X is minimal, both actions $G \curvearrowright H$ and $G \curvearrowright Y$ are minimal. Observe it suffices to prove the result for $G \curvearrowright Y$ because $\partial Y \subseteq \partial X$ and if F is a flat of Y then $H \times F$ is a flat of X . We may assume that $X = Y$, that is X has no Euclidean de Rham factor of positive dimension.

Thus we reduce the proof of the following case: X is separable, has trivial Euclidean de Rham factor and the action of G is minimal.

We consider the constant fields \mathbf{X}^\pm over B_\pm with fibers X . These fields are endowed naturally with actions of G and we can apply Theorem 5.1. For each \mathbf{X}^\pm we get a map from B_\pm to ∂X or an invariant Euclidean subfield of \mathbf{X}^\pm . In the first case, one gets two G -maps $B_\pm \rightarrow \partial X$, we are done. Up to permuting B_+ and B_- , one may assume either that we get two invariant Euclidean subfields \mathbf{E}^- and \mathbf{E}^+ of respectively \mathbf{X}^- and \mathbf{X}^+ and there is no G -map $B_\pm \rightarrow \partial X$ (**Case (I)**) or we get a G -map $b \mapsto \xi_b$ from B_- to ∂X and a Euclidean subfield \mathbf{E} of \mathbf{X}^+ (**Case (II)**).

Readers interested only in the case of a G -boundary B , that is $B = B_- = B_+$, may read only case (I) with $\mathbf{X}^+ = \mathbf{X}^-$ and $\mathbf{E}^+ = \mathbf{E}^-$.

Case (I): Thanks to Lemma 3.7, the map $b \mapsto \dim(E_b^-)$ is measurable and G -invariant. Thanks to ergodicity of $G \curvearrowright B_-$, this dimension is essentially constant. We may assume that this dimension is minimal among dimensions of invariant Euclidean subfields of \mathbf{X}_- . For simplicity, we note $E_b = E_b^-$ and $E_{b'} = E_{b'}^+$, for $b \in B_-$ and $b' \in B_+$.

We aim to apply Proposition 2.1 to each pair $(E_b, E_{b'})$ for $(b, b') \in B_- \times B_+$. First, we show that the four conditions in the proposition are measurable and G -invariant; thanks to ergodicity of $G \curvearrowright B_- \times B_+$, exactly one condition will be satisfied for almost every $(b, b') \in B_- \times B_+$.

Let $\{x_b^n\}_{b \in B_-}$ be a fundamental family of \mathbf{E}^- , and $\{x_{b'}^n\}_{b' \in B_+}$ be a fundamental family of \mathbf{E}^+ . We define $d(b, b') = \inf_{n,m} d(x_b^n, x_{b'}^m)$. The function $d : B_- \times B_+ \rightarrow \mathbb{R}$ is measurable. The infimum distance between E_b and $E_{b'}$ is not achieved if and only if for any $N \in \mathbb{N}$ and any n such that $d(x_b^0, x_{b'}^n) < N$, one has $\inf_m d(x_b^n, x_{b'}^m) > d(b, b')$. This condition is measurable and G -invariant.

If the minimal distance between E_b and $E_{b'}$ is achieved, then the subset of E_b where the distance is achieved is not bounded if and only if for any $N \in \mathbb{N}$, there exists n_1, n_2 such that $d(x_b^{n_1}, x_b^{n_2}) > N$ and $\inf_m d(x_b^{n_i}, x_{b'}^m) \leq 1 + d(b, b')$.

Now, we know that exactly one of these four relative positions given in Proposition 2.1 between E_b and $E_{b'}$ happens for almost every (b, b') . We treat the four cases independently.

Case (i): If the infimum distance is not achieved, choose a section y of \mathbf{E}^- and consider (for b' in a set of full measure) the nonincreasing sequence of subfields \mathbf{X}^n where we define X_b^n as $X_b^n = \left\{ x \in E_b, d(x, E_{b'}) \leq d(b, b') + \frac{1}{n} \right\}$. We can now apply [Duc13, Proposition 8.10] and obtain a section $\xi_b(b')$ of the boundary field $\partial\mathbf{E}^-$. This field is a metric field with a G -action by Lemma 3.11. By construction, ξ satisfies $\alpha(g, b)\xi_b(b') = \xi_{gb}(gb')$. This means that $\xi : B_- \times B_+ \rightarrow \sqcup\partial E_b$ is an invariant section relative to the first projection $B_- \times B_+ \rightarrow B_-$. Now, thanks to relative metric ergodicity, ξ does not depends on b' . So we can interpret $b \mapsto \xi_b$ as a measurable G -map from B_- to ∂X . This yields a contradiction.

If the minimal distance is achieved, we define $Y_b(b') = \{x \in X_b, d(x, E_{b'}) = d(b, b')\}$. Then the family $\mathbf{Y} = (Y_b(b'))_b$ is a subfield of \mathbf{X} (which depends on b').

Case (ii): Assume the minimal set $Y_b(b')$ is bounded. Let $x_b(b')$ be its circumcenter. Then for almost all b' , the family of circumcenters $\{x_b(b')\}$ is a section of \mathbf{E}^- [Duc13, Lemma 8.7] which satisfies $\alpha(g, b)x_b(b') = x_{gb}(gb')$, that is an invariant relative section. Relative metric ergodicity shows that this map is essentially constant. So its essential image is a point which is fixed by G , which is excluded.

In the other case, thanks to [Duc13, Proposition 9.2], we can write $Y_b(b') = F_b(b') \times T_b(b')$ where $\mathbf{F} = (F_b(b'))_b$ is a maximal Euclidean subfield of \mathbf{Y} and $(T_b(b'))_b$ is a subfield of \mathbf{Y} ; both subfields depending on b' .

Before attacking Case (iii), which is harder, let us treat Case (iv).

Case (iv): If for almost (b, b') , $T_b(b')$ is not bounded then its boundary is not empty and its circumradius $< \pi/2$ thanks to Proposition 2.1. Let $\xi_b(b')$ be the circumradius of $\partial T_b(b')$. Thanks to Lemma 3.4, we get a measurable G -map from $B_- \times B_+$ to the metric field $\partial\mathbf{E}^-$ over B_- . Once again relative metric ergodicity implies that $\xi_b(b')$ coincides with a G -map $b \mapsto \xi_b$ and we are done.

Case (iii): If $T_b(b')$ is bounded for almost every (b, b') then we set $t_b(b')$ to be its circumcenter and $E'_b(b') = F_b(b') \times \{t_b(b')\}$.

The dimension of $E'_b(b')$ is measurable (Lemma 3.7) in (b, b') and G -invariant. Thus, this dimension is essentially constant equal to some $k \in \mathbb{N}$.

Now $(b, b') \mapsto \partial E'_b(b')$ is a G -section of the metric field \mathbf{S} of Euclidean subspheres of dimension $k - 1$ associated to \mathbf{E}^- (see Lemma 3.13). Relative metric ergodicity implies that $\partial E'_b(b')$ is essentially equal to some s_b and $s = (s_b)$ is a G -invariant section of \mathbf{S} . Using the second part of Lemma 3.13, let us consider the field \mathbf{E}^s such that for any $b \in B$, E_b^s is the set of Euclidean subspaces of E_b with boundary s_b . By definition $(b, b') \mapsto E'_b(b')$ is a relative invariant section of \mathbf{E}^s and relative metric ergodicity implies that there is a section of \mathbf{E}^s , that is a Euclidean subfield \mathbf{E}' of \mathbf{E}^- such that for almost every (b, b') , $E'_b(b') = E'_b$.

Our assumption on the minimality of the dimension of \mathbf{E}^- implies that $E'_b = E_b$ for almost every $b \in B_-$.

Going back to the definition of $E'_b = E_b(b')$, we see that the fact that $E'_b = E_b$ implies that the distance function $d(\cdot, E_{b'})$ must be constant in restriction to E_b . That is to say that E_b and $E_{b'}$ are parallel for almost all $b, b' \in B_- \times B_+$.

Fubini's theorem tells us that there is $b \in B_-$ such that for a set $B' \subseteq B_+$ of full measure and for any $b' \in B'$, $E_{b'}$ is parallel to E_b . Fix $\Gamma \leq G$ a dense countable subgroup of G and set $B_0 = \cap_{\gamma \in \Gamma} \gamma B'$ which is a Γ -invariant subset of full measure in B_+ . Lemma 2.3 implies that the closed convex hull of the union of the $\{E_b\}_{b \in B_0}$ splits as some product $E \times T$ where each E_b is parallel to E . Observe that Γ preserves this convex set. By continuity of the action, the group G preserves this set as well and by minimality this set is X . Now the assumption that X has trivial Euclidean de Rham factor implies that E is a point and thus the dimension of E_b for almost every b is zero. Finally $E_b, E_{b'}$ are points of X and double metric ergodicity (Remark 4.11) implies there is a fixed point.

Case (II): Fix a point $x \in X$ and for $b \in B_-$ let β_b be the Busemann function associated to ξ_b vanishing at x . Now for $(b, b') \in B_- \times B_+$, look at the restriction $f_{b,b'}$ of β_b to $E_{b'}$. It is a convex function and arguing as above, if it is not constant, one gets a relative section of \mathbf{E} or of $\partial\mathbf{E}$. Once again relative ergodicity gives an invariant Euclidean subspace of X or a fixed point at infinity.

If $f_{b,b'}$ is constant, the situation is different. Actually since this situation is a measurable G -invariant condition on (b, b') thanks to double ergodicity it holds for almost all $(b, b') \in B_- \times B_+$. Applying Lemma 3.6 to \mathbf{E} , we get that the subfield \mathbf{Y} of flats parallel to \mathbf{E} splits as $\mathbf{E} \times \mathbf{Z}$ and G acts on \mathbf{Z} with an invariant section z . Lemma 2.4 shows that ξ is actually a section of $\partial\mathbf{Z}$.

Let \mathbf{B}^r be the subfield of \mathbf{Z} consisting of closed balls of radius $r > 0$ around z . Thanks to Lemma 3.5, G acts on this CAT(0) field. Let $z_{b'}^r(b)$ be the point on the ray from $z_{b'}$ to ξ_b at distance r from $z_{b'}$. This is an invariant relative section of \mathbf{B}^r and thanks to relative ergodicity, there is an invariant section $z_{b'}^r$ of \mathbf{B}^r such that $z_{b'}^r(b) = z_{b'}^r$ for almost $(b, b') \in B_- \times B_+$.

Let us define $\eta_{b'}$ to be $\lim_{r \rightarrow \infty} z_{b'}^r$. By construction this is a invariant section of $\partial\mathbf{X}_+$ that is a G -map $\varphi_+ : B_+ \rightarrow \partial X$ as desired. \square

APPENDIX A. NON-EMPTYNESS OF THE BOUNDARY OF MINIMAL CAT(0) SPACES OF FINITE TELESCOPIC DIMENSION

P. Py pointed out that the proof of Proposition 1.8 in [CL10] uses implicitly the fact that the boundary of an unbounded CAT(0) space of finite telescopic dimension is non empty as soon as its group of isometries acts minimally (meaning that is there is no non-trivial closed convex subset which is invariant under all isometries). The proof of [CL10, Proposition 1.8] was completed in [CL] but the question about the non-emptiness of the boundary still remained.

We answer positively this question. The example of a rooted tree with an edge of length n attached to the root for any $n \in \mathbb{N}$ shows that the hypothesis of no fixed point for all isometries is essential. Observe that if the boundary is empty then any isometry is elliptic since any hyperbolic or parabolic isometry has a fixed point at infinity. Moreover, [CL10, Theorem 1.6] shows that the isometry group of such a minimal space cannot be amenable.

We emphasize that Theorem 1.2 relies on Theorem 5.1 which only uses [CL10, Proposition 1.8(ii)] where there is no implicit assumption. So, the gap in the cited paper is filled in.

Proof of Theorem 1.2. First of all, we prove the result for finitely generated groups. Let G be a finitely generated group acting on some CAT(0) space of finite telescopic dimension X with empty boundary. This last assumption implies that there is a closed convex subset on which G acts minimally. We may assume this subset is X itself. Since G is countable then the closed convex hull of any orbit is G -invariant and separable. The minimality assumption implies X coincides with any such closed convex hull and thus X is separable. Since G is finitely generated then G has a symmetric probability measure μ uniformly supported on a symmetric finite set of generators and thus the Poisson boundary B associated to μ is a G -boundary.

We can apply Theorem 5.1 for the constant field of CAT(0) spaces X over B and we obtain a measurable map $B \rightarrow \partial X$ or an invariant subfield of flats. In any case, we get an non empty boundary or an equivariant map $B \rightarrow X$. In this last case double metric ergodicity for $G \curvearrowright B$




FIGURE 1. An unbounded CAT(0) space of dimension 1 with empty boundary

implies that this map is essentially constant and we get a G -fixed point.

We conclude in the general case in the following way: assume X has empty boundary and let G be the group of all isometries of X . Since $\partial X = \emptyset$, X is \mathcal{T}_c -compact (see §2.1). Let \mathcal{F} be the collection of all finite subsets of G . For $F \in \mathcal{F}$ let G_F be the subgroup generated by F . We just proved that G_F has a non-empty, closed and convex set of fixed points that we denote by X_F . The collection $\{X_F\}_{F \in \mathcal{F}}$ is then a filtering family of \mathcal{T}_c -closed non-empty subspaces, by compactness $\cap_{F \in \mathcal{F}} X_F \neq \emptyset$ and thus one gets a G -fixed point. \square

Proof of Corollary 1.3. The proof goes as in [CM09, Proposition 1.5]. We reproduce it for reader's convenience. Theorem 1.2 shows that ∂X is not empty. Assume ∂X has radius at most $\pi/2$ then there is a canonical point $\xi \in \partial X$ fixed by all isometries and such that the closed ball of radius $\pi/2$ — for the Tits distance — around ξ coincides with ∂X . If $g \in \text{Isom}(X)$ does not let the Busemann function β_ξ invariant, then g is an hyperbolic isometry and translates along some geodesic line with extremities $\eta_-, \eta_+ \in \partial X$. One of this point does not belong to the closed ball of radius $\pi/2$ around ξ yielding a contradiction.

Now, if ∂X has radius larger than $\pi/2$, there is a minimal closed convex subset $Y \subseteq X$ with $\partial Y = \partial X$. The union X_0 of all such minimal spaces is closed convex and can be decomposed as $X_0 = Y \times Z$. Moreover X_0 is $\text{Isom}(X)$ -invariant with a diagonal action. Minimality implies that $X = X_0$. If $\partial Z \neq \emptyset$ one has $\partial Y = \partial X = \partial Y * \partial Z$ leading to a contradiction. Thus $\partial Z = \emptyset$ and Theorem 1.2 implies $\text{Isom}(X) \curvearrowright Z$ has a fixed point. This fixed point is the whole of Z by minimality and thus $X = Y$. \square

REFERENCES

- [AB94] Norbert A'Campo and Marc Burger. Réseaux arithmétiques et commensurateur d'après G. A. Margulis. *Invent. Math.*, 116(1-3):1–25, 1994. [1](#)
- [AB98] Scot Adams and Werner Ballmann. Amenable isometry groups of Hadamard spaces. *Math. Ann.*, 312(1):183–195, 1998. [12](#)
- [AH14] Martin Anderegg and Philippe Henry. Actions of amenable equivalence relations on CAT(0) fields. *Ergodic Theory Dynam. Systems*, 34(1):21–54, 2014. [6](#)
- [BF] Uri Bader and Alex Furman. Boundary pairs, algebraic gates and their applications. In preparation. [9, 10](#)
- [BF12] Uri Bader and Alex Furman. Boundaries, Weyl groups, and superrigidity. *Electron. Res. Announc. Math. Sci.*, 19:41–48, 2012. [9](#)
- [BF13] U. Bader and A. Furman. Algebraic Representations of Ergodic Actions and Super-Rigidity. *ArXiv e-prints*, November 2013, 1311.3696. [1, 9](#)

- [BH99] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999. [1](#), [3](#), [4](#), [5](#), [6](#), [7](#)
- [BM02] M. Burger and N. Monod. Continuous bounded cohomology and applications to rigidity theory. *Geom. Funct. Anal.*, 12(2):219–280, 2002. [1](#)
- [Bou04] Nicolas Bourbaki. *Integration. I. Chapters 1–6*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004. Translated from the 1959, 1965 and 1967 French originals by Sterling K. Berberian. [9](#)
- [CL] Pierre-Emmanuel Caprace and Alexander Lytchak. Erratum to ‘At Infinity of finite-dimensional CAT(0) spaces’. Unpublished. [14](#)
- [CL10] Pierre-Emmanuel Caprace and Alexander Lytchak. At infinity of finite-dimensional CAT(0) spaces. *Math. Ann.*, 346(1):1–21, 2010. [1](#), [2](#), [3](#), [12](#), [14](#)
- [CM09] Pierre-Emmanuel Caprace and Nicolas Monod. Isometry groups of non-positively curved spaces: structure theory. *J. Topol.*, 2(4):661–700, 2009. [2](#), [15](#)
- [Duc12] B. Duchesne. Infinite dimensional Riemannian symmetric spaces with fixed-sign curvature operator. *ArXiv e-prints*, April 2012, 1204.6012. [1](#)
- [Duc13] Bruno Duchesne. Infinite-dimensional nonpositively curved symmetric spaces of finite rank. *Int. Math. Res. Not. IMRN*, (7):1578–1627, 2013. [6](#), [7](#), [12](#), [13](#)
- [Fre81] D. H. Fremlin. Measurable functions and almost continuous functions. *Manuscripta Math.*, 33(3-4):387–405, 1980/81. [9](#)
- [Fur63] Harry Furstenberg. A Poisson formula for semi-simple Lie groups. *Ann. of Math. (2)*, 77:335–386, 1963. [1](#), [10](#)
- [Fur73] Harry Furstenberg. Boundary theory and stochastic processes on homogeneous spaces. In *Harmonic analysis on homogeneous spaces (Proc. Sympos. Pure Math., Vol. XXVI, Williams Coll., Williamstown, Mass., 1972)*, pages 193–229. Amer. Math. Soc., Providence, R.I., 1973. [1](#)
- [Jaw94] Wojciech Jaworski. Strongly approximately transitive group actions, the Choquet-Deny theorem, and polynomial growth. *Pacific J. Math.*, 165(1):115–129, 1994. [10](#)
- [Jun01] H.W.E. Jung. Über die kleinste kugel, die eine räumliche figur einschliesst. *Journal für die Reine und Angewandte Mathematik*, 123:241–257, 1901. [2](#)
- [Kai00] Vadim A. Kaimanovich. The Poisson formula for groups with hyperbolic properties. *Ann. of Math. (2)*, 152(3):659–692, 2000. [2](#)
- [Kai03] V. A. Kaimanovich. Double ergodicity of the Poisson boundary and applications to bounded cohomology. *Geom. Funct. Anal.*, 13(4):852–861, 2003. [1](#), [10](#)
- [Kec95] Alexander S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. [5](#)
- [Kle99] Bruce Kleiner. The local structure of length spaces with curvature bounded above. *Math. Z.*, 231(3):409–456, 1999. [1](#), [3](#)
- [KM99] Anders Karlsson and Gregory A. Margulis. A multiplicative ergodic theorem and nonpositively curved spaces. *Comm. Math. Phys.*, 208(1):107–123, 1999. [2](#)
- [LS97] Urs Lang and Viktor Schroeder. Jung’s theorem for Alexandrov spaces of curvature bounded above. *Ann. Global Anal. Geom.*, 15(3):263–275, 1997. [6](#)
- [Mar91] G. A. Margulis. *Discrete subgroups of semisimple Lie groups*, volume 17 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1991. [1](#)
- [Mon06] Nicolas Monod. Superrigidity for irreducible lattices and geometric splitting. *J. Amer. Math. Soc.*, 19(4):781–814, 2006. [3](#)
- [Val77] M. Valadier. Sur le plongement d’un champ mesurable d’espaces métriques dans un champ trivial. *Travaux Sém. Anal. Convexe*, 7(3, Exp. No. 14):4, 1977. [5](#)
- [Zim78] Robert J. Zimmer. Amenable ergodic group actions and an application to Poisson boundaries of random walks. *J. Functional Analysis*, 27(3):350–372, 1978. [11](#)
- [Zim84] Robert J. Zimmer. *Ergodic theory and semisimple groups*, volume 81 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1984. [9](#)