

Lecture Notes¹ for the course
Distributions and Partial Differential Equations

Master Mathématiques Fondamentales et Applications
Université Paris-Saclay, Sep-Dec 2020

Stéphane Nonnenmacher

September 2020

1. following earlier versions by Thierry Ramond and Patrick Gérard

Contents

1 Distributions in one space dimension	8
1.1 Background on differential calculus	8
1.1.1 Basic properties of smooth functions on I	8
1.1.2 Support of a continuous function	9
1.2 Test functions	9
1.2.1 The space of test functions	9
1.2.2 Smooth partitions of unity	12
1.2.3 Convergence in the space of test functions	15
1.3 Definition of distributions on I	16
1.3.1 Definitions	16
1.3.2 Distributions defined by locally integrable functions	18
1.3.3 The Dirac mass	20
1.3.4 A distribution involving an infinite number of derivatives	21
1.3.5 Cauchy's principal value of $1/x$	21
1.4 Differentiating distributions	22
1.5 Product by a smooth function	27
1.6 Restriction and support	31
1.6.1 Definitions	31
1.6.2 Some properties of the support	34
1.7 Sequences of distributions	36
1.7.1 Convergence in \mathcal{D}'	36
1.7.2 The Uniform Boundedness Principle	38
1.8 An introduction to Sobolev spaces	40
1.8.1 Generalised derivatives.	40

1.8.2	The Sobolev space $H^1(I)$.	40
1.8.3	The special case of a bounded interval	41
1.9	Further properties	45
1.9.1	Characterisation of Lipschitz functions	45
1.9.2	Derivation and integration under the bracket	46
1.9.3	Convolution and regularisation	48
1.9.4	Positive distributions and increasing functions	50
1.9.5	The structure of distributions	53
2	Distributions in several variables	56
2.1	A brief review of several variables differential calculus	56
2.1.1	Scalar product, norm, distance, topology	56
2.1.2	Partial derivative, C^1 functions, differential, gradient	57
2.1.3	The Chain rule	58
2.1.4	Higher order partial derivatives	58
2.1.5	Multi-indices	59
2.2	Test functions	61
2.2.1	Definitions. Examples	61
2.2.2	Partitions of unity	62
2.3	Distributions on an open subset of \mathbb{R}^d	63
2.3.1	Définitions. Examples	63
2.3.2	Restriction and support	64
2.3.3	Multiplication by a smooth function	67
2.3.4	Derivation	68
2.3.5	Convergence	69
2.4	Superficial measures and the jump formula	75
2.4.1	Motivation	75
2.4.2	Preliminaries on non-negative distributions	75
2.4.3	The measure $\delta(f)$ and the superficial measure associated to $\{f = 0\}$.	76
2.4.4	Integration on level sets : the smooth coarea formula	79
2.4.5	Superficial measure on a closed hypersurface and the jump formula for a regular open subset	80
2.5	Sobolev spaces in an open set	83

2.5.1	Definition and general facts	84
2.5.2	Variational formulation of some elliptic problems	84
2.5.3	Approximation by smooth functions	87
2.5.4	The trace theorem	91
2.5.5	Solving the Dirichlet problem	92
2.6	The uniform boundedness principle	93
2.6.1	Step1. Realizing \mathcal{D}_K as a complete metric space	93
2.6.2	Step 2. The Baire lemma	95
2.6.3	Step 3. Proof of Theorem 2.6.1	96
3	The Fourier Transformation	97
3.1	The Schwartz space	97
3.1.1	Definitions and examples	97
3.1.2	Convergence in $\mathcal{S}(\mathbb{R}^d)$ and density results	98
3.2	The Fourier transformation in $\mathcal{S}(\mathbb{R}^d)$	100
3.2.1	Definition and first properties	100
3.2.2	Gaussians	101
3.2.3	The Inversion formula	102
3.2.4	The "change of head" lemma	103
3.2.5	The Plancherel identity	103
3.3	The space $\mathcal{S}'(\mathbb{R}^d)$ of tempered distributions	104
3.3.1	Definition, examples	104
3.3.2	The fundamental characterization of \mathcal{S}'	106
3.3.3	Convergence in $\mathcal{S}'(\mathbb{R}^d)$	107
3.4	The Fourier Transformation in $\mathcal{S}'(\mathbb{R}^d)$	108
3.4.1	Definition	108
3.4.2	The Fourier Transformation on L^1 and L^2	109
3.4.3	The Fourier Transform of compactly supported distributions	110
3.4.4	Fourier characterization of Sobolev spaces on \mathbb{R}^d	111
3.5	Some applications	114
3.5.1	Partial differential equations with constant coefficients	114
3.5.2	Local results	115

3.5.3 Global results	118
3.6 Fourier series	124
3.7 An application to probability theory	131

Introduction : Why Distributions ?

The notion of *distribution* emerged during the twentieth century, as a powerful tool in the study of partial differential equations (PDEs). The need to generalize the concept of a "function" also emerged from different parts of theoretical physics, in particular electromagnetism and quantum mechanics. On the mathematical side, major contributors to the formal definition of distributions are Jean Leray, Serguei Sobolev and Laurent Schwartz.

In this introduction, we present four simple motivations for the emergence of distributions.

Deriving non differentiable functions

Taking the derivative of non-differentiable functions

In differential calculus, one immediately encounters the unpleasant fact that not every function is differentiable. The purpose of distribution theory is to remedy this flaw. Recall that, given a function f on an open interval $I \subset \mathbb{R}$, the derivative function f' is defined as

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h},$$

provided this limit exists at every point $x \in I$. Let us mention a classical elementary application of this notion. If f' exists, f is a nondecreasing function on I if and only if¹ its derivative $f'(x) \geq 0$ at every point $x \in I$. However, there are many examples of nondecreasing functions such that f' cannot be defined on the whole of I : a typical example is the Heaviside function,

$$H(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases}$$

Of course, H admits a derivative at every point $x \neq 0$, and this derivative is 0. But this is not sufficient to conclude that H is nondecreasing, since the derivative of $-H$ enjoys the same property ! It is therefore necessary to extend to $x = 0$ the derivative of H , in a way which takes into account the discontinuity of H at $x = 0$. In fact, to every locally integrable function f , we shall associate a mathematical object — a distribution — called the derivative of f , with the property that f is nondecreasing iff its derivative is nonnegative (the nonnegativity of a distribution will need to be properly defined).

1. in the sequel, the expression "if and only if" will be abbreviated by *iff*

Fourier series

In his famous memoir *Théorie analytique de la chaleur* (1822), Joseph Fourier introduced, for every “reasonable” 2π -periodic function f , the coefficients

$$c_n(f) = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-inx} dx, \quad n \in \mathbb{Z}$$

and he stated that

$$f(x) = \sum_{n=-\infty}^{+\infty} c_n(f) e^{inx}.$$

Throughout the nineteenth and the early twentieth century, many mathematicians have tried to give sense to the above equality for the largest possible class of functions f . Dirichlet has been proved it for any C^1 function, yet Kolmogorov made the striking observation that there exists locally integrable functions f such that the above Fourier series is divergent for every $x \in \mathbb{R}$! We shall see below that, even in such an unfavorable situation, the above series is convergent, but in a different sense, namely *in the sense of distributions*. In fact, for any element u in the class of 2π -periodic distribution (this class includes, in particular, the locally integrable functions), we shall define a sequence $(c_n(u))_{n \in \mathbb{Z}}$ of Fourier coefficients such that the corresponding series is convergent in the sense of distributions, and its sum equals u . Furthermore, the derivative of the distribution u (which is also a 2π -periodic distribution) can be obtained by summing the series of derivatives, namely the series with coefficients $(inc_n(u))$. This interplay between Fourier series and differentiation was a major reason for their introduction by Fourier back in 1822; the theory of distributions manages to extend the applicability of this connection to a much larger class of objects.

Electrostatics

If f is a function on \mathbb{R}^3 which represents an electric charge distribution, the electric potential u generated by this charge distribution satisfies the Poisson equation

$$-\Delta u = f.$$

If f is smooth enough — say C^1 — and small enough at infinity (for instance it vanishes outside of some ball), one can prove that there exists a unique C^2 solution u of this equation which goes to 0 at infinity: this solution is given by the integral

$$(0.0.1) \quad u(x) = \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{f(y)}{|x-y|} dy.$$

This formula still has a meaning if f is bounded and has compact support (we will note $f \in L_{comp}^\infty(\mathbb{R}^3)$), but is not necessarily continuous. In that case u is no more C^2 in general, yet it is tempting to try to interpret the Poisson equation in this more general situation as well.

Actually, in Physics charge distributions are often modeled as being supported on surfaces, curves, points, or forming microscopic dipoles... Such charge distributions cannot be described by L_{comp}^∞ functions. Yet, in all these cases, the formula (0.0.1) still makes sense, provided one integrates on the appropriate subset of \mathbb{R}^3 ; what is then the status of the Poisson equation ? We will show that it is possible to interpret the Poisson equation in all these cases, *in the sense of distributions*.

Let us remark that the word *distributions* actually takes its origin in this application to electrostatics, a (mathematical) distribution primarily represented a distribution of electric charge.

Quantum mechanics

Quantum mechanics is a theory based on the Hilbert space $L^2(\mathbb{R}^3, \mathbb{C})$: the state of a quantum particle² is represented by a wavefunction, which is a square integrable function $u \in L^2(\mathbb{R}^3, \mathbb{C})$, with L^2 norm equal to unity. Nevertheless, the founding fathers of the theory (in particular Paul Dirac) felt the need to extend the notion of quantum state to objects which are not in L^2 . In particular, Dirac introduced his famous “delta function”, namely a “function” $\delta(x)$ which vanishes at each point $x \neq 0$, but such that, *formally*

$$\int_{\mathbb{R}^3} \delta(x) dx = 1.$$

If one had to give a value to $\delta(0)$, this value should be $+\infty$ for the above identity to hold. So this δ cannot be a function: it is not an element of L^2 . Mathematically, this “delta function” will appear as one of the simplest nontrivial distributions, the delta distribution at the origin, often denoted by δ_0 .

Dirac also invented the “bracket” notation $\langle u|v \rangle$ for the scalar bracket on L^2 , and extended the notation to his singular objects: his notation $\langle x_0|v \rangle$ will represent the distributional bracket between the delta distribution δ_{x_0} and the “test function” v , which we will denote by similar brackets $\langle \delta_{x_0}, u \rangle_{\mathcal{D}', \mathcal{D}}$.

2. Here I only consider *scalar* particles, that is particles without spin.

Chapter 1

Distributions in one space dimension

We have chosen to start our presentation of the mathematical theory of distributions by the one dimensional setting. This will give simpler expressions, yet will already convey most of the ideas and methods of the theory. Note that this one dimensional theory is already useful in practice, for instance in signal analysis, or in 1-dimensional models in quantum or wave mechanics.

Distributions will appear below as “duals” of smooth functions (more precisely, linear forms acting on spaces of smooth functions), a distribution will be described by the way it acts on all smooth functions. For this reason, these smooth functions will be called *test functions*, since their rôle will be to test the distributions.

We will start our by describing in some detail these classes of test functions.

1.1 Background on differential calculus

In this section, we recall elementary facts about smooth functions of one real variable. The functions will be defined on some open (nonempty) interval $I \subset \mathbb{R}$. This interval may be bounded ($I =]a, b[$), semibounded ($I =]-\infty, b[$ or $I =]a, \infty[$), or simply $I = \mathbb{R}$. The theory will essentially be identical in all cases. For the reader's ease, at first read it can be convenient to take $I = \mathbb{R}$.

1.1.1 Basic properties of smooth functions on I

A smooth function on an open interval $I \subset \mathbb{R}$ is a function $\varphi : I \rightarrow \mathbb{C}$ whose derivatives of any order $\varphi', \varphi'', \dots, \varphi^{(k)}, \dots$ exist and are continuous on I . A linear combination of smooth functions is a smooth function, and we denote $\mathcal{C}^\infty(I)$ the vector space formed by all smooth functions on I .

If $\varphi \in \mathcal{C}^\infty(I)$ and J is an open subset of I , the function defined on J by $x \mapsto \varphi(x)$ is a smooth function on J , that we denote by $\varphi|_J$. This function is called the restriction of φ on J .

The product of two smooth functions is smooth, and the derivatives of the product are expressed by *Leibniz formula*:

$$(\varphi_1 \varphi_2)^{(k)} = \sum_{j=0}^k \binom{k}{j} \varphi_1^{(j)} \varphi_2^{(k-j)}.$$

A smooth function $\varphi : I \rightarrow \mathbb{C}$ satisfies the Taylor formula with integral remainder, for any order m and any

base point $x_0 \in I$:

$$\forall x \in I, \varphi(x) = \sum_{k=0}^m \frac{(x-x_0)^k}{k!} \varphi^{(k)}(x_0) + \frac{(x-x_0)^{m+1}}{m!} \int_0^1 (1-s)^m \varphi^{(m+1)}(x_0 + s(x-x_0)) ds.$$

This formula can be proved by integrating by parts the last term on the right hand side.

Exercise 1.1.1 Prove Hadamard's lemma: if $\varphi \in \mathcal{C}^\infty(I)$ satisfies $\varphi(x_0) = 0$, there exists a function $\psi \in \mathcal{C}^\infty(I)$ such that $\varphi(x) = (x-x_0)\psi(x)$ for any $x \in I$.

In other words, we can factorize from φ the monomial $(x-x_0)$, and the quotient is still a smooth function.

1.1.2 Support of a continuous function

If f is continuous on I and J is an open subset of I , we say that f vanishes on J if it vanishes at every point of J , or, equivalently, if $f|_J$ is the null function.

Definition 1.1.2 Let $f : I \rightarrow \mathbb{C}$ be a continuous function. The support of f is the complement of the union of all the open sets in I where f vanishes. This set is denoted by $\text{supp } f$.

Note that the support of f is a closed set. It is also the closure of the set of $x \in I$ such that $f(x) \neq 0$. The following characterization is often useful:

$$x_0 \notin \text{supp } f \iff \exists V \text{ neighborhood of } x_0 \text{ such that } f|_V = 0.$$

Exercise 1.1.3 Show that

$$\text{supp}(f_1 f_2) \subset \text{supp } f_1 \cap \text{supp } f_2.$$

Are these two sets equal?

Of course, if $\varphi \in \mathcal{C}^\infty(I)$ vanishes on an open set $J \subset I$, all its derivatives vanish as well on J , and, therefore, for any integer k ,

$$\text{supp } \varphi^{(k)} \subset \text{supp } \varphi.$$

1.2 Test functions

In this section we introduce and manipulate a first class of test functions.

1.2.1 The space of test functions

We start by choosing I an open (nonempty) interval of \mathbb{R} .

Definition 1.2.1 We denote by $\mathcal{D}(I) = \mathcal{C}_0^\infty(I)$ the vector space of functions which are \mathcal{C}^∞ on I , and whose support is a compact subset of I . Equivalently, a smooth function belongs to $\mathcal{D}(I)$ if it vanishes outside some closed segment $[a, b] \subset I$.

If J is an open subinterval of I , one may identify a function $\varphi \in \mathcal{D}(J)$ with its extension $\tilde{\varphi} \in \mathcal{D}(I)$, that is the function defined as

$$\tilde{\varphi}(x) = \varphi(x) \text{ for } x \in J, \quad \tilde{\varphi}(x) = 0 \text{ for } x \in I \setminus J.$$

Indeed, $\tilde{\varphi}$ is automatically smooth and with compact support on I for $\varphi \in \mathcal{D}(J)$. On the other hand, a function $\varphi \in \mathcal{C}_0^\infty(\mathbb{R})$ can be identified with its restriction $\varphi|_I \in \mathcal{D}(I)$ for any open interval I that contains $\text{supp } \varphi$.




Figure 1.1: Schematic plot of a test function on \mathbb{R} .

Exercise 1.2.2 Let f and φ be two functions in $L^1(\mathbb{R})$. Show that the convolution $f * \varphi$ of f and φ , given by

$$f * \varphi(x) = \int f(x - y)\varphi(y)dy,$$

is defined almost everywhere, and is an L^1 function.

Suppose moreover that $\varphi \in \mathcal{C}_0^\infty(\mathbb{R})$, show that $f * \varphi$ is smooth. At last, if f is also continuous, show that

$$\text{supp } f * \varphi \subset \text{supp } f + \text{supp } \varphi.$$

Proofs: (cf TD1, Ex. 1). First of all, $f * \varphi$ is defined almost everywhere, and is integrable. Indeed, using Fubini's theorem for non-negative functions,

$$\begin{aligned} \int |f * \varphi(x)|dx &\leq \iint |f(x - y)\varphi(y)|dydx \\ &\leq \int |\varphi(y)| \left(\int |f(x - y)|dx \right) dy \leq \|\varphi\|_{L^1} \|f\|_{L^1} < +\infty \end{aligned}$$

Therefore $f * \varphi$ is finite almost everywhere, and belongs to L^1 .

(cf TD 1, Ex. 5). Suppose now that φ is smooth with compact support. Changing variables, we have

$$f * \varphi(x) = \int f(y)\varphi(x-y)dy.$$

Thus $f * \varphi$ is also smooth thanks to the Lebesgue theorem, since

- the function $x \mapsto f(y)\varphi(x-y)$ is smooth for all y , and

$$\partial_x^k(f(y)\varphi(x-y)) = f(y)\varphi^{(k)}(x-y),$$

- we have the domination

$$|f(y)\varphi^{(k)}(x-y)| \leq |f(y)| \sup |\varphi^{(k)}| \in L^1.$$

(cf TD1, Ex. 4) Finally, assume that f is continuous. If $x \notin \text{supp } f + \text{supp } \varphi$, then for any $y \in \text{supp } f$, $x-y$ does not belong to $\text{supp } \varphi$, thus

$$f * \varphi(x) = \int f(y)\varphi(x-y)dy = 0. \square$$

It is not immediately clear that $\mathcal{D}(I)$ is not reduced to the null function. One knows for example that the only compactly supported *real analytic function* on \mathbb{R} is the null function, due to the principle of isolated zeroes. Yet, one has the following

Proposition 1.2.3 The space $\mathcal{D}(I)$ is nontrivial. More precisely, for every $x_0 \in I$ and $r > 0$ such that $[x_0 - r, x_0 + r] \subset I$, there exists a function $\varphi_{x_0, r} \in \mathcal{C}^\infty(I)$ such that $\text{supp } \varphi_{x_0, r} = [x_0 - r, x_0 + r]$.

Proof.— Here we give a “standard” construction, which uses the the function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$\varphi(t) = \begin{cases} e^{-1/t} & \text{as } t > 0, \\ 0 & \text{as } t \leq 0. \end{cases}$$

Let us check that φ is smooth on \mathbb{R} . Indeed, it is easily seen to be \mathcal{C}^∞ on \mathbb{R}^* , with $\varphi^{(k)}(t) = 0$ for all $t < 0$. On the other hand, for $t > 0$, one can prove by induction that

$$\forall k \in \mathbb{N}, \varphi^{(k)}(t) = P_k \left(\frac{1}{t} \right) e^{-1/t}$$

where P_k is a polynomial of degree $2k$. Therefore, for any $k \geq 0$, $\varphi^{(k)}(t) \rightarrow 0$ as $t \rightarrow 0^+$, so that $\varphi^{(k)}$ is continuous on \mathbb{R} .

Now let $x_0 \in I$, and $r > 0$ such that $[x_0 - r, x_0 + r] \subset I$. We then define the function $\varphi_{x_0, r} : I \rightarrow \mathbb{R}^+$ by

$$\varphi_{x_0, r}(x) = \varphi(r^2 - |x - x_0|^2), \quad \forall x \in \mathbb{R}.$$

This function is smooth, with $\text{supp } \varphi_{x_0, r} = [x_0 - r, x_0 + r]$. In particular it belongs to $\mathcal{D}(I)$.

(A different construction of a function in $\mathcal{D}(\mathbb{R})$ is given in TD1, Ex. 7). □

We will often use a particular type of test function, equal to unity on some interval.

Proposition 1.2.4 (Cutoff functions) Let $I \subset \mathbb{R}$ an open set, and a segment $[a, b] \subset I$. There exists a function $\psi \in \mathcal{D}(I)$ such that

- i) $\psi = 1$ on $[a, b]$,
- ii) $\forall x \in I, \psi(x) \in [0, 1]$.

In French, such a function is called “fonction plateau”, while in English it is rather referred to as a “cut-off function”.

Proof.— First of all, we prove that, for every $\alpha < \beta$, there exists $\chi_{\alpha, \beta} \in \mathcal{C}^\infty(\mathbb{R})$ such that

- i) $\chi_{\alpha, \beta} = 0$ on $] -\infty, \alpha]$,
 - ii) $\forall x \in \mathbb{R}, \chi_{\alpha, \beta}(x) \in [0, 1]$,
 - iii) $\chi_{\alpha, \beta} = 1$ on $[\beta, +\infty[$.
- (such a $\chi_{\alpha, \beta}$ is called a “smooth step function”).

Let $\varphi_{x_0, r} \in \mathcal{C}^\infty(\mathbb{R})$ be the function defined in Prop. 1.2.3, with the parameters

$$x_0 = \frac{\alpha + \beta}{2}, \quad r = \frac{\beta - \alpha}{2}.$$

Then it is easy to check that the denominator below is strictly positive, and that

$$\chi_{\alpha, \beta}(x) \stackrel{\text{def}}{=} \frac{\int_{-\infty}^x \varphi_{x_0, r}(t) dt}{\int_{-\infty}^{+\infty} \varphi_{x_0, r}(t) dt}$$

satisfies the required properties.

Coming back to $[a, b] \subset I$, select $a', b' \in I$ such that $a' < a < b < b'$. Then we may just take the product

$$\psi(x) = \chi_{a', a}(x)(1 - \chi_{b, b'}(x))$$

□

1.2.2 Smooth partitions of unity

An important use of these cutoff functions lies in the construction of *smooth partitions of unity*. We start by defining the notion of *open cover* of a compact set.

Definition 1.2.5 (Open cover) Fix I an open interval, and let $K \subset I$ be a compact (bounded and closed) subset of I . A finite open cover of K inside I is a family of open intervals $I_1, \dots, I_n \subset I$, such that

$$K \subset \bigcup_{i=1}^n I_n.$$

To any open cover we may associate a smooth partition of unity.

Proposition 1.2.6 (Smooth partitions of unity) Let $K \subset I$ be a compact subset of the open interval I , and let $I_1, \dots, I_n \subset I$ be a finite open cover of K .

Then, there exist functions $\chi_1 \in \mathcal{C}_0^\infty(I_1, [0, 1]), \dots, \chi_n \in \mathcal{C}_0^\infty(I_n, [0, 1])$, such that their sum

$$\chi = \sum_{j=1}^n \chi_j$$

is equal to unity on all of K .

As a consequence of this smooth partition of unity, we easily draw the following

Corollary 1.2.7 Let $\varphi \in \mathcal{C}_0^\infty(I)$, and assume $I_1, \dots, I_n \subset I$ form a finite open cover of $\text{supp } \varphi$ inside I :

$$\text{supp } \varphi \subset I_1 \cup \dots \cup I_n .$$

Then there exist $\varphi_1 \in \mathcal{C}_0^\infty(I_1), \dots, \varphi_n \in \mathcal{C}_0^\infty(I_n)$, such that

$$\varphi = \varphi_1 + \dots + \varphi_n .$$

Indeed, once we have available a partition of unity χ_i, \dots, χ_n associated with the cover of $\text{supp } \varphi$, it suffices to take $\varphi_j = \chi_j \varphi$ for $j = 1, \dots, n$. \square




Figure 1.2: Two cutoff functions realizing a partition of unity associated with the open cover $K \subset I_1 \cup I_2$, as in Proposition 1.2.6.

Proof.— Let us now prove Proposition 1.2.6. We start with an elementary

Lemma 1.2.8 (“Shrinking lemma”) Take I_1, \dots, I_n a family of open subintervals of I . Then, for every compact subset K such that

$$K \subset I_1 \cup \dots \cup I_n ,$$

there exist segments¹ $[a_1, b_1] \subset I_1, \dots, [a_n, b_n] \subset I_n$ such that

$$K \subset]a_1, b_1[\cup \dots \cup]a_n, b_n[.$$

To prove this Lemma we proceed by induction on n . For $n = 1$, K is a compact subset of the open interval I_1 , hence there exists a segment $[\alpha_1, \beta_1] \subset I_1$ such that $K \subset [\alpha_1, \beta_1]$. For instance, one may choose $\alpha_1 = \min K$, $\beta_1 = \max K$. Then we just choose $a_1, b_1 \in I$ such that $a_1 < \alpha_1 < \beta_1 < b_1$.

Assume now that for some $n \geq 2$, the result is true at the level $n - 1$; let us and prove that it holds at the level n . The set

$$K' = K \setminus I_n$$

is closed and contained in K , hence it is a compact subset of I , and

$$K' \subset I_1 \cup \cdots \cup I_{n-1}.$$

Applying the induction assumption, there exist $[a_1, b_1] \subset I_1, \dots, [a_{n-1}, b_{n-1}] \subset I_{n-1}$ such that

$$K' \subset]a_1, b_1[\cup \cdots \cup]a_{n-1}, b_{n-1}[.$$

Now consider the compact subset

$$K'' = K \setminus \left(]a_1, b_1[\cup \cdots \cup]a_{n-1}, b_{n-1}[\right) \subset I_n.$$

Applying at the level 1, there exists a segment $[a_n, b_n] \subset I_n$ such that

$$K'' \subset]a_n, b_n[.$$

As a consequence, we obtain

$$K \subset]a_1, b_1[\cup \cdots \cup]a_n, b_n[,$$

as announced. □

Let us use this Lemma to prove Proposition 1.2.6. Let $[a_1, b_1], \dots, [a_n, b_n]$ be the segments provided by the Lemma. From Proposition 1.2.4 we may construct cutoff functions $\psi_1 \in \mathcal{C}_0^\infty(I_1, [0, 1]), \dots, \psi_n \in \mathcal{C}_0^\infty(I_n, [0, 1])$ such that $\psi_1 = 1$ on $[a_1, b_1], \dots, \psi_n = 1$ on $[a_n, b_n]$. We define

$$\begin{aligned} \chi_1 &= \psi_1, \\ \chi_2 &= \psi_2(1 - \psi_1), \\ &\dots \\ \chi_n &= \psi_n(1 - \psi_{n-1}) \dots (1 - \psi_1). \end{aligned}$$

Then $\chi_1 \in \mathcal{D}(I_1, [0, 1]), \dots, \chi_n \in \mathcal{D}(I_n, [0, 1])$, and the difference

$$\begin{aligned} 1 - (\chi_1 + \cdots + \chi_n) &= 1 - \psi_1 - \psi_2(1 - \psi_1) - \cdots - \psi_n(1 - \psi_{n-1}) \dots (1 - \psi_1) \\ &= (1 - \psi_1)(1 - \psi_2 - \cdots - \psi_n(1 - \psi_{n-1}) \dots (1 - \psi_2)) \\ &= (1 - \psi_1) \dots (1 - \psi_n). \end{aligned}$$

By construction, the function $(1 - \psi_1) \dots (1 - \psi_n)$ vanishes identically on $[a_1, b_1] \cup \cdots \cup [a_n, b_n] \supset K$, so we get the result. □

1.2.3 Convergence in the space of test functions

Once we have defined the space of test functions $\mathcal{D}(I)$, it is important to define a proper notion of convergence on this space, or equivalently a topology on $\mathcal{D}(I)$. Test functions are smooth and compactly supported, it is natural that the topology on $\mathcal{D}(I)$ takes into account all these properties.

The natural notion of convergence for continuous functions is that of uniform convergence, since it is the simplest one for which the limit of a sequence of continuous functions is continuous. For test functions, the notion of convergence is provided by the next definition.

Definition 1.2.9 Let $(\varphi_j)_{j \in \mathbb{N}}$ be a sequence of functions in $\mathcal{D}(I)$, and $\varphi \in \mathcal{D}(I)$. We say that (φ_j) tends to φ in $\mathcal{D}(I)$ (or in the $\mathcal{D}(I)$ -sense), if:

- i) *Uniform support* There exists a segment $[a, b] \subset I$ such that $\text{supp } \varphi_j \subset [a, b]$ for all $j \in \mathbb{N}$.
- ii) *Uniform convergence of all derivatives* For any $k \in \mathbb{N}$, $\|\varphi_j^{(k)} - \varphi^{(k)}\|_\infty := \sup_{x \in I} |\varphi_j^{(k)}(x) - \varphi^{(k)}(x)| \rightarrow 0$ as $j \rightarrow \infty$.

We denote this convergence by

$$\varphi = \mathcal{D} - \lim_{j \rightarrow +\infty} \varphi_j.$$

Obviously, when it exists, the limit function φ of a sequence $(\varphi_j)_{j \in \mathbb{N}}$ is unique.

Remark 1.2.10 Under the conditions of the above definition, the limit function necessarily satisfies $\text{supp } \varphi \subset [a, b]$.

Exercise 1.2.11 Assuming the uniform convergence $\|\varphi_j - \varphi\|_\infty \rightarrow 0$ alone does not imply, a priori, the uniform convergence of derivatives.

Construct a counterexample.

Exercise 1.2.12 Let $\varphi \in \mathcal{D}(\mathbb{R})$, and, for $t \neq 1$, denote by $\psi_t \in \mathcal{D}(\mathbb{R})$ the function given by

$$\psi_t(x) = \frac{\varphi(tx) - \varphi(x)}{t - 1}.$$

Show that the family (ψ_t) converges in $\mathcal{D}(\mathbb{R})$ as t tends to 1.

Remark 1.2.13 The definition 1.2.9 of the convergence of sequences in $\mathcal{D}(I)$ corresponds to a slightly subtle topology on $\mathcal{D}(I)$, called the *strict inductive topology*. Why “inductive”? Because one first defines a topology on the space of smooth functions supported on a given segment $K \subset I$ (let us denote this space by $\mathcal{D}_K(I)$). This topology is defined through the family of seminorms

$$N_{K,k}(\varphi) = \max_{0 \leq j \leq k} \|\varphi^{(j)}\|_\infty, \quad \varphi \in \mathcal{D}_K(I),$$

namely a local base of open neighbourhoods in $\mathcal{D}_K(I)$ of the zero function can be defined by

$$U_{K,k} = \{\varphi \in \mathcal{D}_K(I), N_{K,k}(\varphi) < 1/k\}, \quad k \geq 1.$$

With this base, one recovers the fact that a sequence $(\varphi_j) \subset \mathcal{D}_K(I)$ converges to zero in $\mathcal{D}_K(I)$ iff all seminorms $N_{K,k}(\varphi_j)$ converge to zero when $j \rightarrow \infty$.

To induce a topology on $\mathcal{D}(I)$, one considers a growing sequence of segments $K_1 \subset \cdots \subset K_i \subset K_{i+1} \subset \cdots$, such that

$$I = \bigcup_{i \geq 1} K_i.$$

It is obvious that for any i and any function $\varphi \in \mathcal{D}_{K_i}(I)$, that function is also an element of $\mathcal{D}_{K_{i+1}}(I)$, and the seminorms $N_{K_{i+1},k}(\varphi) = N_{K_i,k}(\varphi)$. As a result, the intersection between the open set $U_{K_{i+1},k} \subset \mathcal{D}_{K_{i+1}}(I)$ and $\mathcal{D}_{K_i}(I)$ is just the set $U_{K_i,k}$, which is open in $\mathcal{D}_{K_i}(I)$; one says that the topology on $\mathcal{D}_{K_i}(I)$ is induced by the topology on $\mathcal{D}_{K_{i+1}}(I)$.

There exists a single topology on $\mathcal{D}(I)$, the strict inductive topology, such that for each open set $U \subset \mathcal{D}(I)$ of this topology, the of $U \cap \mathcal{D}_{K_i}(I)$ is an open set for the topology of $\mathcal{D}_{K_i}(I)$. A family of neighbourhood of the zero function in $\mathcal{D}(I)$ is given by the sets

$$U_k = \bigcup_{i \geq 1} U_{K_i,k}.$$

Let us quote a few properties of this inductive topology:

1. each $\mathcal{D}_{K_i}(I)$ is a closed subspace of $\mathcal{D}(I)$.
2. each space $\mathcal{D}_{K_i}(I)$ is separated and complete w.r.to its topology, as a consequence $\mathcal{D}(I)$ is separated and complete in the inductive topology.
3. although the topology on each $\mathcal{D}_{K_i}(I)$ is metrizable, the strict inductive topology on $\mathcal{D}(I)$ is *not* metrizable.
4. a sequence $(\varphi_j) \subset \mathcal{D}(I)$ converges to some $\varphi_0 \in \mathcal{D}(I)$ iff the two properties of Definition 1.2.9 are satisfied.

1.3 Definition of distributions on I

Equipped with our test functions, we are now able to formally define the notion of distribution on an open interval I . As mentioned in the Introduction, distributions correspond to a certain class of linear forms acting on test functions.

1.3.1 Definitions

Definition 1.3.1 Let $I \subset \mathbb{R}$ an open subset, and T a complex valued linear form on $\mathcal{D}(I)$. One says that T is a distribution on I if, for every segment $[a, b] \subset I$,

$$(1.3.1) \quad \exists C > 0, \exists m \in \mathbb{N}, \forall \varphi \in \mathcal{D}(I) \text{ with } \text{supp } \varphi \subset [a, b], |T(\varphi)| \leq C \sum_{\alpha \leq m} \|\varphi^{(\alpha)}\|_{\infty}.$$

We denote by $\mathcal{D}'(I)$ the set of distributions on I , and for $T \in \mathcal{D}'(I)$, $\varphi \in \mathcal{D}(I)$, we use the bracket notations

$$\langle T, \varphi \rangle = \langle T, \varphi \rangle_{\mathcal{D}'(I), \mathcal{D}(I)} := T(\varphi).$$

From the bound (1.3.1), we say that the distributional bracket $\langle T, \varphi \rangle$ is *controlled by m derivatives of φ* .

The above definition may appear a bit artificial. The following Proposition provides a more natural characterization, in terms of the topology of the space of test functions.

Proposition 1.3.2 (Continuity) A linear form T on $\mathcal{D}(I)$ is a distribution on I if and only if, for any sequence (φ_j) of functions in $\mathcal{D}(I)$ that converges to φ in the $\mathcal{D}(I)$ -sense, one has $T(\varphi_j) \rightarrow T(\varphi)$. In other words, the linear form $T : \mathcal{D}(I) \rightarrow \mathbb{C}$ is a distribution iff it is *continuous*² with respect to the topology of $\mathcal{D}(I)$.

Proof.— Let T be a distribution on I , and (φ_j) a sequence in $\mathcal{D}(I)$ which converges to φ in $\mathcal{D}(I)$. There is a segment $[a, b] \subset I$ such that $\text{supp } \varphi_j \subset [a, b]$ for all $j \in \mathbb{N}$, and $\text{supp } \varphi \subset [a, b]$. There exist $C > 0$ and $m \in \mathbb{N}$ such that

$$\forall \psi \in \mathcal{D}(I) \text{ with } \text{supp } \psi \subset [a, b], \quad |T(\psi)| \leq C \sum_{\alpha \leq m} \|\psi^{(\alpha)}\|_{\infty}.$$

In particular, for any $j \in \mathbb{N}$,

$$|T(\varphi_j) - T(\varphi)| = |T(\varphi_j - \varphi)| \leq C \sum_{\alpha \leq m} \|\varphi_j^{(\alpha)} - \varphi^{(\alpha)}\|_{\infty}.$$

Therefore $T(\varphi_j) \rightarrow T(\varphi)$ as $j \rightarrow +\infty$, and we have proved the *only if* part of the proposition.

We now want to prove the *if* part. Suppose that for any sequence (φ_j) of functions which converges in $\mathcal{D}(I)$, we have $T(\varphi_j) \rightarrow T(\varphi)$, where $\varphi = \mathcal{D} - \lim \varphi_j$. We will reason *ab absurdo*: let us assume that the linear form T is *not* a distribution: the converse of (1.3.1) gives the long statement:

$$(1.3.2) \quad \exists [a, b] \subset I, \forall C > 0, \forall m \in \mathbb{N}, \exists \varphi \in \mathcal{D}(\Omega) \text{ with } \text{supp } \varphi \subset [a, b],$$

$$\text{such that } |T(\varphi)| > C \sum_{\alpha \leq m} \|\varphi^{(\alpha)}\|_{\infty}.$$

(Notice that φ cannot be the null function). In particular, for any $j \in \mathbb{N}$, choosing $C = m = j$, there is a test function $\varphi_j \in \mathcal{D}(I)$ such that $\text{supp } \varphi_j \subset [a, b]$ and

$$|T(\varphi_j)| > j \sum_{\alpha \leq j} \|\varphi_j^{(\alpha)}\|_{\infty}.$$

Let $\psi_j \in \mathcal{D}(I)$ be defined by $\psi_j := \varphi_j / |T(\varphi_j)|$. One obviously has $|T(\psi_j)| = 1$, with $\text{supp } \psi_j \subset [a, b]$. On the other hand, for any given $\alpha \in \mathbb{N}$ and any $j \geq \alpha$, one has the bound

$$\|\psi_j^{(\alpha)}\|_{\infty} \leq \sum_{\beta \leq j} \|\psi_j^{(\beta)}\|_{\infty} < \frac{1}{j},$$

therefore $\lim_{j \rightarrow \infty} \|\psi_j^{(\alpha)}\|_{\infty} = 0$; this proves the limit

$$\mathcal{D} - \lim_{j \rightarrow \infty} \psi_j = 0.$$

The continuity of T then imposes $T(\psi_j) \rightarrow 0$, which contradicts the normalization $|T(\psi_j)| = 1$. Hence the *if* part of the Proposition is proved. \square

Remark 1.3.3 As a set of continuous linear forms, $\mathcal{D}'(I)$ forms a vector space on \mathbb{C} : if $T_1, T_2 \in \mathcal{D}'(I)$ and $\lambda_1, \lambda_2 \in \mathbb{C}$, then $\lambda_1 T_1 + \lambda_2 T_2$ is the linear form given by

$$\langle \lambda_1 T_1 + \lambda_2 T_2, \varphi \rangle := \lambda_1 \langle T_1, \varphi \rangle + \lambda_2 \langle T_2, \varphi \rangle,$$

and this form is obviously continuous since T_1, T_2 are so.

For $T \in \mathcal{D}'(I)$, we may also denote by \bar{T} or T^* the distribution given by

$$\langle \bar{T}, \varphi \rangle = \overline{\langle T, \bar{\varphi} \rangle}.$$

Then, any distribution T can be written as $T = T_1 + iT_2$ where T_1 and T_2 are *real distributions*, that is such that $\langle T, \varphi \rangle \in \mathbb{R}$ for any real valued function φ . Indeed, this relation holds with

$$T_1 = \frac{1}{2}(T + \bar{T}) = \operatorname{Re}(T) \quad \text{and} \quad T_2 = \frac{1}{2i}(T - \bar{T}) = \operatorname{Im}(T).$$

Equipped with this general definition, we will now provide some examples of distributions.

1.3.2 Distributions defined by locally integrable functions

The first distributions we meet are well-known objects, namely locally integrable functions.

If $f \in L^1_{\text{loc}}(I)$, the function $f\varphi$ is integrable for any $\varphi \in \mathcal{D}(I)$, and

$$T_f : \varphi \rightarrow \int_I f(x)\varphi(x) dx$$

is a linear form on $\mathcal{D}(I)$. Moreover, if $[a, b] \subset I$ is a segment, for any $\varphi \in \mathcal{D}(I)$ supported in $[a, b]$, one easily checks that

$$|T_f(\varphi)| \leq \|f\|_{L^1([a,b])} \sup |\varphi|.$$

This bound shows that T_f is a distribution on I , and that $T_f(\varphi)$ is controlled by 0 derivative of φ .

As a matter of fact, one can identify $L^1_{\text{loc}}(I)$ with a part of $\mathcal{D}'(I)$, that is identify f with T_f . This is the content of the

Proposition 1.3.4 The linear map

$$f \in L^1_{\text{loc}}(I) \mapsto T_f \in \mathcal{D}'(I)$$

is injective.

Remark 1.3.5 The statement $f = g$ for $f, g \in L^1_{\text{loc}}(I)$ should be understood in the sense of classes for the equivalence on functions : $f \sim g$ if $f(x) = g(x)$ almost everywhere (a.e.) In other words, if f is a locally integrable function, $T_f = 0$ if and only if $f = 0$ almost everywhere. However, in the special case $f \in \mathcal{C}^0(I)$, this is equivalent to $f = 0$ everywhere. Indeed, the only open set with zero Lebesgue measure is the empty set.

Proof.— Assume $T_f = 0$. It is enough to prove that $f = 0$ a.e. on every segment of I , or equivalently that $\psi f = 0$ for every $\psi \in \mathcal{D}(I)$.

Our first task is to approximate the function f by a smooth function. For this aim, we will use a family of *convolution kernels* [see Ex. 6 in TD1].

Let $\rho \in \mathcal{D}(\mathbb{R})$, non negative, supported in $[-1, 1]$, and such that

$$\int_{\mathbb{R}} \rho(x) dx = 1 .$$

We then rescale ρ by a factor³ $\varepsilon > 0$, into the function

$$\rho_{\varepsilon}(x) = \frac{1}{\varepsilon} \rho\left(\frac{x}{\varepsilon}\right) .$$

We notice that the integral of ρ_{ε} remains equal to unity, while $\rho_{\varepsilon} \in \mathcal{D}(\mathbb{R})$ is supported in $[-\varepsilon, \varepsilon]$.

Lemma 1.3.6 (Smoothing by convolution) Let $g \in L^1(\mathbb{R})$. Then the convolution product

$$g_{\varepsilon} = g * \rho_{\varepsilon}$$

converges to g in $L^1(\mathbb{R})$ when ε tends to 0.

Indeed, we have

$$\begin{aligned} |g_{\varepsilon}(x) - g(x)| &\leq \left| \int g(y) \rho_{\varepsilon}(x - y) dy - g(x) \right| \\ &\leq \left| \int g(x - \varepsilon z) \rho(z) dz - g(x) \right| \leq \int |g(x - \varepsilon z) - g(x)| \rho(z) dz \end{aligned}$$

Then by Fubini-Tonelli,

$$\|g_{\varepsilon} - g\|_{L^1} \leq \int \left(\int |g(x - \varepsilon z) - g(x)| \rho(z) dz \right) dx \leq \int \|\tau_{\varepsilon z} g - g\|_{L^1} \rho(z) dz,$$

where $\tau_a g$ denotes the translation of the function g given by $\tau_a g(x) = g(x - a)$. We now show that the above expression vanishes in the limit $\varepsilon \rightarrow 0$. For this we will use the dominated convergence theorem. Firstly, the domination is obvious:

$$\|\tau_{\varepsilon z} g - g\|_{L^1} \rho(z) \leq 2\|g\|_{L^1} \rho(z).$$

Secondly, the continuity of the translation group acting on the L^p spaces for $p \in [1, \infty[$, namely

$$\|\tau_{\varepsilon z} g - g\|_{L^1} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0 \text{ .[see Ex.1 in TD1],}$$

shows the simple convergence of the integrand. The dominated convergence theorem then implies the convergence of the integral.

Coming back to the proof of the proposition, we have for any $x \in \mathbb{R}$:

$$(\psi f) * \rho_{\varepsilon}(x) = \int f(y) \psi(y) \rho_{\varepsilon}(x - y) dy = \langle T_f, \psi \rho_{\varepsilon}(x - \cdot) \rangle = 0,$$

where we used the fact that $y \mapsto \psi(y) \rho_{\varepsilon}(x - y)$ belongs to $\mathcal{D}(I)$. Applying the lemma to $g = \psi f \in L^1$, we conclude $\psi f = 0$ in L^1 , which completes the proof. \square

3. In general, in analysis the letter ε is used for a "small" parameter.

1.3.3 The Dirac mass

The next example of distribution is a genuinely new object, in the sense that it is more singular than a function. It admits various names: Dirac distribution, Dirac mass, Dirac measure, Dirac delta function,..., all associated with the physicist Paul Dirac, which introduced it in quantum mechanics (and called it “ δ ”).

For a point $x_0 \in I$, we denote by $\delta_{x_0} : \mathcal{D}(I) \rightarrow \mathbb{C}$ the linear form defined by

$$\delta_{x_0}(\varphi) := \varphi(x_0).$$

For any function $\varphi \in \mathcal{D}(I)$, one obviously has

$$|\delta_{x_0}(\varphi)| \leq \|\varphi\|_\infty,$$

so that δ_{x_0} is a distribution on I . It is called the *Dirac mass* at x_0 .

We notice that $\delta_{x_0}(\varphi)$ is controlled by zero derivative of φ , like in the case of distributions T_f associated with an L^1_{loc} function f . Yet, we claim that the Dirac mass at x_0 cannot be defined by a locally integrable function f . Indeed, otherwise we would have, for any function $\varphi \in \mathcal{D}(I)$ such that $x_0 \notin \text{supp } \varphi$,

$$\varphi(x_0) = 0 = \int f(x)\varphi(x)dx.$$

Let us split the interval I at the point x_0 , getting the disjoint union

$$I = I_1 \sqcup \{x_0\} \sqcup I_2.$$

Then for any $\varphi_1 \in \mathcal{D}(I_1)$ we will have

$$\int_I f(x)\varphi_1(x) dx = \int_{I_1} f|_{I_1}(x)\varphi_1(x) dx = 0.$$

The second integral is the distributional bracket associated with the function $f|_{I_1} \in L^1_{loc}(I_1)$. The Proposition 1.3.4 applied to f_1 , then implies that $f|_{I_1} = 0$ a.e. in I_1 . The same reasoning applies to the restriction $f|_{I_2}$, which hence vanishes a.e. on I_2 . Finally, $f = 0$ a.e. in I .

Now, if we select a plateau function ψ such that $\psi(x_0) = 1$, we would have

$$1 = \psi(x_0) = \langle T_f, \psi \rangle = \int_I f\psi = 0,$$

which is a contradiction.

The Dirac mass is relevant in many contexts. Among the 4 motivations we presented in the introduction, 3 were involving the Dirac mass:

- i) the derivative of the Heaviside function is the Dirac distribution at the origin (see below section (1.4))
- ii) in electrostatics, a point charge at the origin can be modeled by a “density” given by the Dirac mass
- iii) in quantum mechanics, we already explained that the quantum state localized at a point x is modeled by the Dirac mass at x .

1.3.4 A distribution involving an infinite number of derivatives

In this section we want to explain why, in the bound (1.3.1) in the definition of a distribution, the number m of derivatives needed to control $T(\varphi)$ may depend on the support of φ (hence on the segment $[a, b]$ containing this support).

Indeed, we construct below a distribution such that $m = m([a, b])$ converges to infinity when $[a, b]$ is made larger and larger, up to exhausting all of I .

We take $I =]0, 1[$. Consider a sequence $(x_n)_{n \in \mathbb{N}}$ of points of $]0, 1[$ converging to 0 as n tends to infinity. Let $(a_n)_{n \in \mathbb{N}}$ a sequence of complex numbers. For every $\varphi \in \mathcal{D}(I)$, we claim that

$$(1.3.3) \quad \langle T, \varphi \rangle = \sum_{n=0}^{\infty} a_n \varphi^{(n)}(x_n)$$

defines a distribution on I . Indeed, let $a < b$ be points in $]0, 1[$ such that $\text{supp}(\varphi) \subset [a, b]$. Then there exists an integer N such that, for every $n > N$, $x_n < a$, so that

$$\sum_{n=0}^{\infty} a_n \varphi^{(n)}(x_n) = \sum_{n=0}^N a_n \varphi^{(n)}(x_n)$$

is well defined, and controlled by N derivatives of φ . For the same reason, if a sequence $(\varphi_k)_{k \in \mathbb{N}}$ converges to φ in $\mathcal{D}(I)$, then there exists a segment $[a, b] \subset I$ containing the supports of all the φ_k and of φ , so there will exist some N such that

$$\langle T, \varphi_k \rangle = \sum_{n=0}^N a_n \varphi_k^{(n)}(x_n) \xrightarrow{k \rightarrow \infty} \sum_{n=0}^N a_n \varphi^{(n)}(x_n) = \langle T, \varphi \rangle.$$

Remark 1.3.7 In this example it is crucial that the sequence $(x_n)_{n \in \mathbb{N}}$ does not admit an accumulation point in I . Indeed, if there were such an accumulation point $x_a \in I$, then for a given choice of parameters (a_n) (assuming infinitely many of these parameters are nonzero), one can construct a test function $\varphi \in \mathcal{D}(I)$ such that the derivatives $\varphi^{(n)}(x_a)$ grow arbitrarily fast when $n \rightarrow \infty$, in particular such as to make the (1.3.3) divergent. Hence the linear form T would not be defined on this function φ .

1.3.5 Cauchy's principal value of $1/x$

We now introduce a distribution which looks similar to the distributions associated with L_{loc}^1 functions. However, in the present case the function is $1/x$, which is not locally integrable at the origin. For this reason, one has to be a bit clever and proceed by a limiting argument, removing smaller and smaller neighbourhoods of the origin. We will see that the resulting distribution is more singular than distributions T_f associated with L_{loc}^1 functions.

Precisely, we consider the linear form $T : \mathcal{D}(\mathbb{R}) \rightarrow \mathbb{C}$ given by

$$T(\varphi) = \lim_{\varepsilon \rightarrow 0^+} \int_{|x| > \varepsilon} \frac{\varphi(x)}{x} dx.$$

Let us show that this limit exists for any $\varphi \in \mathcal{D}(\mathbb{R})$. Indeed, for such a function, we have

$$\begin{aligned} \int_{|x| > \varepsilon} \frac{\varphi(x)}{x} dx &= \int_{\varepsilon}^{+\infty} \frac{\varphi(x)}{x} dx + \int_{-\infty}^{-\varepsilon} \frac{\varphi(x)}{x} dx \\ &= \int_{\varepsilon}^{+\infty} \frac{\varphi(x) - \varphi(-x)}{x} dx \xrightarrow{\varepsilon \rightarrow 0} \int_0^{+\infty} \frac{\varphi(x) - \varphi(-x)}{x} dx. \end{aligned}$$

where we have used that the function

$$x \in \mathbb{R} \mapsto \frac{\varphi(x) - \varphi(-x)}{x}$$

is C^∞ and compactly supported (cf Ex. 1.1.1). In effect, both integrals on \mathbb{R}_+ and \mathbb{R}_- diverge, but with opposite asymptotics, such that their sum remains bounded.

Now we show that the well-defined linear form T is a distribution on \mathbb{R} . For any $\varphi \in \mathcal{D}(\mathbb{R})$, we have the bound

$$\left| \frac{\varphi(x) - \varphi(-x)}{x} \right| \leq 2\|\varphi'\|_\infty$$

and, if $\text{supp } \varphi \subset [-A, A]$,

$$\left| \int_0^{+\infty} \frac{\varphi(x) - \varphi(-x)}{x} dx \right| \leq 2A\|\varphi'\|_\infty.$$

This shows that T satisfies the estimate in Definition 1.3.1. Notice that the constant in the estimate is here $2A$, it does indeed depend on the support of φ . Another striking fact is that this distribution is controlled by the first derivative of φ . In this sense, it is more singular than the distributions T_f associated with locally integrable functions.

This distribution is called the *Cauchy principal value* of $1/x$,⁴ and we denote it by

$$\langle \text{pv} \left(\frac{1}{x} \right), \varphi \rangle = \lim_{\varepsilon \rightarrow 0^+} \int_{|x| > \varepsilon} \frac{\varphi(x)}{x} dx = \int_0^{+\infty} \frac{\varphi(x) - \varphi(-x)}{x} dx.$$

There are other examples of extensions of functions near a singular point (Hadamard finite parts,...), which can often be seen as a *renormalization* procedure of a divergent expression: one subtracts the "trivial" divergent behaviour, to identify a bounded remainder, the "renormalized" expression.

1.4 Differentiating distributions

This section introduces one of the most important operation on distributions, namely the derivative. In order to understand the definition below, let us notice that, if $f \in C^1(I)$, a simple integration by parts yields

$$\forall \varphi \in \mathcal{C}_0^\infty(I), \quad \langle T_{f'}, \varphi \rangle = \int_I f'(x)\varphi(x) dx = - \int_I f(x)\varphi'(x) dx = -\langle T_f, \varphi' \rangle.$$

This suggest the following definition.

Proposition 1.4.1 Let $T \in \mathcal{D}'(I)$. The linear form on $\mathcal{D}(I)$ defined by

$$\varphi \mapsto -\langle T, \varphi' \rangle$$

is a distribution, that we call the derivative of T , and that we denote by T' .

Proof.— Let $[a, b] \subset I$ be a segment, and $C > 0$, $m \in \mathbb{N}$ the constants given by the fact that $T \in \mathcal{D}'(I)$. For $\varphi \in \mathcal{D}(I)$, supported in $[a, b]$, we have

$$|\langle T', \varphi \rangle| = |\langle T, \varphi' \rangle| \leq C \sum_{\alpha \leq m} \sup |\varphi^{(1+\alpha)}| \leq C \sum_{\alpha \leq m+1} \sup |\varphi^{(\alpha)}|,$$

4. In French, it is called "valeur principale de $1/x$ ", hence denoted as $\text{vp} \left(\frac{1}{x} \right)$.

which shows that T' is a distribution, a little more singular than T . □

Of course the above calculation gives immediately the nice property:

Proposition 1.4.2 If $T = T_f$ with $f \in C^1(I)$, then $T' = T_{f'}$.

The next examples are much more interesting.

Example 1.4.3 Let $H : \mathbb{R} \rightarrow \mathbb{C}$ denote the Heaviside function, namely $H(x) = \mathbf{1}_{\mathbb{R}^+}(x)$. The function H belongs to $L^1_{\text{loc}}(\mathbb{R})$, so we can denote by $T = T_H$ the associated distribution. For $\varphi \in \mathcal{D}(\mathbb{R})$,

$$\langle T', \varphi \rangle = -\langle T, \varphi' \rangle = -\int_0^{+\infty} \varphi'(x) dx = \varphi(0),$$

so that $T'_H = \delta_0$, as we stated above: in the sense of distributions, the derivative of the Heaviside function is the delta mass at the origin.

Example 1.4.4 Let $f \in L^1_{\text{loc}}(I)$ and $a \in I$. Consider

$$F(x) = \int_a^x f(t) dt .$$

A classical application of the dominated convergence theorem claims that F is continuous on I . In particular, F is locally integrable. We claim that

Proposition 1.4.5

$$T'_F = T_f .$$

Proof.— Introduce, for every $t \in I$,

$$I_{<t} = \{x \in I : x < t\} , \quad I_{>t} = \{x \in I : x > t\} .$$

Let us compute, for every $\varphi \in \mathcal{C}_0^\infty(I)$,

$$\begin{aligned} \langle T'_F, \varphi \rangle &= -\langle T_F, \varphi' \rangle = -\int_I \left(\int_a^x f(t) dt \right) \varphi'(x) dx \\ &= \int_{I_{<a}} \left(\int_{]x,a[} f(t) dt \right) \varphi'(x) dx - \int_{I_{>a}} \left(\int_{]a,x[} f(t) dt \right) \varphi'(x) dx . \end{aligned}$$

Let us apply the Fubini theorem to both integrals in the right hand side. Notice that, if the support of φ is included into $[\alpha, \beta]$ with $a \in [\alpha, \beta]$, the integrand of both integrals is supported by $[\alpha, \beta] \times [\alpha, \beta]$ and is bounded by

$$|f(t)| |\varphi'(x)|$$

which is integrable on $[\alpha, \beta] \times [\alpha, \beta]$. Hence the Fubini theorem allows us to interchange the orders of integration. This yields

$$\begin{aligned} \langle T'_F, \varphi \rangle &= \int_{I_{<a}} f(t) \left(\int_{I_{<t}} \varphi'(x) dx \right) dt - \int_{I_{>a}} f(t) \left(\int_{I_{>t}} \varphi'(x) dx \right) dt \\ &= \int_{I_{<a}} f(t) \varphi(t) dt + \int_{I_{>a}} f(t) \varphi(t) dt \\ &= \int_I f(t) \varphi(t) dt = \langle T_f, \varphi \rangle . \end{aligned}$$

□ In this proof we notice a “meta-strategy” which will be used at many points in this course: transfer the computations (differentiation, product with a smooth function) from the distribution side to the test function side.

Remark 1.4.6 [Going further than Lebesgue !] The statement of Prop. 1.4.5 looks very similar with the one of Prop. 1.4.2. However the extension from $f \in C^1$ to $f \in L^1_{loc}$ contains some subtleties.

1. The derivative f is not defined everywhere, yet the corresponding distribution is unique, as explained in Prop. 1.3.4. On the other hand, the function F is continuous, hence unambiguously defined.

2. Lebesgue proved that the function $F(x) = \int^x f$ is almost everywhere differentiable, with a derivative equal to f ; in particular F can be recovered from its derivative. This is not a general property of continuous functions⁵. Indeed, there exists continuous functions G , differentiable almost everywhere, but such that their (a.e. defined) derivative G' does not allow to reconstruct G . For instance, the “devil’s staircase” function G is continuous on $[0, 1]$, almost everywhere differentiable with $G'(x) = 0$ a.e., yet G is not a constant function (see fig. 1.3). This uncomfortable phenomenon does not happen with distributional derivatives, as we shall see below in Proposition 1.4.8: for such a function G the distributional derivative T'_G is not associated with an L^1_{loc} function, but is more singular; yet it vanishes on a union of intervals of full measure on $]0, 1[$.




Figure 1.3: Left: the “devil staircase” function G , which satisfies $G'(x) = 0$ a.e. Right: $f(x) = \log(|x|)$ and its derivative.

Example 1.4.7 Let $f \in L^1_{loc}(\mathbb{R})$ be the function given by $f(x) = \ln(|x|)$, and $T = T_f \in \mathcal{D}'(\mathbb{R})$ be the associated distribution. We want to compute T' ; it seems reasonable that T' is related to the function $x \mapsto 1/x$, but this one is not in $L^1_{loc}(\mathbb{R})$. However, let $\varphi \in C^\infty_0([-A, A])$. We compute

$$\begin{aligned} \langle T', \varphi \rangle &= - \int_{\mathbb{R}} \ln|x| \varphi'(x) dx = - \lim_{\varepsilon \rightarrow 0} \int_{|x| > \varepsilon} \varphi'(x) \ln|x| dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_{|x| > \varepsilon} \frac{\varphi(x)}{x} dx + \lim_{\varepsilon \rightarrow 0} (\varphi(\varepsilon) - \varphi(-\varepsilon)) \ln(\varepsilon) \end{aligned}$$

5. This property actually characterizes *absolute continuity* of the function F , which is a stronger property than continuity

and the second limit in the right hand side is 0, since φ is smooth. We conclude that

$$T' = \text{pv}(1/x),$$

the notation introduced in the previous paragraph.

The only functions with an identically null derivative on a full interval are constant functions. The same result holds true for distributions.

Proposition 1.4.8 If $T \in \mathcal{D}'(I)$ satisfies $T' = 0$, then T is associated with a constant function.

The second point of Remark 1.4.6 shows that this statement is not completely obvious. Notice that this is the first differential equation we are solving in the space of distributions. **Proof.**— We start the proof by a remark that may be useful in other contexts. A function $\varphi \in \mathcal{D}(I)$ is the derivative of a function $\psi \in \mathcal{D}(I)$ if and only if $\int_I \varphi = 0$. Indeed, if $\varphi = \psi'$ for ψ supported in $[a, b] \subset I$, then

$$\int_I \varphi(x) dx = \int_a^b \psi'(x) dx = [\psi(x)]_a^b = 0.$$

Conversely, if $\int_I \varphi = 0$, the function $\psi : x \mapsto \int_{I \cap]-\infty, x[} \varphi(t) dt$ is compactly supported, with support included in any compact interval containing that of φ , and satisfies $\psi' = \varphi$.

Now let $\chi \in \mathcal{D}(I)$ a function such that $\int_I \chi = 1$. For any $\varphi \in \mathcal{D}(I)$, we split the function into the sum

$$\varphi = \varphi_1 + \varphi_2 \quad \text{with} \quad \varphi_1 = \varphi - \left(\int_I \varphi \right) \chi, \quad \text{and} \quad \varphi_2 = \left(\int_I \varphi \right) \chi,$$

that is a “massless” part and a “massive” one. Since $\int_I \varphi_1 = 0$, there exists $\psi \in \mathcal{D}(I)$ such that $\varphi_1 = \psi'$. Hence, by linearity we may write

$$\begin{aligned} \langle T, \varphi \rangle &= \langle T, \psi' \rangle + \left(\int_I \varphi \right) \langle T, \chi \rangle \\ &= -\langle T', \psi \rangle + \left(\int_I \varphi \right) \langle T, \chi \rangle. \end{aligned}$$

If we use the hypothesis $T' = 0$, this yields

$$\langle T, \varphi \rangle = C \int_I \varphi = \langle T_C, \varphi \rangle,$$

where $C = \langle T, \chi \rangle$ is independent of φ , and T_C is the distribution associated with the constant function equal to C . \square

Proposition 1.4.2 implies that the derivative of a distribution associated with a constant function is null. Therefore, the above statement is actually an equivalence.

This result can be easily extended to a differential equation with a nonhomogeneous term.

Corollary 1.4.9 Let $f \in L^1_{\text{loc}}(I)$. The distributions $T \in \mathcal{D}'(I)$ solving the equation

$$T' = T_f$$

are all of the form $T = T_F$, where

$$F(x) = \int_a^x f(t) dt + c, \quad a \in I, \quad c \in \mathbb{C}.$$

Proof.— Let $a \in I$ and define

$$F_1(x) = \int_a^x f(t) dt.$$

By Proposition 1.4.5, $T'_{F_1} = T_f = T'$. Then just apply Proposition 1.4.9 to $T - T_{F_1}$. \square

Using arguments from the proof of Proposition 1.4.8, we can actually establish the surjectivity of the mapping $T \mapsto T'$ on the whole of $\mathcal{D}'(I)$.

Proposition 1.4.10 If $S \in \mathcal{D}'(I)$, there exists $T \in \mathcal{D}'(I)$ such that $T' = S$.

Proof.— As is becoming usual, the strategy will be to push the computations to the test functions. The identity $T' = S$ is equivalent to

$$(1.4.4) \quad \forall \varphi \in \mathcal{D}(I), \quad \langle T, \varphi' \rangle = -\langle S, \varphi \rangle.$$

In view of the proof of Proposition 1.4.8, the above equation already imposes the value of T on any massless test functions φ .

To deal with massive test functions, we fix a “mass function” $\chi \in \mathcal{D}(I)$ such that $\int_I \chi dx = 1$. Given any $\varphi \in \mathcal{D}(I)$, we define its massless part as before as $\varphi_0(x) = \varphi(x) - \chi(x) \int_I \varphi$, and then define its primitive:

$$P(\varphi)(x) = \int_{t < x} \varphi_0(t) dt.$$

The function $P(\varphi)$ belongs to $\mathcal{D}(I)$, hence P defines a linear mapping

$$P : \mathcal{D}(I) \rightarrow \mathcal{D}(I), \quad \text{such that} \quad \text{supp}[P(\psi)] \subset \text{supp}(\psi) \cup \text{supp}(\chi).$$

It is easy to check that this mapping is (sequentially) continuous: if a sequence (φ_n) converges to φ in $\mathcal{D}(I)$, then $P(\varphi_n)$ converges to $P(\varphi)$ in $\mathcal{D}(I)$. Therefore the linear form T on $\mathcal{D}(I)$ defined by

$$\forall \varphi \in \mathcal{D}(I), \quad \langle T, \varphi \rangle = -\langle S, P(\varphi) \rangle,$$

is itself continuous, hence it is a distribution.

If we now apply this distribution to φ' , we check that $P(\varphi') = \varphi$, hence T indeed solves (1.4.4). \square

Since T' is itself a distribution, it can be differentiated. Iterating this process, one can define the successive derivatives $T^{(k)}$ of T for all $k \in \mathbb{N}$. This means that any distribution can be differentiated at any order, and we obtain the formula

$$\forall \varphi \in \mathcal{D}(I), \quad \langle T^{(k)}, \varphi \rangle = (-1)^k \langle T, \varphi^{(k)} \rangle.$$

For instance, the k -th derivative of the Dirac mass at the origin is defined as

$$\langle \delta_{x_0}^{(k)}, \varphi \rangle = (-1)^k \varphi^{(k)}(x_0).$$

The distribution described in subsection 1.3.4 can be expressed as

$$T = \sum_{n \geq 1} (-1)^n a_n \delta_{x_n}^{(n)}.$$

1.5 Product by a smooth function

One important property of continuous (or C^k) functions is that they form an *algebra*: one can multiply functions with one another, staying the same class of continuous (or C^k) functions. This is already not the case for two L^1 functions, yet the product of an L^1 function with a bounded (L^∞) function remains in L^1 .

Two distributions can in general not be multiplied with one another. However, we may multiply a distribution with smooth functions, as shown in the following

Proposition 1.5.1 Let $T \in \mathcal{D}'(I)$, and $f \in \mathcal{C}^\infty(I)$. The linear form

$$\varphi \mapsto \langle T, f\varphi \rangle$$

is a distribution on $D(I)$. We denote it by fT .

Proof.— Suppose (φ_j) is a sequence of functions in $\mathcal{D}(I)$, that converges to 0 in the $\mathcal{D}(I)$ -sense. There is a segment $[a, b] \subset\subset I$ such that $\text{supp } \varphi_j \subset [a, b]$ for all j , which implies $\text{supp } f\varphi_j \subset [a, b]$ for all j . Moreover, for any $\alpha \in \mathbb{N}$,

$$(f\varphi)^{(\alpha)} = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} f^{(\beta)} \varphi^{(\alpha-\beta)}.$$

so that if we denote

$$M = \max_{\beta \leq \alpha} \sup_{x \in [a, b]} |f^{(\beta)}(x)| := \|f\|_{C^\alpha([a, b])},$$

we see that

$$\|(f\varphi_j)^{(\alpha)}\|_\infty \leq M \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \|\varphi_j^{(\alpha-\beta)}\|_\infty.$$

Since each of the terms in the sum converges to 0 as $j \rightarrow +\infty$, the right hand side does so too, so that $(f\varphi_j) \rightarrow 0$ in $\mathcal{D}(I)$. Finally, since T is a distribution, $\langle T, f\varphi_j \rangle \rightarrow 0$. This proves that fT is continuous on $\mathcal{D}(I)$, hence is a distribution on I . \square

Exercise 1.5.2 For $f \in \mathcal{C}^\infty(\mathbb{R}^d)$, show that $f\delta_0 = f(0)\delta_0$.

Exercise 1.5.3 For $f \in \mathcal{C}^\infty(\mathbb{R}^d)$ and $g \in L^1_{\text{loc}}(\mathbb{R}^d)$, show that $fT_g = T_{fg}$.

Exercise 1.5.4 Show that $x \text{pv}(1/x) = 1$.

Exercise 1.5.5 For $f, g \in \mathcal{C}^\infty(\mathbb{R}^d)$, show that $f(gT) = (fg)T$.

We have seen that distributions can be differentiated arbitrarily many times; so can smooth functions. In view of the above proposition, it makes sense to differentiate the product fT . Like for the product of two smooth functions, the derivatives of this product satisfy a Leibniz formula (we leave the proof of the following result to the reader).

Proposition 1.5.6 (Leibniz formula) For $f \in \mathcal{C}^\infty(I)$, and $T \in \mathcal{D}'(I)$, we have

$$\forall \alpha \in \mathbb{N}, \quad (fT)^{(\alpha)} = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} f^{(\beta)} T^{(\alpha-\beta)}.$$

Exercise 1.5.2 implies the identity

$$(x - x_0)\delta_{x_0} = 0.$$

Our next statement is a converse to this identity.

Proposition 1.5.7 Let $x_0 \in I$, and let $T \in \mathcal{D}'(I)$ satisfy

$$(x - x_0)T = 0.$$

Then there exists a constant $c \in \mathbb{C}$ such that $T = c\delta_{x_0}$.

Proof.— Let $\chi \in \mathcal{D}(I)$ satisfy $\chi(x_0) = 1$; as in previous proofs, we use it as a “reference function”. For every $\varphi \in \mathcal{D}(I)$, we split it into a multiple of this reference function, and a term vanishing at x_0 , using Hadamard’s lemma (see exercise 1.1.1):

$$\begin{aligned} \varphi(x) &= \varphi(x_0) + (x - x_0)\psi(x), \quad \psi \in \mathcal{C}^\infty(I), \\ &= \varphi(x_0)\chi(x) + (x - x_0) \left(\frac{1 - \chi(x)}{x - x_0} \varphi(x_0) + \psi(x) \right) \\ &= \varphi(x_0)\chi(x) + (x - x_0)\theta(x). \end{aligned}$$

The function θ is smooth from its explicit expression, and is compactly supported since $\varphi - \varphi(x_0)\chi$ is so. We infer

$$\langle T, \varphi \rangle = \langle T, \chi \rangle \varphi(x_0) + \langle (x - x_0)T, \theta \rangle = \langle T, \chi \rangle \varphi(x_0),$$

whence $T = c\delta_{x_0}$ with $c := \langle T, \chi \rangle$. □

Example 1.5.8 We are now in measure to solve another linear first order differential equation on the space of distributions, more complicated than the equation $T' = S$ of the previous section. Let us look for solutions $T \in \mathcal{D}'(\mathbb{R})$ of the equation

$$(1.5.5) \quad xT' + T = 0.$$

By the Leibniz formula, this equation is equivalent to

$$(xT)' = 0.$$

Using Proposition 1.4.8, this is equivalent to

$$xT = T_{c_1} \quad \text{for some } c_1 \in \mathbb{C}.$$

Exercise 1.5.4 shows that $c_1 \text{pv}(1/x)$ is a solution of this equation, so we now have to solve

$$x \left(T - c_1 \text{pv} \left(\frac{1}{x} \right) \right) = 0 .$$

Finally, using Proposition 1.5.7, we obtain the general solution of the differential equation $xT' + T = 0$:

$$T = c_1 \text{pv} \left(\frac{1}{x} \right) + c_2 \delta_0 , \quad (c_1, c_2) \in \mathbb{C}^2 .$$

We have proved that the only global nontrivial solutions to (1.5.5) are nonsmooth distributions; the singularity is due to the fact that the coefficient of the highest derivative (here T') vanishes at a point of the interval.

Finally, we use elements of the proof of Prop. 1.4.10 to prove the surjectivity of the mapping $T \mapsto (x - x_0)T$ on $\mathcal{D}'(I)$.

Proposition 1.5.9 Given $S \in \mathcal{D}'(I)$ and $x_0 \in I$, there exists $T \in \mathcal{D}'(I)$ such that

$$(x - x_0)T = S .$$

Of course, if $x_0 \notin I$, the statement trivially holds, since the function $(x - x_0)^{-1}$ is smooth on I , and $(x - x_0)^{-1}S$ is well-defined.

Proof.— We proceed similarly to the proof of Proposition 1.4.10. Indeed, $(x - x_0)T = S$ reads

$$(1.5.6) \quad \forall \varphi \in \mathcal{D}(I) , \quad \langle T, (x - x_0)\varphi \rangle = \langle S, \varphi \rangle ,$$

which, in view of Hadamard's lemma, imposes the value of T on test functions vanishing at x_0 . As in the proof of Proposition 1.5.7, we fix a reference function $\chi \in \mathcal{D}(I)$ such that $\chi(x_0) = 1$. For every $\psi \in \mathcal{D}(I)$, we define

$$Q(\psi)(x) = \frac{\psi(x) - \psi(x_0)\chi(x)}{x - x_0} = \int_0^1 \left[\psi'(tx + (1-t)x_0) - \psi(x_0)\chi'((tx + (1-t)x_0)) \right] dt .$$

This defines a linear mapping $Q : \mathcal{D}(I) \rightarrow \mathcal{D}(I)$, with the properties

$$\text{supp } Q(\psi) \subset \text{supp}(\psi) \cup \text{supp}(\chi) ,$$

and, if (ψ_n) converges to ψ in $\mathcal{D}(I)$, then $Q(\psi_n)$ converges to $Q(\psi)$ in $\mathcal{D}(I)$. We can then define $T \in \mathcal{D}'(I)$ by

$$\forall \psi \in \mathcal{D}(I) , \quad \langle T, \psi \rangle = \langle S, Q(\psi) \rangle .$$

Since, for every $\varphi \in \mathcal{D}(I)$,

$$Q((x - x_0)\varphi) = \varphi ,$$

this distribution T satisfies (1.5.6). □

Similarly, the differential equation of Example 1.5.8 can also be made nonhomogeneous.

Exercise 1.5.10 For a given $S \in \mathcal{D}'(I)$, describe the solutions $T \in \mathcal{D}'(I)$ of the equation

$$xT' + T = S .$$

To generalize Proposition 1.5.9, we may as well multiply T' by higher order monomials.

Corollary 1.5.11 If $x_0 \in I$ and m is a positive integer, the mapping $T \in \mathcal{D}'(I) \mapsto (x - x_0)^m T \in \mathcal{D}'(I)$ is surjective, and its kernel is the m -dimensional vector space generated by $\{\delta_{x_0}^{(j)}, j = 0, 1, \dots, m - 1\}$.

Proof.— We proceed by induction on m . The case $m = 1$ has been solved in Propositions 1.5.7 and 1.5.9. Assume $m \geq 2$ and that the statement is true for $m - 1$. The surjectivity of the mapping $T \in \mathcal{D}'(I) \mapsto (x - x_0)^m T \in \mathcal{D}'(I)$ follows immediately from the similar property at the level $m - 1$ and from Proposition 1.5.9.

Concerning the kernel, the identity

$$(x - x_0)^m T = (x - x_0)^{m-1}((x - x_0)T)$$

is solved by the induction hypothesis by

$$(x - x_0)T = \sum_{j=0}^{m-2} c_j \delta_{x_0}^{(j)}, \quad \text{with arbitrary coefficients } c_j \in \mathbb{C}.$$

Prop. 1.5.9 claims that this has a solution. We can actually find an explicit solution to this equation. For this, we will use the following lemma, which generalizes Prop. 1.5.7:

Lemma 1.5.12 For any point $x_0 \in I$ and any integer $k \geq 1$,

$$(x - x_0)\delta_{x_0}^{(k)} = -k\delta_{x_0}^{(k-1)}.$$

Indeed, for every test function φ , we have

$$\langle (x - x_0)\delta_{x_0}^{(k)}, \varphi \rangle = (-1)^k \langle (x - x_0)\varphi \rangle^{(k)}(x_0) = (-1)^k k \varphi^{(k-1)}(x_0) = -k \langle \delta_{x_0}^{(k-1)}, \varphi \rangle.$$

Coming back to the proof of Prop. 1.5.11, we may rewrite

$$(x - x_0)T = -(x - x_0) \sum_{j=0}^{m-2} \frac{c_j}{j+1} \delta_{x_0}^{(j+1)},$$

Finally, applying Prop. 1.5.7 to the distribution

$$T + \sum_{j=0}^{m-2} \frac{c_j}{j+1} \delta_{x_0}^{(j+1)}$$

adds a possible term $c\delta_{x_0}$, which completes the proof. \square

After multiplying T by monomials, we may generalize to products with polynomials (we leave the proof to the reader).

Corollary 1.5.13 Let P be a non identically zero polynomial function. Then the mapping

$$T \in \mathcal{D}'(I) \mapsto PT \in \mathcal{D}'(I)$$

is surjective, and its kernel is the vector space generated by the distributions $\delta_a^{(j)}$, where a ranges over the zeroes of P inside I , and $j = 0, 1, \dots, m(a) - 1$, where $m(a)$ denotes the multiplicity of a as a zero of P .

Finally, let us solve another (very simple) differential equation on $\mathcal{D}'(\mathbb{R})$.

Proposition 1.5.14 Let $I \subset \mathbb{R}$ be an open interval, and $a \in C^\infty(I)$. The distributions in $\mathcal{D}'(I)$ that satisfy the differential equation

$$T' + aT = 0$$

are exactly the C^∞ solutions, that is the regular distributions T_f with $f : x \mapsto Ce^{-A(x)}$, for some constant $C \in \mathbb{C}$, where A is primitive of a in I .

In other words, the distribution solutions of this differential equation coincide with the classical solutions. Notice that this is not always the case if the coefficient of the highest derivative vanishes — see example 1.5.8 above.

Proof.— Let A be a primitive of a on I . For $T \in \mathcal{D}'(I)$, we have, using Leibniz formula,

$$(e^AT)' = ae^AT + e^AT' = e^A(T' + aT).$$

Thus

$$T' + aT = 0 \iff (e^AT)' = 0 \iff e^AT = T_C \iff T = e^{-A}T_C = T_{Ce^{-A}}.$$

□

Exercise 1.5.15 Solve in $\mathcal{D}'(I)$ the inhomogeneous equation $T' + aT = f$, for $f \in L^1_{\text{loc}}(I)$. Furthermore, prove that, if $S \in \mathcal{D}'(I)$, there exists $T \in \mathcal{D}'(I)$ such that

$$T' + aT = S.$$

1.6 Restriction and support

1.6.1 Definitions

Definition 1.6.1 Let $T \in \mathcal{D}'(I)$, and $J \subset I$ an open subinterval of I . The restriction of T to J is the distribution $T|_J \in \mathcal{D}'(J)$ defined as

$$\forall \varphi \in \mathcal{D}(J), \langle T|_J, \varphi \rangle = \langle T, \underline{\varphi} \rangle,$$

where $\underline{\varphi}$ denotes the extension of φ by 0 on $I \setminus J$.

We say that T vanishes in J if $T|_J = 0$.

Let us state without proof the following elementary properties of restriction with respect to derivation and multiplication by a smooth function.

$$(T|_J)' = T'|_J, (fT)|_J = f|_J T|_J.$$

As a first application of this important notion, let us state a very useful result.

Proposition 1.6.2 Let $(I_n)_{n \in \mathbb{N}}$ be a sequence of intervals in I such that $I_n =]a_n, b_n[$, $[a_n, b_n] \subset I_{n+1}$ for every n , and

$$\bigcup_{n \in \mathbb{N}} I_n = I .$$

Suppose we are given for every $n \in \mathbb{N}$, a distribution T_n on I_n such that

$$\forall n \in \mathbb{N} , T_{n+1}|_{I_n} = T_n .$$

Then there exists a unique distribution T on I such that

$$\forall n \in \mathbb{N} , T|_{I_n} = T_n .$$

Proof.— The key of the proof is the following elementary fact. For every segment $[a, b]$ included in I , there exists n such that $[a, b] \subset I_n$. Indeed, from the assumptions, the sequence (a_n) is decreasing, tending to the left boundary of I , while the sequence (b_n) is increasing, tending to the right boundary of I . This fact is also a consequence of the Borel–Lebesgue property for the compact set $[a, b]$.

If such a distribution T exists, for every $\varphi \in \mathcal{D}(I)$, there exists n such that $\text{supp}(\varphi) \subset I_n$, and therefore we should have

$$\langle T, \varphi \rangle = \langle T_n, \varphi \rangle .$$

This proves the uniqueness of T , and suggests a way of constructing it. Indeed, given $\varphi \in \mathcal{D}(I)$, the assumption $T_{n+1}|_{I_n} = T_n$ implies that, for every n such that $\text{supp}(\varphi) \subset I_n$, the quantity $\langle T_n, \varphi \rangle$ does not depend on n . We can therefore define

$$\langle T, \varphi \rangle = \langle T_n, \varphi \rangle$$

for every such n . This clearly defines a linear form on $\mathcal{D}(I)$. Furthermore, if φ_j converges to φ in $\mathcal{D}(I)$, then there exist n such that,

$$\forall j , \text{supp}(\varphi_j) \subset I_n .$$

Therefore

$$\langle T, \varphi_j \rangle = \langle T_n, \varphi_j \rangle$$

and, because T_n is a distribution on I_n , $\langle T, \varphi_j \rangle$ tends to $\langle T, \varphi \rangle$. □

Let us state a first consequence of Proposition 1.6.2.

Corollary 1.6.3 Let $f \in C^\infty(I)$ admitting only finite order zeroes in I , namely, if $f(a) = 0$, there exists $m \geq 1$ such that $f^{(m)}(a) \neq 0$. Then the mapping $T \in \mathcal{D}'(I) \mapsto fT \in \mathcal{D}'(I)$ is surjective.

This result generalizes the product with polynomials of Corollary 1.5.13, to the case of a smooth function f with possibly infinitely many zeros on I ; yet, a crucial point is the fact that on any segment $[a, b] \subset I$, the number of zeros of f is at most finite.

Proof.— If $f(a) = 0$ and m is the smallest integer such that $f^{(m)}(a) \neq 0$, the Taylor formula yields

$$f(x) = \frac{(x-a)^m}{(m-1)!} \int_0^1 (1-t)^{m-1} f^{(m)}(a+t(x-a)) dt .$$

From the continuity of $f^{(m)}$, we infer that there exists an open interval $J_a \subset I$ containing a such that $f(x) \neq 0$ for every $x \in J_a \setminus \{a\}$. If $[\alpha, \beta]$ is a segment of I , we conclude that the compact set of zeroes of f in $[\alpha, \beta]$ is covered by the family of open intervals J_a 's, hence – from the Borel–Lebesgue property – by a finite subfamily of J_a 's, hence this set is finite. Consequently, in $] \alpha, \beta [$, we can write

$$f = Pg ,$$

where P is a polynomial and g is a smooth function which does not have any zero in $] \alpha, \beta [$. Consequently, the Corollary 1.5.13 shows that the mapping

$$T \in \mathcal{D}'(] \alpha, \beta [) \mapsto fT \in \mathcal{D}'(] \alpha, \beta [)$$

is surjective, and its kernel consists of finite linear combinations of Dirac masses on the zeroes of f in $] \alpha, \beta [$ and of some of their derivatives. In particular, any distribution $U_{] \alpha, \beta [}$ in this kernel is the restriction to $] \alpha, \beta [$ of some distribution U on I , satisfying $fU = 0$ on I .

Let us now construct a sequence $(I_n)_{n \in \mathbb{N}}$ as in Proposition 1.6.2. Let $S \in \mathcal{D}'(I)$. In view of the above observation, there exists $T_0 \in \mathcal{D}'(I_0)$ such that $f|_{I_0} T_0 = S|_{I_0}$, and there exists $\tilde{T}_1 \in \mathcal{D}'(I_1)$ such that $f|_{I_1} \tilde{T}_1 = S|_{I_1}$. Denote by $\tilde{T}_{1,0}$ the restriction of \tilde{T}_1 to I_0 . Of course,

$$f|_{I_0}(\tilde{T}_{1,0} - T_0) = 0 ,$$

and from the above observation, there exists $U_1 \in \mathcal{D}'(I_1)$ such that $U_1|_{I_0} = \tilde{T}_{1,0} - T_0$ and $f|_{I_1} U_1 = 0$. If we set

$$T_1 = \tilde{T}_1 - U_1 \in \mathcal{D}'(I_1) ,$$

we observe that

$$T_1|_{I_0} = T_0 , f|_{I_1} T_1 = S|_{I_1} .$$

By induction, we can therefore construct, for every $n \in \mathbb{N}$, a distribution T_n on I_n such that

$$T_{n+1}|_{I_n} = T_n , f|_{I_n} T_n = S|_{I_n} .$$

Applying Proposition 1.6.2, there exists $T \in \mathcal{D}'(I)$ such that, for every n , $T|_{I_n} = T_n$. Consequently,

$$\forall n \in \mathbb{N} , (fT)|_{I_n} = f|_{I_n} T_n = S|_{I_n} ,$$

hence, by the uniqueness part of Proposition 1.6.2, we conclude that

$$fT = S .$$

□

We now introduce a crucial notion.

Definition 1.6.4 The support of a distribution $T \in \mathcal{D}'(I)$ is the complement of the union of all the open subintervals on which the restriction of T is 0. We denote it by $\text{supp } T$.

Notice that $\text{supp } T$ is closed, and the following characterizations are convenient.

- $x_0 \notin \text{supp } T$ if and only if there is an open neighborhood J of x_0 such that $T|_J = 0$.
- $x_0 \in \text{supp } T$ if and only if for any open neighborhood J of x_0 , one can find $\varphi \in \mathcal{C}_0^\infty(J)$ such that $\langle T, \varphi \rangle \neq 0$.

Example 1.6.5 *i)* Let $T = \delta_0$. If J is an open interval that does not contain $\{0\}$, then $\langle T, \varphi \rangle = \varphi(0) = 0$ for any $\varphi \in \mathcal{D}(\mathbb{R})$ such that $\text{supp } \varphi \subset J$. Thus $\text{supp } T \subset \{0\}$. On the other hand, if J is an open subinterval that contains 0, we can find a plateau function ψ over $[-r, r]$ in $\mathcal{D}(J)$, and for this function we have $\langle T, \psi \rangle = \psi(0) = 1$. Therefore we have $0 \in \text{supp } T$ and finally $\text{supp } T = \{0\}$.

ii) If $T = T_f$ for some $f \in \mathcal{C}^0(I)$, with I an open interval of \mathbb{R} , we have

$$\text{supp } T = \text{supp } f = \overline{\{x \in I, f(x) \neq 0\}}.$$

Indeed, suppose that $x_0 \notin \text{supp } f$. There is an open neighborhood J of x_0 such that $f|_J = 0$. For $\varphi \in \mathcal{D}(J)$, we have thus $\langle T_f, \varphi \rangle = 0$, so that T_f vanishes on J , and $x_0 \notin \text{supp } T_f$. Conversely, if $x_0 \notin \text{supp } T_f$, there is a neighborhood J of x_0 such that, for all $\varphi \in \mathcal{C}_0^\infty(J)$, we have $\int f\varphi dx = \langle T_f, \varphi \rangle = 0$. We have seen in Proposition 1.3.4 that this implies $f = 0$ in J , thus $x_0 \notin \text{supp } f$.

iii) If $T = T_f$ for some $f \in L^1_{\text{loc}}(I)$, the same argument leads to

$$\text{supp}(T_f) = \text{ess - supp}(f) = \{x_0 \in I, \forall r > 0, \text{meas}\{x \in [x_0 - r, x_0 + r], f(x) \neq 0\} > 0\}.$$

1.6.2 Some properties of the support

Lemma 1.6.6 Let $I \subset \mathbb{R}$ be an open set, and $T \in \mathcal{D}'(I)$.

- i)* For $k \in \mathbb{N}$, $\text{supp } T^{(k)} \subset \text{supp } T$.
- ii)* For $f \in \mathcal{C}^\infty(I)$, $\text{supp } fT \subset \text{supp } f \cap \text{supp } T$.

Proof.— Let $x_0 \notin \text{supp } T$. There exists a neighborhood V of x_0 such that for all $\psi \in \mathcal{C}_0^\infty(V)$, $\langle T, \psi \rangle = 0$. But if $\varphi \in \mathcal{C}_0^\infty(V)$, $\psi = \varphi^{(k)} \in \mathcal{C}_0^\infty(V)$, thus

$$\langle T^{(k)}, \varphi \rangle = (-1)^k \langle T, \varphi^{(k)} \rangle = 0.$$

Therefore $x_0 \notin \text{supp } T^{(k)}$, which proves *i)*. Now we prove the second point. If $x_0 \notin \text{supp } f \cap \text{supp } T$, x_0 either belongs to $(\text{supp } f)^c$ or to $(\text{supp } T)^c$. In the first case, there exists a neighborhood V of x_0 such that $f|_V = 0$. For $\varphi \in \mathcal{C}_0^\infty(V)$, we have $\langle fT, \varphi \rangle = \langle T, f\varphi \rangle = 0$, thus $x_0 \notin \text{supp}(fT)$. In the latter case, there exists a neighborhood V of x_0 such that, for all $\psi \in \mathcal{C}_0^\infty(V)$, $\langle T, \psi \rangle = 0$. For $\varphi \in \mathcal{C}_0^\infty(V)$, $f\varphi$ is a smooth function, which vanishes out of a compact set included in V . Thus $\langle fT, \varphi \rangle = \langle T, f\varphi \rangle = 0$, and $x_0 \notin \text{supp } fT$. \square

The following result is fundamental.

Proposition 1.6.7 Let $\varphi \in \mathcal{D}(I)$ and $T \in \mathcal{D}'(I)$. If $\text{supp } \varphi \cap \text{supp } T = \emptyset$, then $\langle T, \varphi \rangle = 0$.

Proof.— Let $x \in \text{supp } \varphi$. We have, by assumption, $x \notin \text{supp } T$, thus there is an open subinterval J_x containing x on which T vanishes. From the covering of the compact subset $\text{supp}(\varphi)$ with the open sets J_x , one can extract a finite covering

$$\text{supp } \varphi \subset \bigcup_{j=1}^n J_{x_j}.$$

By Proposition 1.2.7, one can find a smooth partition of φ , namely functions $\varphi_j \in \mathcal{D}(J_{x_j})$ for $j = 1, \dots, n$, such that

$$\varphi = \varphi_1 + \dots + \varphi_n.$$

Therefore

$$\langle T, \varphi \rangle = \sum_{j=1}^n \langle T, \varphi_j \rangle = 0.$$

□

A word of caution: one may have $\varphi = 0$ on $\text{supp } T$ and $\langle T, \varphi \rangle \neq 0$. For example, this is the case for $T = \delta'_0$ and $\varphi \in \mathcal{D}(\mathbb{R})$ such that $\varphi(0) = 0$, $\varphi'(0) = 1$.

An immediate consequence of Proposition 1.6.7 is the following important

Corollary 1.6.8 The only distribution with empty support is 0.

We conclude this section by the important characterisation of distributions $T \in \mathcal{D}'(I)$ supported on a single point.

Proposition 1.6.9 Let $T \in \mathcal{D}'(I)$, with $T \neq 0$, and $x_0 \in I$. If $\text{supp } T \subset \{x_0\}$, there exists $m \in \mathbb{N}$ and $m + 1$ complex numbers a_k for $0 \leq k \leq m$ such that

$$T = \sum_{k=0}^m a_k \delta_{x_0}^{(k)}.$$

Proof.— We write the proof for $x_0 = 0$. Let $[a, b]$ be a segment of I containing 0 in its interior, and $\chi \in \mathcal{D}([a, b])$ a plateau function equal to unity on a smaller segment containing 0 in its interior. Since T is a distribution, there exist $C > 0$, $m \in \mathbb{N}$ such that

$$\forall \psi \in \mathcal{D}(I), \quad \text{supp } \psi \subset [a, b] \implies |\langle T, \psi \rangle| \leq C \sum_{k=0}^m \sup |\psi^{(k)}|,$$

and from now on we denote by m the smallest integer for which this property holds⁶. For any function $\varphi \in \mathcal{D}(I)$, since $\text{supp}(\varphi - \chi\varphi) \cap \text{supp } T = \emptyset$, we have

$$\langle T, \varphi \rangle = \langle T, \chi\varphi \rangle + \langle T, \varphi - \chi\varphi \rangle = \langle T, \chi\varphi \rangle.$$

Since $\text{supp}(\chi\varphi) \subset [a, b]$, we have

$$|\langle T, \varphi \rangle| \leq |\langle T, \chi\varphi \rangle| \leq C \sum_{k=0}^m \sup |(\chi\varphi)^{(k)}|.$$

6. This value m is called the *order* of the distribution $T|_{[a,b]}$

At last, the Leibniz formula gives

$$\sup |(\chi\varphi)^{(k)}| = \sup_{[a,b]} |(\chi\varphi)^{(k)}| \leq C \sum_{\ell=0}^k \sup_{[a,b]} |\varphi^{(\ell)}|,$$

We first claim that, if $\varphi \in \mathcal{D}(I)$ satisfies $\varphi^{(k)}(0) = 0$ for every $k \leq m$, then $\langle T, \varphi \rangle = 0$. Indeed, let χ be a plateau function on $[-\frac{1}{2}, \frac{1}{2}]$, with $\text{supp}(\chi) \subset [-1, 1]$. For $\varepsilon > 0$ small enough, set

$$\chi_\varepsilon(x) = \chi\left(\frac{x}{\varepsilon}\right).$$

Since $\varphi - \chi_\varepsilon\varphi$ vanishes in a neighborhood of 0, we have

$$\langle T, \varphi \rangle = \langle T, \chi_\varepsilon\varphi \rangle.$$

By the Taylor formula and the assumption on φ

$$\varphi(x) = x^{m+1}\psi(x)$$

for some $\psi \in \mathcal{D}(I)$. Hence

$$\chi_\varepsilon\varphi(x) = \chi\left(\frac{x}{\varepsilon}\right)x^{m+1}\psi(x) = \varepsilon^{m+1}\rho\left(\frac{x}{\varepsilon}\right)\psi(x),$$

where we introduced the notation $\rho(y) = y^{m+1}\chi(y)$. By the Leibniz formula, we have

$$\sup_{k \leq m} \sup_x \left| \left(\rho\left(\frac{x}{\varepsilon}\right)\psi(x) \right)^{(k)} \right| \leq \frac{B}{\varepsilon^m}$$

and therefore

$$|\langle T, \varphi \rangle| = |\langle T, \chi_\varepsilon\varphi \rangle| \leq C \sup_{k \leq m} \sup |(\chi_\varepsilon\varphi)^{(k)}| \leq CB\varepsilon.$$

As ε tends to 0, we obtain the claimed result

$$\langle T, \varphi \rangle = 0.$$

Let us now study the case of a general $\varphi \in \mathcal{D}(I)$. Applying the Taylor formula, we have

$$\varphi(x) = \sum_{k \leq m} \frac{x^k}{k!} \varphi^{(k)}(0) \chi_{\varepsilon_0}(x) + r(x),$$

where $\varepsilon_0 > 0$ is small enough so that $[-\varepsilon_0, \varepsilon_0] \subset I$, and $r \in \mathcal{C}_0^\infty(I)$ satisfies

$$r^{(k)}(0) = 0, \text{ for all } k \leq m.$$

We can therefore apply the above statement to r , and get

$$\langle T, \varphi \rangle = \sum_{k \leq m} \varphi^{(k)}(0) \langle T, \frac{x^k}{k!} \chi_{\varepsilon_0}(x) \rangle.$$

This is precisely what we have claimed, if we set $a_k = \langle T, \frac{x^k}{k!} \chi_{\varepsilon_0} \rangle$. □

1.7 Sequences of distributions

1.7.1 Convergence in \mathcal{D}'

Definition 1.7.1 Let (T_j) be a sequence of distributions in $\mathcal{D}'(I)$. We say that (T_j) converges to $T \in \mathcal{D}'(I)$ when, for any function $\varphi \in \mathcal{C}_0^\infty(I)$, the sequence of complex numbers $(\langle T_j, \varphi \rangle)$ converges to $\langle T, \varphi \rangle$. In this case, we write $T_j \rightarrow T$ in $\mathcal{D}'(I)$.

Example 1.7.2 If (f_j) is a sequence of locally integrable functions on I such that, for some $f \in L^1_{\text{loc}}(I)$,

$$\forall [a, b] \subset I, \int_a^b |f_j(x) - f(x)| dx \rightarrow 0,$$

then

$$T_{f_j} \rightarrow T_f$$

in $\mathcal{D}'(I)$. Indeed, for every test function φ on I with $\text{supp } \varphi \subset [a, b]$,

$$\langle T_{f_j}, \varphi \rangle - \langle T_f, \varphi \rangle = \int_I f_j(x) \varphi(x) dx - \int_I f(x) \varphi(x) dx$$

so that

$$|\langle T_{f_j}, \varphi \rangle - \langle T_f, \varphi \rangle| \leq \sup |\varphi(x)| \int_a^b |f_j(x) - f(x)| dx$$

which tends to 0 as j tends to the infinity.

Notice that a special case of this is the following situation. The sequence f_j converges to f almost everywhere on I , and there exists a locally integrable function h on I such that $\sup_j |f_j| \leq h$ almost everywhere on I . Indeed, the connection to the above condition of L^1 convergence is provided by the dominated convergence theorem.

Example 1.7.3 Let $\rho \in L^1(\mathbb{R})$ such that $\int_{\mathbb{R}} \rho(x) dx = 1$, and (ρ_ε) the sequence of functions defined by

$$\rho_\varepsilon(x) = \varepsilon^{-1} \rho\left(\frac{x}{\varepsilon}\right).$$

We also denote $(T_\varepsilon) = (T_{\rho_\varepsilon})$ the associated family of distributions. For $\varphi \in \mathcal{C}_0^\infty(\mathbb{R})$, we have

$$\langle T_\varepsilon, \varphi \rangle = \int_{\mathbb{R}} \rho_\varepsilon(x) \varphi(x) dx = \int_{\mathbb{R}} \rho(y) \varphi(\varepsilon y) dy.$$

By the dominated convergence theorem, $\langle T_\varepsilon, \varphi \rangle \rightarrow \varphi(0)$ as $\varepsilon \rightarrow 0$. Therefore the sequence (T_{ρ_ε}) converges to δ_0 in $\mathcal{D}'(\mathbb{R})$.

Exercise 1.7.4 Let (f_j) be a sequence of L^1 functions on I such that

$$\sup_j \int_I |f_j| dx < \infty, \int_I f_j dx \rightarrow c, \text{supp}(T_{f_j}) \subset [x_0 - \varepsilon_j, x_0 + \varepsilon_j], \varepsilon_j \rightarrow 0.$$

Prove that

$$T_{f_j} \rightarrow c \delta_{x_0}.$$

Exercise 1.7.5 Let

$$f_\varepsilon(x) = \begin{cases} \frac{1}{\varepsilon^2} & \text{if } x \in]0, \varepsilon[\\ \frac{-1}{\varepsilon^2} & \text{if } x \in]-\varepsilon, 0[\\ 0 & \text{if } |x| > \varepsilon \end{cases}$$

Prove that

$$T_{f_\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} -\delta'_0.$$

Exercise 1.7.6 Let

$$f_\varepsilon(x) = \frac{x}{x^2 + \varepsilon}, \quad \varepsilon > 0.$$

Prove that

$$T_{f_\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} \text{pv} \left(\frac{1}{x} \right)$$

in $\mathcal{D}'(\mathbb{R})$.

The operations that we have defined on distributions are continuous with respect to the notion of convergence. More precisely,

Proposition 1.7.7 If (T_j) converges to T in $\mathcal{D}'(I)$, then

- i) For any $k \in \mathbb{N}$, $(T_j^{(k)})$ converges to $T^{(k)}$.
- ii) For any $f \in \mathcal{C}^\infty(I)$, (fT_j) converges to fT .
- iii) For any open subinterval $J \subset I$, $(T_j)|_J$ converges to $T|_J$.

Proof.— Let $\varphi \in \mathcal{C}_0^\infty(I)$. We have clearly

$$\langle \partial^\alpha T_j, \varphi \rangle = (-1)^{|\alpha|} \langle T_j, \partial^\alpha \varphi \rangle \rightarrow (-1)^{|\alpha|} \langle T, \partial^\alpha \varphi \rangle = \langle \partial^\alpha T, \varphi \rangle,$$

and

$$\langle fT_j, \varphi \rangle = \langle T_j, f\varphi \rangle \rightarrow \langle T, f\varphi \rangle = \langle fT, \varphi \rangle.$$

The last statement is trivial. □

Exercise 1.7.8 Show that if (f_j) converges to f in $\mathcal{C}^\infty(I)$, in the sense of the uniform convergence of all derivatives on every compact subset, then $(f_j T)$ converges to fT in $\mathcal{D}'(I)$.

1.7.2 The Uniform Boundedness Principle

Let us motivate the main statement of this paragraph by the following questions.

- Assume that, for a sequence (T_j) of distributions, and for every $\varphi \in \mathcal{D}(I)$, the sequence $(\langle T_j, \varphi \rangle)$ has a limit $L(\varphi)$. This clearly defines a linear form L on $\mathcal{D}(I)$. Is it true that L is a distribution ?

- Assume $T_j \rightarrow T$ in $\mathcal{D}'(I)$ and $f_j \rightarrow f$ in $\mathcal{C}^\infty(I)$ in the sense of the uniform convergence of all derivatives on every compact subset, does it imply that $f_j T_j \rightarrow fT$? In order to prove such a result, we write

$$\langle f_j T_j, \varphi \rangle - \langle fT, \varphi \rangle = \langle T_j, (f_j - f)\varphi \rangle + \langle T_j, f\varphi \rangle - \langle T, f\varphi \rangle.$$

The second term in the right hand side tends to 0 in view of the assumption. In the first term, we notice that $\psi_j = (f_j - f)\varphi$ converges to 0 in $\mathcal{D}(I)$, but it seems difficult to conclude that $\langle T_j, \psi_j \rangle \rightarrow 0$ without a bound on the action of T_j which is uniform with respect to j .

We state below, without proof, an important theoretical result, which is a “distributional version” of the Banach-Steinhaus theorem, and which provides such a uniform bound.

Proposition 1.7.9 Let (T_j) be a sequence in $\mathcal{D}'(I)$, and $[a, b] \subset I$ a segment. If, for any function $\varphi \in \mathcal{C}_0^\infty(I)$ with support in $[a, b]$,

$$\sup_j |\langle T_j, \varphi \rangle| < +\infty,$$

then there exists $C > 0$ and $m \in \mathbb{N}$, independent of j , such that

$$\forall \varphi \in \mathcal{C}_0^\infty(I), \text{ supp } \varphi \subset [a, b] \Rightarrow |\langle T_j, \varphi \rangle| \leq C \sum_{|\alpha| \leq m} \sup |\varphi^{(\alpha)}|.$$

As a consequence, we get the following somewhat surprising result.

Corollary 1.7.10 Let (T_j) be a sequence of distributions on I . If, for all functions $\varphi \in \mathcal{C}_0^\infty(I)$, the sequence $(\langle T_j, \varphi \rangle)$ converges in \mathbb{C} , then there exists a distribution $T \in \mathcal{D}'(I)$ such that $T_j \rightarrow T$ in $\mathcal{D}'(I)$.

Proof.— Let $T : \mathcal{C}_0^\infty(I) \rightarrow \mathbb{C}$ be the linear form given by

$$T(\varphi) = \lim_{j \rightarrow +\infty} \langle T_j, \varphi \rangle.$$

We want to show that T is a distribution. So pick a segment $[a, b] \subset I$. Proposition 1.7.9 ensures that there is a constant $C > 0$ and a natural number m such that, for any $\varphi \in \mathcal{C}_0^\infty(I)$ with $\text{supp } \varphi \subset [a, b]$, we have

$$|\langle T_j, \varphi \rangle| \leq C \sum_{|\alpha| \leq m} \sup |\varphi^{(\alpha)}|.$$

Then we can pass to the limit $j \rightarrow +\infty$, and we get the required estimate. \square

The second motivation can also be addressed successfully. The proof is left to the reader.

Corollary 1.7.11 Let (T_j) be a sequence of distributions on I which converges to T in $\mathcal{D}'(I)$. Then, for every sequence (ψ_j) which converges to ψ in $\mathcal{D}(I)$, we have $\langle T_j, \psi_j \rangle \rightarrow \langle T, \psi \rangle$. In particular, if $f_j \rightarrow f$ in $\mathcal{C}^\infty(I)$ in the sense of the uniform convergence of all derivatives on every compact subset, then $f_j T_j \rightarrow fT$.

A proof of the uniform boundedness principle as well as other applications will be given in the next chapter.

1.8 An introduction to Sobolev spaces

In this section, we give a short overview of some functional spaces, introduced by the Russian mathematician Sergei Sobolev (1908-1989), which allowed to solve differential equations in a wide context. Here we concentrate on a very special case in one variable. A more general presentation will be provided in the next chapters.

1.8.1 Generalised derivatives.

Definition 1.8.1 Let I be an open interval of \mathbb{R} , and let $u \in L^1_{\text{loc}}(I)$. We shall say that u admits a generalised derivative if there exists $f \in L^1_{\text{loc}}(I)$ such that, in the sense of distributions in I ,

$$T'_u = T_f .$$

The element $f \in L^1_{\text{loc}}(I)$ is then unique, is called the generalised derivative of u , and is denoted by $f = u'$.

Notice that uniqueness of f immediately follows from Proposition 1.3.4, and that the notation $f = u'$ is precisely made for leading to following generalised integration by parts formula, which is nothing but a reformulation of $T'_u = T_f$,

$$\forall \varphi \in C_0^\infty(I) , \int_I u(x)\varphi'(x) dx = - \int_I u'(x)\varphi(x) dx .$$

From Corollary 1.4.9, we infer the following useful statement.

Proposition 1.8.2 Assume $u \in L^1_{\text{loc}}(I)$ admits a generalised derivative $u' \in L^1_{\text{loc}}(I)$. Let $x_0 \in I$. Then there exists $c \in \mathbb{C}$ such that

$$u(x) = \int_{x_0}^x u'(t) dt + c \text{ a.e.}$$

In particular, u is almost everywhere equal to a continuous function.

Notice that, as a consequence of the Proposition 1.8.2, if the generalised derivative u' of u turns out to be a continuous function, then u is a C^1 function, and u' is merely its derivative in the usual sense.

1.8.2 The Sobolev space $H^1(I)$.

Definition 1.8.3 Let I be an open interval of \mathbb{R} . We denote by $H^1(I)$ the subspace of $u \in L^2(I)$ which admit a generalised derivative $u' \in L^2(I)$.

We endow the space $H^1(I)$ with the following inner product,

$$(1.8.7) \quad (u, v)_{H^1} = \int_I u'(x)\overline{v'(x)} dx + \int_I u(x)\overline{v(x)} dx, \quad u \in H^1(I), \quad v \in H^1(I).$$

The following proposition shows how the notion of convergence in $\mathcal{D}'(I)$ is useful in this context.

Proposition 1.8.4 Endowed with the inner product (1.8.7), the space $H^1(I)$ is a Hilbert space.

Proof.— Let (u_j) be a Cauchy sequence in $H^1(I)$ for the norm associated to (1.8.7). This precisely means that (u_j) and (u'_j) are Cauchy sequences of $L^2(I)$. Since $L^2(I)$ is a Hilbert space, there exist u, v in $L^2(I)$ such that

$$u_j \rightarrow u, \quad u'_j \rightarrow v$$

in $L^2(I)$. Since the convergence in $L^2(I)$ implies the convergence on $L^1(\cdot, a, b]$ for every segment $[a, b] \subset I$, we infer that, in $\mathcal{D}'(I)$,

$$T_{u_j} \rightarrow T_u, \quad T_{u'_j} \rightarrow T_v.$$

But, by definition of a generalised derivative, and using the continuity of derivation for the convergence of distributions,

$$T_{u'_j} = T'_{u_j} \rightarrow T'_u.$$

This implies $T'_u = T_v$, in other words u admits v as a generalised derivative, $u' = v \in L^2(I)$, which means that $u \in H^1(I)$ and

$$u_j \rightarrow u, \quad u'_j \rightarrow u'.$$

This precisely means that u_j tends to u in $H^1(I)$. Hence $H^1(I)$ is a complete normed vector space. \square

1.8.3 The special case of a bounded interval

In this section, $I =]a, b[$, $a, b \in \mathbb{R}$, is an open bounded interval. In this case, we get more information from Proposition 1.8.2 .

Proposition 1.8.5 If $u \in H^1(\cdot, a, b[)$, then u is almost everywhere equal to a continuous function on $[a, b]$, and we have the Sobolev inequality,

$$\|u\|_{L^\infty} \leq C \|u\|_{H^1},$$

where C is a constant depending only on $b - a$.

Proof.— We start from the identity in Proposition 1.8.2,

$$u(x) = \int_{x_0}^x u'(t) dt + c \quad a.e.$$

Since $L^2(\cdot, a, b[) \subset L^1(\cdot, a, b[)$, we infer that the right hand side has a limit as x tends to a and as x tends to b . Hence u is almost everywhere equal to a continuous function on $[a, b]$. Furthermore, by the Cauchy-Schwarz inequality,

$$\left| \int_y^x u'(t) dt \right| \leq \sqrt{b-a} \|u'\|_{L^2}.$$

Consider

$$m_u = \frac{1}{b-a} \int_a^b u(y) dy .$$

By the Cauchy–Schwarz inequality,

$$|m_u| \leq \frac{1}{\sqrt{b-a}} \|u\|_{L^2} .$$

On the other hand,

$$u(x) - m_u = \frac{1}{b-a} \int_a^b (u(x) - u(y)) dy = \frac{1}{b-a} \int_a^b \left(\int_y^x u'(t) dt \right) dy .$$

Therefore

$$|u(x) - m_u| \leq \frac{1}{b-a} \int_a^b \left| \int_y^x u'(t) dt \right| dy \leq \sqrt{b-a} \|u'\|_{L^2} .$$

Summing up, we obtain

$$\|u\|_{L^\infty} \leq \|u - m_u\|_{L^\infty} + |m_u| \leq (b-a)^{-1/2} \|u\|_{L^2} + (b-a)^{1/2} \|u'\|_{L^2} \leq C \|u\|_{H^1} .$$

□

Our next step is the introduction of an important closed subspace of $H^1(\]a, b[)$.

Definition 1.8.6 We denote by $H_0^1(\]a, b[)$ the closure of $C_0^\infty(\]a, b[)$ in $H^1(I)$.

Recall from Proposition 1.8.5 that $u \mapsto u(a)$ and $u \mapsto u(b)$ are continuous linear forms on $H^1(\]a, b[)$. A remarkable fact is that $H_0^1(\]a, b[)$ can be characterised by these linear forms.

Proposition 1.8.7 Given $u \in H^1(\]a, b[)$, u belongs to $H_0^1(\]a, b[)$ if and only if $u(a) = u(b) = 0$.

Proof.—Since elements of $H^1(\]a, b[)$ are continuous functions on $[a, b]$ and the H^1 norm controls the L^∞ norm on $[a, b]$ from Proposition 1.8.5, any element u of the H^1 closure of test functions can be uniformly approximated on $[a, b]$ by test functions. This immediately implies $u(a) = u(b) = 0$.

Conversely, consider $u \in H^1(\]a, b[)$ such that $u(a) = u(b) = 0$. For $\varepsilon > 0$ small enough, we consider a function $\chi_\varepsilon \in C_0^\infty(I)$, supported into $[a + \varepsilon, b - \varepsilon]$, and such that $\chi_\varepsilon = 1$ on $[a + 2\varepsilon, b - 2\varepsilon]$, with the estimate

$$\|\chi'_\varepsilon\|_{L^\infty} = O\left(\frac{1}{\varepsilon}\right) ,$$

as $\varepsilon \rightarrow 0$. An example of such a family of functions is

$$\chi_\varepsilon(x) = \chi\left(\frac{x-a}{\varepsilon}\right) \chi\left(\frac{b-x}{\varepsilon}\right) ,$$

where $\chi \in C^\infty(\mathbb{R})$ satisfies

$$\chi(t) = \begin{cases} 0 & \text{if } t \leq 1 \\ 1 & \text{if } t \geq 2. \end{cases}$$

Then we claim that $\chi_\varepsilon u \rightarrow u$ in $H^1(]a, b[)$ as $\varepsilon \rightarrow 0$. Indeed, $\chi_\varepsilon u \rightarrow u$ in $L^2(]a, b[)$ by dominated convergence, and

$$(\chi_\varepsilon u)' = \chi_\varepsilon u' + \chi'_\varepsilon u .$$

We are therefore reduced to proving that $\chi'_\varepsilon u \rightarrow 0$ in $L^2(]a, b[)$. We have

$$\|\chi'_\varepsilon u\|_{L^2} \leq O\left(\frac{1}{\varepsilon}\right) \left(\int_a^{a+2\varepsilon} |u(x)|^2 dx + \int_{b-2\varepsilon}^b |u(x)|^2 dx \right)^{\frac{1}{2}} ,$$

and, for $a \leq x \leq a + 2\varepsilon$,

$$\begin{aligned} |u(x)| &= \left| u(a) + \int_a^x u'(t) dt \right| = \left| \int_a^x u'(t) dt \right| \\ &\leq \sqrt{2\varepsilon} \left(\int_a^{a+2\varepsilon} |u'(x)|^2 dx \right)^{\frac{1}{2}} , \end{aligned}$$

with a similar inequality on $[b - 2\varepsilon, b]$. Finally,

$$\|\chi'_\varepsilon u\|_{L^2} \leq O(1) \left(\int_a^{a+2\varepsilon} |u'(x)|^2 dx + \int_{b-2\varepsilon}^b |u'(x)|^2 dx \right)^{\frac{1}{2}} \rightarrow 0 .$$

Summing up we have proved that u can be approximated in H^1 by compactly supported elements. The proof is completed by observing that any compactly supported element of $H^1(]a, b[)$ can be approximated by a sequence of $\mathcal{C}_0^\infty(]a, b[)$. This relies on the following regularisation argument. Let $v \in H^1(]a, b[)$ such that $\text{supp}(v) \subset [\alpha, \beta] \subset]a, b[$. We extend v by 0 as an element of $H^1(\mathbb{R})$ supported in $[\alpha, \beta]$. Let $\rho \in \mathcal{C}_0^\infty(\mathbb{R})$ of integral 1, and

$$\rho_\varepsilon(x) = \frac{1}{\varepsilon} \rho\left(\frac{x}{\varepsilon}\right) .$$

Then $\rho_\varepsilon * v \in \mathcal{C}_0^\infty(\mathbb{R})$, is compactly supported in $]a, b[$ if ε is small enough, and $\rho_\varepsilon * v \rightarrow v$ in L^2 as ε tends to 0. Furthermore, by the Leibniz rule,

$$(\rho_\varepsilon * v)'(x) = \int_{\mathbb{R}} \rho'_\varepsilon(x - y) v(y) dy = \int_{\mathbb{R}} -\frac{d}{dy} [\rho_\varepsilon(x - y)] v(y) dy = \int_{\mathbb{R}} \rho_\varepsilon(x - y) v'(y) dy ,$$

by definition of the generalised derivative. In other words,

$$(\rho_\varepsilon * v)' = \rho_\varepsilon * v'$$

and therefore $(\rho_\varepsilon * v)' \rightarrow v'$ in L^2 as ε tends to 0. Summing up, we have proved that $\rho_\varepsilon * v \rightarrow v$ in H^1 . \square

Proposition 1.8.7 implies an important inequality for elements of $H_0^1(]a, b[)$.

Proposition 1.8.8 (The Poincaré inequality) For every $u \in H^1(]a, b[)$ such that $u(a) = 0$ or $u(b) = 0$, we have

$$(1.8.8) \quad \|u\|_{L^2} \leq (b - a) \|u'\|_{L^2} .$$

In particular,

$$(u, v)_{H_0^1} = \int_a^b u'(x) \overline{v'(x)} dx$$

is an inner product on $H_0^1(]a, b[)$, which is equivalent to the H^1 inner product.

Proof.— Again we refer to Proposition 1.8.2,

$$u(x) = \int_{x_0}^x u'(t) dt + c .$$

Assume for instance $u(a) = 0$. Then making x tend to a , we infer

$$c = \int_a^{x_0} u'(t) dt$$

and consequently

$$u(x) = \int_a^x u'(t) dt .$$

Inequality (1.8.8) then follows from the Cauchy–Schwarz inequality. In view of Proposition 1.8.7, inequality (1.8.8) applies in particular to elements of $H_0^1(]a, b[)$, so that

$$\|u\|_{L^2}^2 \leq (b-a)^2 (u, u)_{H_0^1} .$$

Since $(u, u)_{H^1} = (u, u)_{H_0^1} + \|u\|_{L^2}^2$, we conclude that the inner products $(u, v)_{H^1}$ and $(u, v)_{H_0^1}$ are equivalent on H_0^1 . \square

The Hilbert structure on H_0^1 leads to a very efficient strategy for solving second order linear differential equations with homogeneous boundary conditions.

Theorem 1.8.9 Let $q \in L^1(]a, b[)$ such that $q \geq 0$, and let $f \in L^1(]a, b[)$. There exists a unique $u \in H^1(]a, b[)$ such that u' admits a generalised derivative u'' satisfying

$$-u'' + qu = f , \quad u(a) = u(b) = 0 .$$

Proof.— Notice that qu is well defined in L^1 since $q \in L^1$ and $u \in L^\infty$. Decomposing f into its real and imaginary parts, we may assume f is real valued, so that u is to be real valued as well. So we shall work in the real Hilbert space made of real valued elements of H^1 . This simple reduction allows to avoid the complex conjugation in the inner product, so that the connection with the distribution bracket is clearer.

In view of Proposition 1.8.7, the above problem is equivalent to

$$-T_{u'} + T_{qu} = T_f , \quad u \in H_0^1(]a, b[) ,$$

or, for every $\varphi \in \mathcal{C}_0^\infty(]a, b[)$,

$$\int_a^b u' \varphi' dx + \int_a^b qu \varphi dx = \int_a^b f \varphi dx , \quad u \in H_0^1(]a, b[) .$$

Since the left hand side and the right hand side of the above identity are continuous linear forms of φ for the H^1 norm, and since H_0^1 is the closure of \mathcal{C}_0^∞ in H^1 , we infer that the problem is equivalent to finding $u \in H_0^1(]a, b[)$ such that

$$\forall v \in H_0^1(]a, b[) , \quad \int_a^b u' v' dx + \int_a^b quv dx = \int_a^b f v dx .$$

Now observe that the left hand side is an inner product on the real space H_0^1 , which is equivalent to the H^1 inner product. Indeed,

$$\begin{aligned} (u, u)_{H_0^1} &\leq \int_a^b (u')^2 dx + \int_a^b q u^2 dx \leq \int_a^b (u')^2 dx + \left(\int_a^b q dx \right) \|u\|_{L^\infty}^2 \\ &\leq \max \left(1, C^2 \left(\int_a^b q dx \right) \right) (u, u)_{H^1} \end{aligned}$$

in view of Proposition 1.8.5. Hence, from Proposition 1.8.8, the real space $H_0^1(]a, b[)$ endowed with

$$(u, v)_q = \int_a^b u'v' dx + \int_a^b q uv dx$$

is a (real) Hilbert space. Since, in view of Proposition 1.8.5, the linear form

$$v \mapsto \int_a^b f v dx$$

is continuous on this Hilbert space, the theorem follows from the Riesz representation theorem. \square

Remark 1.8.10 Since u'' belongs to $L^1(]a, b[)$, we infer that u' extends as a continuous function on $[a, b]$. Furthermore, if q, f are continuous functions on $[a, b]$, then u'' is continuous on $[a, b]$, which means that u is C^2 on $[a, b]$, so that the differential equation is satisfied in the usual sense.

1.9 Further properties

1.9.1 Characterisation of Lipschitz functions

Definition 1.9.1 Let k be a positive number and I be an open interval. A function $u : I \rightarrow \mathbb{C}$ is k -Lipschitz if, for every $x, y \in I$,

$$|u(x) - u(y)| \leq k|x - y| .$$

A typical example of a k -Lipschitz function is a C^1 function u such $\|u'\|_\infty \leq k$. Of course, there are Lipschitz functions which are not derivable, like function $x \mapsto |x|$. The following result give a complete description of k -Lipschitz functions.

Theorem 1.9.2 A function $u : \mathbb{R} \rightarrow \mathbb{C}$ is k -Lipschitz if and only if there exists $f \in L^\infty(\mathbb{R})$ such that

$$(1.9.9) \quad \forall x \in \mathbb{R} , u(x) = u(0) + \int_0^x f(t) dt .$$

Remark 1.9.3 A similar result holds on any open interval I , from an adaptation of the proof below on \mathbb{R} .

Proof.— If $\forall x \in \mathbb{R}$, $u(x) = u(0) + \int_0^x f(t) dt$, we have

$$|u(x) - u(y)| = \left| \int_y^x f(t) dt \right| \leq \|f\|_\infty |x - y| .$$

Conversely, let u be a k -Lipschitz function. We are going to prove that u admits a generalised derivative f with $\|f\|_\infty \leq k$. According to Proposition 1.8.2, this will imply property (1.9.9).

As a first step, we are going to prove the following inequality,

$$(1.9.10) \quad \forall \varphi \in \mathcal{C}_0^\infty(I), \quad |\langle T'_u, \varphi \rangle| \leq k \|\varphi\|_{L^1} .$$

Indeed, we have,

$$\begin{aligned} \langle T'_u, \varphi \rangle &= - \int_{\mathbb{R}} u(x) \varphi'(x) dx = - \int_{\mathbb{R}} u(x) \lim_{h \rightarrow 0} \frac{\varphi(x+h) - \varphi(x)}{h} dx \\ &= - \lim_{h \rightarrow 0} \int_{\mathbb{R}} u(x) \frac{\varphi(x+h) - \varphi(x)}{h} dx , \end{aligned}$$

by using either dominated convergence or uniform convergence. Next we decompose the integral as follows,

$$\begin{aligned} \int_{\mathbb{R}} u(x) \frac{\varphi(x+h) - \varphi(x)}{h} dx &= \frac{1}{h} \left(\int_{\mathbb{R}} u(x) \varphi(x+h) dx - \int_{\mathbb{R}} u(x) \varphi(x) dx \right) \\ &= \frac{1}{h} \left(\int_{\mathbb{R}} u(y-h) \varphi(y) dy - \int_{\mathbb{R}} u(x) \varphi(x) dx \right) \\ &= \int_{\mathbb{R}} \frac{u(y-h) - u(y)}{h} \varphi(y) dy , \end{aligned}$$

and, by the Lipschitz property of u , the modulus of the latter expression is bounded by $k \|\varphi\|_{L^1}$. This leads to inequality (1.9.10). At this stage we appeal to the following functional analytic lemma.

Lemma 1.9.4 Let \mathcal{E} be a normed vector space, and let \mathcal{D} be a dense vector subspace of \mathcal{E} . Let $L : \mathcal{D} \rightarrow \mathbb{C}$ be a linear form, which is continuous for the norm on \mathcal{E} . Then L admits a unique continuation \tilde{L} as a continuous linear form on \mathcal{E} , and $\|\tilde{L}\| = \|L\|$.

Applying Lemma 1.9.4 to $\mathcal{E} = L^1(\mathbb{R})$, $\mathcal{D} = \mathcal{D}(\mathbb{R})$ and $L = T'_u$, we infer that T'_u extends as a continuous linear form on $L^1(\mathbb{R})$, of norm at most k . From the $L^1 - L^\infty$ duality theorem, there exists $f \in L^\infty(\mathbb{R})$ such that

$$\forall \varphi \in \mathcal{C}_0^\infty(I), \quad \langle T'_u, \varphi \rangle = \int_{\mathbb{R}} f(x) \varphi(x) dx ,$$

and $\|f\|_\infty \leq k$. This precisely means that f is the generalised derivative of u , whence (1.9.9). \square

1.9.2 Derivation and integration under the bracket

This paragraph provides two very useful rules of calculations which turn out to be crucial when dealing with convolution of distributions — see the next paragraph.

Proposition 1.9.5 (Derivation under the bracket) Let I, J be open intervals, $\psi : I \times J \rightarrow \mathbb{C}$ be a \mathcal{C}^∞ function such that there exists a segment $[a, b] \subset I$ for which

$$\forall z \in J, \text{supp } \psi(\cdot, z) \subset [a, b],$$

where we have set $\psi(\cdot, z) : x \in I \rightarrow \psi(x, z) \in \mathbb{C}$. Define

$$g(z) = \langle T, \psi(\cdot, z) \rangle, \quad z \in J.$$

Then $g \in \mathcal{C}^\infty(J)$ and

$$\forall z \in J, g'(z) = \left\langle T, \frac{\partial \psi}{\partial z}(\cdot, z) \right\rangle.$$

Proof.— Let us first prove that g is derivable on J . We calculate, for $z \in J$ and $h \neq 0$ small enough,

$$\frac{g(z+h) - g(z)}{h} = \left\langle T, \frac{\psi(\cdot, z+h) - \psi(\cdot, z)}{h} \right\rangle.$$

Now we observe that, since $\psi \in \mathcal{C}^\infty(I \times J)$,

$$\forall x \in I, \frac{\psi(x, z+h) - \psi(x, z)}{h} \xrightarrow{h \rightarrow 0} \frac{\partial \psi}{\partial z}(x, z).$$

We claim that the convergence takes place in $\mathcal{C}_0^\infty(I)$. Indeed, we already know that the left hand side is a smooth function of x which is supported in $[a, b]$. We claim that it converges uniformly on I to the right hand side. Indeed,

$$\frac{\psi(x, z+h) - \psi(x, z)}{h} = \int_0^1 \frac{\partial \psi}{\partial z}(x, z+th) dt.$$

Fix $\varepsilon_0 > 0$ such that $|h| \leq \varepsilon_0$. Then the continuous function $\frac{\partial \psi}{\partial z}$ is uniformly continuous on the compact subset $[a, b] \times [z - \varepsilon_0, z + \varepsilon_0]$. Consequently,

$$\sup_{x \in [a, b]} \left| \int_0^1 \frac{\partial \psi}{\partial z}(x, z+th) dt - \frac{\partial \psi}{\partial z}(x, z) \right| \xrightarrow{h \rightarrow 0} 0.$$

The same argument holds for every derivative in x ,

$$\frac{\psi^{(k)}(x, z+h) - \psi^{(k)}(x, z)}{h} \xrightarrow{h \rightarrow 0} \frac{\partial \psi^{(k)}}{\partial z}(x, z)$$

uniformly for $x \in I$.

Using the continuity property of T , we infer

$$\frac{g(z+h) - g(z)}{h} \xrightarrow{h \rightarrow 0} \left\langle T, \frac{\partial \psi}{\partial z}(\cdot, z) \right\rangle,$$

which shows that g is derivable with the claimed formula for $g'(z)$. Finally, by applying this result, one easily proves by induction on n that g is n times derivable with

$$g^{(n)}(z) = \left\langle T, \frac{\partial^n \psi}{\partial z^n}(\cdot, z) \right\rangle.$$

This completes the proof. □

Proposition 1.9.6 (Integration under the bracket) Let I, J be open intervals, $\psi : I \times J \rightarrow \mathbb{C}$ be a C^∞ function such that there exists segments $[a, b] \subset I$ and $[c, d] \subset J$ for which

$$\forall (x, z) \notin [a, b] \times [c, d], \psi(x, z) = 0.$$

Then

$$\int_J \langle T, \psi(\cdot, z) \rangle dz = \left\langle T, \int_J \psi(\cdot, z) dz \right\rangle.$$

Proof.— Consider

$$\tilde{\psi}(x, z) = \int_{t \in J, t < z} \psi(x, t) dt.$$

Then $\tilde{\psi}$ satisfies the assumptions of Proposition 1.9.5, and

$$\frac{\partial \tilde{\psi}}{\partial z}(x, z) = \psi(x, z).$$

Applying this proposition leads to

$$\frac{d}{dz} \langle T, \tilde{\psi}(\cdot, z) \rangle = \langle T, \psi(\cdot, z) \rangle.$$

We integrate both sides on J . This gives

$$\langle T, \tilde{\psi}(\cdot, d) \rangle - \langle T, \tilde{\psi}(\cdot, c) \rangle = \int_J \langle T, \psi(\cdot, z) \rangle dz.$$

In view of the assumptions on ψ , we have

$$\langle T, \tilde{\psi}(\cdot, d) \rangle = \left\langle T, \int_J \psi(\cdot, z) dz \right\rangle, \quad \langle T, \tilde{\psi}(\cdot, c) \rangle = 0.$$

This completes the proof. □

1.9.3 Convolution and regularisation

In this section, we generalise the convolution with a test function to a distribution.

Definition 1.9.7 Let $T \in \mathcal{D}'(\mathbb{R})$ and $\varphi \in \mathcal{D}(\mathbb{R})$. For every $x \in \mathbb{R}$, we denote by $\varphi(x - \cdot)$ the element of $\mathcal{D}(\mathbb{R})$ which maps $y \in \mathbb{R}$ to $\varphi(x - y) \in \mathbb{C}$. We set

$$T * \varphi(x) = \langle T, \varphi(x - \cdot) \rangle.$$

Notice that, if $T = T_f$ for $f \in L^1_{\text{loc}}(\mathbb{R})$, we have

$$T_f * \varphi(x) = \int_{\mathbb{R}} f(y) \varphi(x - y) dy = f * \varphi(x),$$

in the usual sense of convolution of functions.

Proposition 1.9.8 For every $(T, \varphi) \in \mathcal{D}'(\mathbb{R}) \times \mathcal{D}(\mathbb{R})$, $T * \varphi \in \mathcal{C}^\infty(\mathbb{R})$, and

$$(T * \varphi)' = T * \varphi' = T' * \varphi .$$

Proof.— It is enough to prove that $T * \varphi$ is smooth on every finite open interval $]a, b[$. If $\text{supp } \varphi \subset [a, b]$ and $x \in]a, b[$, we have $\text{supp } \varphi(x - \cdot) \subset [\alpha - b, \beta - a]$, so that the assumptions of Proposition 1.9.5 are fulfilled with $I = \mathbb{R}$, $J =]\alpha, \beta[$ and

$$\psi(y, x) = \varphi(x - y) .$$

Consequently, $T * \varphi \in \mathcal{C}^\infty(\mathbb{R})$, and

$$(T * \varphi)'(x) = \langle T, \partial_x \varphi(x - \cdot) \rangle = \langle T, \varphi'(x - \cdot) \rangle = T * \varphi'(x) .$$

Furthermore, we have

$$\varphi'(x - y) = -\partial_y \varphi(x - y) ,$$

so that

$$(T * \varphi)'(x) = \langle T, -\partial_y \varphi(x - \cdot) \rangle = \langle T', \varphi(x - \cdot) \rangle = T' * \varphi(x) .$$

□

Let us come to regularisation. Let $\rho \in \mathcal{C}_0^\infty(\mathbb{R})$ with

$$\int_{\mathbb{R}} \rho(x) dx = 1 .$$

For every $\varepsilon > 0$, we set

$$\rho_\varepsilon(x) = \frac{1}{\varepsilon} \rho\left(\frac{x}{\varepsilon}\right) .$$

It is well known that, if $\varphi \in \mathcal{C}_0^\infty(\mathbb{R})$,

$$\rho_\varepsilon * \varphi \xrightarrow{\varepsilon \rightarrow 0} \varphi$$

in $\mathcal{D}(\mathbb{R})$. We set $f_\varepsilon = T * \rho_\varepsilon$.

Proposition 1.9.9 For every $T \in \mathcal{D}'(\mathbb{R})$,

$$T_{f_\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} T .$$

Proof.— We have to prove that

$$(1.9.11) \quad \forall \varphi \in \mathcal{D}(\mathbb{R}) , \quad \int_{\mathbb{R}} f_\varepsilon(x) \varphi(x) dx \xrightarrow{\varepsilon \rightarrow 0} \langle T, \varphi \rangle .$$

Note that

$$\int_{\mathbb{R}} f_\varepsilon(x) \varphi(x) dx = \int_{\mathbb{R}} \langle T, \rho_\varepsilon(x - \cdot) \rangle \varphi(x) dx = \int_{\mathbb{R}} \langle T, \rho_\varepsilon(x - \cdot) \varphi(x) \rangle dx ,$$

and that

$$\psi(y, x) := \rho_\varepsilon(x - y) \varphi(x)$$

satisfies the assumption of Proposition 1.9.6 on $\mathbb{R} \times \mathbb{R}$. Indeed, if $\text{supp } \rho \subset [-C, C]$, $\text{supp } \varphi \subset [a, b]$, then

$$\forall (y, x) \notin [a - \varepsilon C, b + \varepsilon C] \times [a, b], \psi(y, x) = 0.$$

Applying Proposition 1.9.6, we obtain

$$\int_{\mathbb{R}} \langle T, \rho_{\varepsilon}(x - \cdot) \varphi(x) \rangle dx = \left\langle T, \int_{\mathbb{R}} \rho_{\varepsilon}(x - \cdot) \varphi(x) dx \right\rangle = \langle T, \tilde{\rho}_{\varepsilon} * \varphi \rangle$$

where $\tilde{\rho}(z) = \rho(-z)$. Since $\tilde{\rho}$ satisfies the same assumption as ρ , we have $\tilde{\rho}_{\varepsilon} * \varphi \xrightarrow{\varepsilon \rightarrow 0} \varphi$ in $\mathcal{D}(\mathbb{R})$, and consequently

$$\int_{\mathbb{R}} f_{\varepsilon}(x) \varphi(x) dx \xrightarrow{\varepsilon \rightarrow 0} \langle T, \varphi \rangle.$$

□

1.9.4 Positive distributions and increasing functions

Definition 1.9.10 A distribution T on I is said positive (we note $T \geq 0$) if, for every $\varphi \in \mathcal{D}(I)$ valued in $[0, +\infty[$, we have $\langle T, \varphi \rangle \in [0, +\infty[$.

For instance, it is easy to check that, if $f \in L^1_{\text{loc}}(I)$, $T_f \geq 0$ if and only if $f \geq 0$ a.e. Another example is $T = c\delta_{x_0}$ with $c \geq 0$ and $x_0 \in I$. On the other hand, whatever $c \neq 0$ is, it is clear that $c\delta'_{x_0}$ cannot be positive. In fact, we have the following general result.

Proposition 1.9.11 If T is a positive distribution, then T has order 0, namely

$$\forall [a, b] \subset I, \exists C > 0, \forall \varphi \in \mathcal{D}(I), \text{supp } \varphi \subset [a, b] \Rightarrow |\langle T, \varphi \rangle| \leq C \|\varphi\|_{\infty}.$$

Proof.— Let $[a, b] \subset I$ and $\chi \in \mathcal{D}(I)$ be a plateau function on $[a, b]$. Let $\varphi \in \mathcal{D}(I)$ be real valued and such that $\text{supp } \varphi \subset [a, b]$. Then

$$-\chi \|\varphi\|_{\infty} \leq \varphi \leq \chi \|\varphi\|_{\infty}.$$

Consequently, the positivity of T implies $\langle T, \varphi \rangle \in \mathbb{R}$ and

$$-\langle T, \chi \rangle \|\varphi\|_{\infty} \leq \langle T, \varphi \rangle \leq \langle T, \chi \rangle \|\varphi\|_{\infty},$$

and therefore

$$|\langle T, \varphi \rangle| \leq \langle T, \chi \rangle \|\varphi\|_{\infty}.$$

If φ is complex valued with $\text{supp } \varphi \subset [a, b]$, we decompose

$$\varphi = \text{Re}(\varphi) + i \text{Im}(\varphi)$$

and we conclude

$$|\langle T, \varphi \rangle| \leq 2 \langle T, \chi \rangle \|\varphi\|_{\infty}.$$

□

Remark 1.9.12 Since any compactly supported (positive) continuous function on I can be approximated uniformly by a sequence of (positive) test functions supported in a fixed segment of I , the above proposition implies that any positive distribution extends as a positive linear form on compactly supported continuous functions. Hence, according to the Riesz representation theorem, there exists a positive Borel measure μ on I , finite on segments, such that

$$\forall \varphi \in \mathcal{D}(I), \langle T, \varphi \rangle = \int_I \varphi(x) d\mu(x).$$

Let us come to the main result of this paragraph, which characterises increasing functions. For simplicity, we state and prove this result on \mathbb{R} , but a similar result holds on any open interval. Recall that an increasing function on \mathbb{R} is a function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\forall x \geq y, f(x) \geq f(y).$$

(sometimes such functions are called nondecreasing functions).

Theorem 1.9.13 If $f : \mathbb{R} \rightarrow \mathbb{R}$ is an increasing function, then $T'_f \geq 0$. Conversely, if $T \in \mathcal{D}'(\mathbb{R})$ is such that $T' \geq 0$ and $\langle T, \varphi \rangle \in \mathbb{R}$ for every real valued test function φ , then there exists an increasing function f on \mathbb{R} such that $T_f = T$.

Proof.— Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be an increasing function. Note that f is bounded on every segment, hence it is locally integrable, and it makes sense to consider T_f . Let $\varphi \in \mathcal{D}(\mathbb{R})$, valued in $[0, +\infty[$. Let us calculate

$$\begin{aligned} \langle T'_f, \varphi \rangle &= - \int_{\mathbb{R}} u(x) \varphi'(x) dx = - \int_{\mathbb{R}} f(x) \lim_{h \rightarrow 0} \frac{\varphi(x+h) - \varphi(x)}{h} dx \\ &= - \lim_{h \rightarrow 0} \int_{\mathbb{R}} f(x) \frac{\varphi(x+h) - \varphi(x)}{h} dx \\ &= \lim_{h \rightarrow 0} \int_{\mathbb{R}} \frac{f(y-h) - f(y)}{h} \varphi(y) dy, \end{aligned}$$

by the same arguments as in subsection 1.9.1. It yields $\langle T'_f, \varphi \rangle \geq 0$, hence T'_f is positive. Conversely, let $T \in \mathcal{D}'(\mathbb{R})$ such that $T' \geq 0$. Choose $\rho \in \mathcal{D}(\mathbb{R})$, supported in \mathbb{R}_+ , satisfying

$$0 \leq \rho, \int_{\mathbb{R}} \rho(x) dx = 1.$$

Recall that, from Proposition 1.9.9, the smooth function

$$f_\varepsilon = T * \rho_\varepsilon$$

is such that $T_{f_\varepsilon} \rightarrow T$ in $\mathcal{D}'(\mathbb{R})$. Notice moreover that, since T takes real values on real valued test functions, f_ε is real valued. Our strategy is to prove that, for an appropriate choice of ρ , f_ε is an increasing function and converges for the simple convergence to a function f , with local dominated convergence. This will imply that $T_{f_\varepsilon} \rightarrow T_f$, hence $T = T_f$.

Let us first prove that f_ε is increasing. We know that

$$f'_\varepsilon = T' * \rho_\varepsilon$$

which is ≥ 0 because $T' \geq 0$ and $\rho_\varepsilon \geq 0$. Therefore f_ε is an increasing function on \mathbb{R} .

We claim that moreover f_ε is an decreasing function of ε . In fact, applying Proposition 1.9.5 of derivation under the bracket, we have that $f_\varepsilon(x)$ is a C^∞ function of $\varepsilon > 0$, and that

$$\frac{d}{d\varepsilon} f_\varepsilon(x) = \left\langle T, \frac{\partial}{\partial \varepsilon} \rho_\varepsilon(x - \cdot) \right\rangle .$$

Observe that

$$\begin{aligned} \frac{\partial}{\partial \varepsilon} \rho_\varepsilon(x - y) &= \frac{\partial}{\partial \varepsilon} \left[\frac{1}{\varepsilon} \rho \left(\frac{x - y}{\varepsilon} \right) \right] \\ &= -\frac{1}{\varepsilon^2} \left[\rho \left(\frac{x - y}{\varepsilon} \right) + \frac{x - y}{\varepsilon} \rho' \left(\frac{x - y}{\varepsilon} \right) \right] \\ &= \frac{1}{\varepsilon^2} \frac{\partial}{\partial y} \left[(x - y) \rho \left(\frac{x - y}{\varepsilon} \right) \right] \end{aligned}$$

Consequently,

$$\frac{d}{d\varepsilon} f_\varepsilon(x) = -\frac{1}{\varepsilon} \left\langle T', \frac{(x - \cdot)}{\varepsilon} \rho \left(\frac{x - \cdot}{\varepsilon} \right) \right\rangle \leq 0 ,$$

because $T' \geq 0$ and $z\rho(z) \geq 0$ in view of the assumptions on ρ .

Now we claim that, as $\varepsilon \rightarrow 0$, for every $x \in I$, $f_\varepsilon(x)$ is bounded from above. Indeed, consider $\psi \in \mathcal{D}(I)$, valued in $[0, +\infty[$, supported in $[x, +\infty[$, and such that

$$\int_{\mathbb{R}} \psi(y) dy = 1 .$$

Then, because $f_\varepsilon(y)$ is an increasing function of y ,

$$f_\varepsilon(x) = f_\varepsilon(x) \int_{\mathbb{R}} \psi(y) dy \leq \int_{\mathbb{R}} f_\varepsilon(y) \psi(y) dy \xrightarrow{\varepsilon \rightarrow 0} \langle T, \psi \rangle$$

because of Proposition 1.9.9. Since $f_\varepsilon(x)$ is increasing as ε decreases to 0, we conclude that

$$f_\varepsilon(x) \xrightarrow{\varepsilon \rightarrow 0} f(x)$$

where f is an increasing function on \mathbb{R} . Furthermore, if $x \in [a, b] \subset \mathbb{R}$

$$f(x) \geq f_\varepsilon(x) \geq f_\varepsilon(a) \xrightarrow{\varepsilon \rightarrow 0} f(a) .$$

Therefore we can apply the dominated convergence theorem and conclude that, for every $\varphi \in \mathcal{D}(\mathbb{R})$,

$$\int_{\mathbb{R}} f_\varepsilon(x) \varphi(x) dx \xrightarrow{\varepsilon \rightarrow 0} \int_{\mathbb{R}} f(x) \varphi(x) dx .$$

Summing up, applying again Proposition 1.9.9, we have proved

$$\langle T, \varphi \rangle = \int_{\mathbb{R}} f(x) \varphi(x) dx ,$$

hence $T = T_f$. □

Remark 1.9.14 The first point of Theorem 1.9.13 says that, if f is increasing, T'_f is a positive distribution, hence is given by a positive Borel measure, finite on segments. This measure is called the Stieltjes measure associated to f , and the formula for the integral of a compactly supported continuous function with respect to this measure is an extension of the definition of the usual integral of a continuous function by using Riemann sums. More precisely, if $\text{supp } \varphi \subset [a, b]$,

$$\langle T'_f, \varphi \rangle = \lim_{N \rightarrow \infty} \sum_{j=0}^{N-1} \varphi \left(a + j \frac{b-a}{N} \right) \left[f \left(a + (j+1) \frac{b-a}{N} \right) - f \left(a + j \frac{b-a}{N} \right) \right].$$

For instance, if $f : \mathbb{R} \rightarrow \mathbb{R}$ is the repartition function of a random variable on \mathbb{R} , then T'_f is nothing but the law of this random variable.

1.9.5 The structure of distributions

Recall that, by Proposition 1.4.10, every distribution T has a primitive S . Furthermore, as can be observed from the proof of 1.4.10, if T is real, then one can choose S real. Now, if T is positive, then S satisfies the assumption of Theorem 1.9.13, and there exists an increasing function f such that $T_f = S$, or $T = T'_f$. In other words, any positive distribution is the derivative of T_f , where f is increasing. This statement extends to any distribution of order 0 as follows.

Theorem 1.9.15 Any distribution of order 0 on \mathbb{R} is of the form T'_f , where $f \in L^\infty_{\text{loc}}(\mathbb{R})$.

Proof.— Let us come back to the proof of Proposition 1.4.10. Let $\chi \in \mathcal{D}(\mathbb{R})$ supported in $] -1, 1[$ and satisfying

$$\int_{\mathbb{R}} \chi(y) dy = 1.$$

For every $\psi \in \mathcal{D}(\mathbb{R})$, we set

$$P(\psi)(x) = \int_{-\infty}^x \left[\psi(t) - \chi(t) \int_{\mathbb{R}} \psi(y) dy \right] dt.$$

Notice that $P(\psi) \in \mathcal{D}(\mathbb{R})$ and, if $\text{supp}(\psi) \subset] -n, n[$ for some $n \geq 1$, then $\text{supp}(P(\psi)) \subset] -n, n[$. Since $P(\psi') = \psi$, we infer that the distribution S defined by

$$\langle S, \psi \rangle = -\langle T, P(\psi) \rangle$$

satisfies $S' = T$. Since T is of order 0, for every $n \geq 1$, there exists $C_n > 0$ such that

$$\forall \varphi \in \mathcal{D}(] -n, n[), |\langle T, \varphi \rangle| \leq C_n \|\varphi\|_\infty.$$

Consequently, if $\psi \in \mathcal{D}(] -n, n[)$,

$$|\langle S, \psi \rangle| = |\langle T, P(\psi) \rangle| \leq C_n \|P(\psi)\|_\infty \leq C_n B \|\psi\|_{L^1}.$$

Arguing as in proof of Theorem 1.9.2, we infer that there exists $f_n \in L^\infty(] -n, n[)$ such that

$$S|_{] -n, n[} = T_{f_n}.$$

Therefore we have constructed a sequence $(f_n)_{n \geq 1}$ of functions $f_n \in L^\infty(]-n, n[)$ such that

$$T_{f_{n+1}|_{]-n, n[}} = (S|_{]-n-1, n+1[})|_{]-n, n[} = S|_{]-n, n[} = T_{f_n} .$$

Hence $f_{n+1}|_{]-n, n[} = f_n$ and we infer that there exists $f \in L^\infty_{\text{loc}}(\mathbb{R})$ such that

$$\forall n \geq 1, f|_{]-n, n[} = f_n .$$

Since

$$S|_{]-n, n[} = T_{f|_{]-n, n[}}$$

for every $n \geq 1$, we conclude $S = T_f$, and $T = S' = T'_f$. □

Let us now consider the case of distributions of finite order. Let m be a nonnegative integer. We shall say that $T \in \mathcal{D}'(I)$ is of order $\leq m$ if

$$\forall [a, b] \subset I, \exists C > 0, \forall \varphi \in \mathcal{D}(I), \text{supp}(\varphi) \subset [a, b] \Rightarrow |\langle T, \varphi \rangle| \leq C \sup_{0 \leq k \leq m} \|\varphi^{(k)}\|_\infty .$$

Theorem 1.9.16 *i)* If $T \in \mathcal{D}'(\mathbb{R})$ is of order $\leq m$, there exists $f \in L^\infty_{\text{loc}}(\mathbb{R})$ such that $T = T_f^{(m+1)}$.

ii) If $T \in \mathcal{D}'(\mathbb{R})$ is arbitrary, there exists a sequence $(f_n)_{n \geq 1}$ of $L^\infty(\mathbb{R})$ such that on every segment of \mathbb{R} , f_n vanishes for n large enough, and

$$T = \sum_{n=1}^{\infty} T_{f_n}^{(n)} .$$

Proof.— The first statement follows from an induction argument on m , based on Theorem 1.9.15 and on the following lemma.

Lemma 1.9.17 If $T \in \mathcal{D}'(\mathbb{R})$ is of order $\leq m$ with $m \geq 1$, there exists $S \in \mathcal{D}'(\mathbb{R})$ of order $\leq m - 1$ such that $S' = T$.

The proof of this lemma is straightforward, taking into account the formula $\langle S, \psi \rangle = -\langle T, P(\psi) \rangle$ and the fact that, for every $m \geq 1$ and $\psi \in \mathcal{D}(]-n, n[)$,

$$\sup_{0 \leq k \leq m} \|P(\psi)^{(k)}\|_\infty \leq B \|\psi\|_{L^1} + \sup_{0 \leq k \leq m-1} \|\psi^{(k)}\|_\infty \leq B_n \sup_{0 \leq k \leq m-1} \|\psi^{(k)}\|_\infty .$$

Let us prove statement ii). For every integer $j \geq 1$, there exists an integer m_j such that $T|_{]-j-1/2, j+1/2[}$ is of order $\leq m_j$. Furthermore, we may impose without loss of generality that $m_{j+1} > m_j$.

We are going to construct a sequence $(g_j)_{j \geq 1}$ of L^∞ functions on \mathbb{R} such that, for every $j \geq 1$, $\text{supp}(T_{g_j}) \subset [-j, -j+1] \cup [j-1, j]$, and

$$\left(T - \sum_{\ell=1}^j T_{g_\ell}^{(m_\ell+1)} \right) |_{]-j, j[} = 0 .$$

Let us first construct g_1 . Since T is of order $\leq m_1$ on $] -3/2, 3/2[$, property i) (adapted to an interval) implies that there exists $h_1 \in L_{\text{loc}}^\infty(] -3/2, 3/2[)$ such that

$$T = T_{h_1}^{(m_1+1)} .$$

Then $g_1 := \mathbf{1}_{]-1,1[} h_1 \in L^\infty(\mathbb{R})$ and satisfies

$$\left(T - T_{g_1}^{(m_1+1)} \right)_{\|]-1,1[} = 0 , \quad \text{supp}(T_{g_1}) \subset [-1, 1] .$$

Assuming g_1, \dots, g_j are constructed, let us construct g_{j+1} . Since the sequence m_j is increasing, the distribution

$$\tilde{T}_{j+1} = \left(T - \sum_{\ell=1}^j T_{g_\ell}^{(m_\ell+1)} \right)_{\|]-j-3/2, j+3/2[}$$

is of order $\leq m_{j+1} + 1$. Furthermore, the restriction of \tilde{T}_{j+1} to $] -j, j[$ is 0. Therefore, using again property i), there exists $h_{j+1} \in L_{\text{loc}}^\infty(] -j-3/2, j+3/2[)$ such that

$$\tilde{T}_{j+1} = T_{h_{j+1}}^{(m_{j+1}+1)} .$$

Furthermore, the restriction of $T_{h_{j+1}}^{(m_{j+1}+1)}$ to $] -j, j[$ is 0, which implies from Corollary 1.4.9 that h_{j+1} coincides with a polynomial function p_{j+1} on $] -j, j[$. We consider

$$g_{j+1} := \mathbf{1}_{\|]-j-1, j+1[} (h_{j+1} - p_{j+1}) .$$

Then $g_{j+1} \in L^\infty(\mathbb{R})$, $T_{g_{j+1}}$ is supported in $[-j-1, -j] \cup [j, j+1]$, and

$$\left(\tilde{T}_{j+1} - T_{g_{j+1}}^{(m_{j+1}+1)} \right)_{\|]-j-1, j+1[} = 0 .$$

Coming back to the expression of \tilde{T}_{j+1} , we have checked the induction assumption at rank $j+1$, so that the sequence $(g_j)_{j \geq 1}$ is constructed by induction on j .

In view of the properties of the supports of T_{g_j} , for every j , only the terms of rank $\ell \leq j$ of the series

$$\sum_{\ell=1}^{\infty} T_{g_\ell}^{(m_\ell+1)}$$

have a nonzero restriction to $] -j, j[$. This proves that this series is convergent in $\mathcal{D}'(\mathbb{R})$. Furthermore, by the construction of the g_j , the sum of this series coincides with T on every interval $] -j, j[$, therefore

$$T = \sum_{\ell=1}^{\infty} T_{g_\ell}^{(m_\ell+1)} .$$

Setting $f_n := g_\ell$ if there exists $\ell \geq 1$ such that $n = m_\ell + 1$, and $f_n := 0$ otherwise, we end up with

$$T = \sum_{n=1}^{\infty} T_{f_n}^{(n)} .$$

□

Chapter 2

Distributions in several variables

2.1 A brief review of several variables differential calculus

2.1.1 Scalar product, norm, distance, topology

From now on we shall work on the vector space \mathbb{R}^d , with typical elements

$$x = \begin{pmatrix} x_1 \\ \cdot \\ \cdot \\ x_d \end{pmatrix}$$

with $x_1, \dots, x_d \in \mathbb{R}$. We denote by (e_1, \dots, e_d) the canonical basis, so that

$$x = \sum_{j=1}^d x_j e_j .$$

This canonical basis is an orthonormal basis for the canonical scalar product

$$x \cdot y = \sum_{j=1}^d x_j y_j ,$$

defining the Euclidean norm

$$|x| = \sqrt{x \cdot x} = \left(\sum_{j=1}^d x_j^2 \right)^{\frac{1}{2}} .$$

This norm classically defines the distance

$$d(x, y) = |x - y| ,$$

and a topology —independent of the chosen norm— for which a set Ω is open if and only if for every $a \in \Omega$, there exists $r > 0$ such that $B(a, r) \subset \Omega$. Here, $B(a, r)$ denotes the open ball

$$\{x \in \mathbb{R}^d, d(x, a) < r\} .$$

As in every metric space, a compact subset K of \mathbb{R}^d can be equivalently defined by the Borel–Lebesgue covering property, or by the Bolzano–Weierstrass extraction property for every sequence in K . Since we are in a finite dimensional vector space, compact subsets of \mathbb{R}^d coincide with closed bounded subsets.

If F is a nonempty closed subset of \mathbb{R}^d , we shall often use the distance function to F ,

$$d(x, F) = \inf_{z \in F} d(x, z) .$$

Notice that $d(x, F) = 0$ if and only if $x \in F$, and that, by the Bolzano–Weierstrass property for closed bounded subsets, this infimum is attained at some $z \in F$. Furthermore, the function $d(\cdot, F)$ is continuous on \mathbb{R}^d . In fact, by the triangle inequality, it is 1-Lipschitz continuous,

$$|d(x, F) - d(y, F)| \leq d(x, y) .$$

If K is a compact subset of \mathbb{R}^d and $\delta > 0$, the set

$$K_\delta = \{x \in \mathbb{R}^d, d(x, K) \leq \delta\}$$

is a compact subset containing K in its interior. The following lemma will be of constant use.

Lemma 2.1.1 If K is a compact subset of an open subset Ω of \mathbb{R}^d , then

$$\inf_{x \in K} d(x, \Omega^c) = \delta_0 > 0 .$$

For every $\delta \in]0, \delta_0[$, K_δ is contained in Ω .

Proof.— The first statement follows from the continuity of $d(\cdot, \Omega^c)$, which consequently attains its minimum on K , and from the assumption $K \cap \Omega^c = \emptyset$. The second statement is an elementary consequence of the triangle inequality, since, for every $z \in \Omega^c$, $y \in K$,

$$d(x, z) \geq |d(y, z) - d(x, y)|$$

hence, if $x \in K_\delta$, choosing $y \in K$ such that $d(x, y) = d(x, K)$, $d(x, \Omega^c) \geq \delta_0 - \delta > 0$. □

Finally, we denote by $C^0(\Omega)$ the space of continuous functions $f : \Omega \rightarrow \mathbb{C}$.

2.1.2 Partial derivative, C^1 functions, differential, gradient

Definition 2.1.2 Given $f : \Omega \rightarrow \mathbb{C}$ and $a \in \Omega$, $j \in \{1, \dots, d\}$, we say that f admits a j -th partial derivative at a if the function $t \mapsto f(a + te_j)$, locally defined for t in a neighbourhood of 0 in \mathbb{R} , has a derivative at $t = 0$. We set

$$\frac{d}{dt} f(a + te_j)|_{t=0} = \frac{\partial f}{\partial x_j}(a) = \partial_j f(a) .$$

We say that f is a C^1 function on Ω if it admits a j -th partial derivative for every $j \in \{1, \dots, d\}$ at every point $a \in \Omega$, and if the functions $\partial_j f$ are continuous on Ω . We denote by $C^1(\Omega)$ the space of C^1 functions on Ω .

If $f \in C^1(\Omega)$, one can prove that f is differentiable at every point a ; using the mean value theorem,

$$f(a+h) = f(a) + L_a(h) + o(|h|) \text{ as } h \rightarrow 0$$

where

$$L_a(h) = \sum_{j=1}^d \partial_j f(a) h_j .$$

The linear map $L_a : \mathbb{R}^d \rightarrow \mathbb{C}$ is called the differential of f at point a , and usually denoted as

$$L_a = d_a f .$$

If f is real valued, $d_a f$ is a linear form on the Euclidean space \mathbb{R}^d , hence it can be represented by the scalar product with a vector, called the gradient of f at a , and denoted by $\nabla f(a)$,

$$d_a f(h) = \nabla f(a) \cdot h, \quad \nabla f(a) = \begin{pmatrix} \partial_1 f(a) \\ \vdots \\ \partial_d f(a) \end{pmatrix} .$$

Notice that the mapping $\nabla f : \Omega \rightarrow \mathbb{R}^d$ is continuous.

2.1.3 The Chain rule

Proposition 2.1.3 Let $\Omega \subset \mathbb{R}^d$, $\Omega' \subset \mathbb{R}^p$ be open sets, and $\psi : \Omega \rightarrow \Omega'$, $\psi = (\psi_1, \psi_2, \dots, \psi_p)$, a function of class C^m , $m \geq 1$. Let $f : \Omega' \rightarrow \mathbb{C}$ be a C^m function.

Then $f \circ \psi$ is C^m on Ω . Moreover

$$d_x(f \circ \psi) = d_{\psi(x)} f \cdot d_x \psi = \nabla \psi(x) \cdot \nabla f(\psi(x)),$$

or, for any $j \in \{1, \dots, n\}$,

$$\partial_j(f \circ \psi)(x) = \sum_{k=1}^p \partial_k f(\psi(x)) \partial_j \psi_k(x).$$

2.1.4 Higher order partial derivatives

More generally, for $m \geq 2$, we denote by $C^m(\Omega)$ the vector space of functions $f \in C^1(\Omega)$ whose partial derivatives $\partial_1 f, \partial_2 f, \dots, \partial_d f$ belong to $C^{m-1}(\Omega)$. Moreover, $C^\infty(\Omega)$ is the intersection of all $C^m(\Omega)$.

In general, the order in which one computes repeated partial derivatives matters, but this is not the case for C^2 functions:

Proposition 2.1.4 (Schwarz Lemma) If $f \in C^2(\Omega)$, then, for any $j, k \in \{1, \dots, d\}$,

$$\partial_j(\partial_k f) = \partial_k(\partial_j f)$$

In particular for C^∞ functions, one can compute partial derivatives of f in any order. It is therefore very convenient to use multi-indices.

2.1.5 Multi-indices

Let $f \in C^\infty(\Omega)$, and $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d) \in \mathbb{N}^d$ a multiindex. We denote $\partial^\alpha f$ the function

$$\partial^\alpha f = \partial_1^{\alpha_1} \partial_2^{\alpha_2} \dots \partial_d^{\alpha_d} f.$$

The number

$$|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_d,$$

is the order of the partial derivative, and it is called the length of α . The context usually avoids any confusion with the Euclidean norm ! We also set

$$\alpha! = \alpha_1! \alpha_2! \dots \alpha_d!$$

and, for $\beta \in \mathbb{N}^d$ such that $\beta_j \leq \alpha_j$ for all j , which we will write $\beta \leq \alpha$,

$$\binom{\alpha}{\beta} = \frac{\alpha!}{\beta! (\alpha - \beta)!} = \binom{\alpha_1}{\beta_1} \binom{\alpha_2}{\beta_2} \dots \binom{\alpha_d}{\beta_d}.$$

With these notations, the Leibniz formula for the derivatives of a product of functions easily extends to the case of partial derivatives of functions of several variables. Its proof is exactly the same.

Proposition 2.1.5 Let f and g be functions in $C^\infty(\Omega)$, and $\alpha \in \mathbb{N}^d$ a multiindex. We have

$$\partial^\alpha (fg) = \sum_{\beta \in \mathbb{N}^d, \beta \leq \alpha} \binom{\alpha}{\beta} \partial^\beta f \partial^{\alpha - \beta} g.$$

Proof.— We prove the result by induction over $|\alpha|$. If $|\alpha| = 1$, $\partial^\alpha = \partial_j$ for some $j \in \{1, \dots, d\}$, and

$$\partial_j (fg) = (\partial_j f)g + f(\partial_j g),$$

which is the above formula. Suppose then that the formula is true for all multiindices of length $\leq m$. Let $\alpha \in \mathbb{N}^d$ such that $|\alpha| = m + 1$. There exists $j \in \{1, \dots, d\}$ et $\beta \in \mathbb{N}^d$ of length m such that

$$\alpha = \beta + 1_j,$$

where $1_j = (0, \dots, 0, 1, 0, \dots, 0)$ with a 1 as j -th coordinate. With these notations

$$\partial^\alpha (fg) = \partial^{\beta+1_j} (fg) = \partial^\beta (\partial_j (fg)) = \partial^\beta ((\partial_j f)g) + \partial^\beta (f(\partial_j g)).$$

Since β is of length m , the induction assumption gives

$$\begin{aligned} \partial^\alpha (fg) &= \sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \partial^\gamma (\partial_j f) \partial^{\beta - \gamma} g + \sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \partial^\gamma f \partial^{\beta - \gamma} (\partial_j g) \\ &= \sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \partial^{\gamma+1_j} f \partial^{\beta - \gamma} g + \sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \partial^\gamma f \partial^{\beta+1_j - \gamma} g \\ &= \sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \partial^{\gamma+1_j} f \partial^{\alpha - (\gamma+1_j)} g + \sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \partial^\gamma f \partial^{\alpha - \gamma} g \end{aligned}$$

We change the multiindex in the first sum: $\gamma \leftarrow \gamma + 1_j$, and we get

$$\begin{aligned}\partial^\alpha(fg) &= \sum_{1_j \leq \gamma \leq \alpha} \binom{\beta}{\gamma - 1_j} \partial^\gamma f \partial^{\alpha - \gamma} g + \sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \partial^\gamma f \partial^{\alpha - \gamma} g \\ &= \partial^\alpha f g + \sum_{1_j \leq \gamma \leq \beta} \left(\binom{\beta}{\gamma - 1_j} + \binom{\beta}{\gamma} \right) \partial^\gamma f \partial^{\alpha - \gamma} g + f \partial^\alpha g \\ &= \partial^\alpha f g + \sum_{1_j \leq \gamma \leq \beta} \binom{\beta + 1_j}{\gamma} \partial^\gamma f \partial^{\alpha - \gamma} g + f \partial^\alpha g,\end{aligned}$$

which is the required result. \square

We can continue the analogy with the 1 variable case: if $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$, we denote by x^α the monomial

$$x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \dots x_d^{\alpha_d}.$$

Then we can show, the same way as for the Leibniz formula, that for $x, y \in \mathbb{R}^d$ and $\alpha \in \mathbb{N}^d$,

$$(x + y)^\alpha = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} x^\beta y^{\alpha - \beta}.$$

With these notations, Taylor's formula can be written as

Proposition 2.1.6 Let $f : \Omega \subset \mathbb{R}^d \rightarrow \mathbb{C}$ a function of class \mathcal{C}^{m+1} . Let $a, b \in \Omega$, such that the segment $[a, b]$ is included in Ω . We have

$$f(b) = \sum_{|\alpha| \leq m} \frac{(b-a)^\alpha}{\alpha!} \partial^\alpha f(a) + (m+1) \sum_{|\alpha|=m+1} \frac{(b-a)^\alpha}{\alpha!} \int_0^1 (1-t)^m \partial^\alpha f(a + t(b-a)) dt.$$

Proof.— We have already seen that if $\varphi : \mathbb{R} \rightarrow \mathbb{C}$ is smooth, we have

$$\varphi(1) = \sum_{k=0}^m \frac{1}{k!} \varphi^{(k)}(0) + \frac{1}{m!} \int_0^1 (1-s)^m \varphi^{(m+1)}(s) ds.$$

We shall use this result for the function $\varphi : t \mapsto f(a + t(b-a))$. Notice that

$$\varphi'(t) = \sum_{j=1}^d (b-a)_j (\partial_j f)(a + t(b-a))$$

and more generally that

$$\varphi^{(k)}(t) = \sum_{j_1, j_2, \dots, j_k=1}^d (b-a)_{j_1} \dots (b-a)_{j_k} (\partial_{j_1} \dots \partial_{j_k} f)(a + t(b-a)).$$

This sum only contains terms of the form $(b-a)^\alpha (\partial^\alpha f)(a + t(b-a))$ with $\alpha \in \mathbb{N}^d$ of length $|\alpha| = k$. Therefore we can write, for some coefficients $c_\alpha \in \mathbb{R}$,

$$\sum_{j_1, j_2, \dots, j_k=1}^d (b-a)_{j_1} \dots (b-a)_{j_k} (\partial_{j_1} \dots \partial_{j_k} f)(a + t(b-a)) = \sum_{|\alpha|=k} c_\alpha (b-a)^\alpha (\partial^\alpha f)(a + t(b-a)).$$

Denoting $x = b - a$, this equality between two polynomials in x implies that the c_α are given by

$$c_\alpha = \#\{(j_1, j_2, \dots, j_k) \in \{1, \dots, d\}^k, x_1^{\alpha_1} x_2^{\alpha_2} \dots x_d^{\alpha_d} = x_{j_1} \dots x_{j_k}\}.$$

thus

$$c_\alpha = \binom{k}{\alpha_1} \binom{k - \alpha_1}{\alpha_2} \dots \binom{k - \alpha_1 - \dots - \alpha_{n-1}}{\alpha_n} = \frac{k!}{\alpha!}.$$

Indeed, one has to choose first α_1 numbers among j_1, \dots, j_k that should be 1, then α_2 among the $k - \alpha_1$ numbers left that should be 2, ...

Since $a + t(b - a)|_{t=0} = a$, and $a + t(b - a)|_{t=1} = b$, we obtain the stated formula. \square

Exercise 2.1.7 Show that, for $k \in \mathbb{N}$ and $(x_1, x_2, \dots, x_d) \in \mathbb{R}^d$, we have

$$(x_1 + x_2 + \dots + x_d)^k = \sum_{|\alpha|=k} \frac{k!}{\alpha!} x^\alpha.$$

Applying the Taylor formula, we immediately obtain.

Corollary 2.1.8 (Hadamard's formula) Let $f \in \mathcal{C}^{m+1}(\Omega)$, where Ω is a convex open subset of \mathbb{R}^d . If $a \in \Omega$ and $f(a) = 0$, there exist d functions g_1, g_2, \dots, g_d of class $\mathcal{C}^m(\Omega)$ such that

$$f(x) = \sum_{j=1}^d (x_j - a_j) g_j(x).$$

Notice that the convexity assumption of Ω plays a role, since we need that $[a, x] \subset \Omega$. In practice, we shall use this lemma when Ω is a ball.

2.2 Test functions

2.2.1 Definitions. Examples

If $f \in \mathcal{C}^0(\Omega)$, the support of f is the closure of $\{x \in \Omega, f(x) \neq 0\}$ for the topology induced on Ω . This is a closed subset of Ω , denoted by $\text{supp}(f)$. If $m \in \mathbb{N} \cup \{\infty\}$, we denote by $\mathcal{C}_0^m(\Omega)$ the space of functions in $\mathcal{C}^m(\Omega)$ with a compact support. Notice that a compact set for the induced topology in Ω is also a compact subset of \mathbb{R}^d contained in Ω . In particular, elements of $\mathcal{D}(\Omega) = \mathcal{C}_0^\infty(\Omega)$ are called test functions on Ω . As in one space dimension, if $V \subset \Omega$ is an open subset, every element φ of $\mathcal{D}(V)$ can be extended as an element $\underline{\varphi}$ of $\mathcal{D}(\Omega)$ by setting $\underline{\varphi} = 0$ in $\Omega \setminus V$. This allows to identify $\mathcal{D}(V)$ as the subspace of $\mathcal{D}(\Omega)$ defined by $\text{supp}(\underline{\varphi}) \subset V$. We shall often make this identification.

Proposition 2.2.1 *i)* For every $a \in \mathbb{R}^d$, for every $r > 0$, there exists $\varphi \in \mathcal{D}(\mathbb{R}^d)$ such that $\varphi(x) \geq 0$ for every $x \in \mathbb{R}^d$ and $\text{supp}(\varphi) = \overline{B(a, r)}$.

ii) For every compact subset K of Ω , there exists $\chi \in \mathcal{D}(\Omega)$, valued in $[0, 1]$, such that $\chi = 1$ on K .

Proof.— The proof of the first statement is quite similar to its one dimensional analogue. Recall that the function f defined by

$$f(t) = \begin{cases} e^{-\frac{1}{t}} & \text{if } t > 0 \\ 0 & \text{if } t \leq 0 \end{cases}$$

is a C^∞ function. Then the function φ defined by

$$\varphi(x) = f(r^2 - |x - a|^2)$$

satisfies the requirements.

The second statement requires a little more work. Let φ be as in the first statement with $a = 0$ and $r = 1$. Since $\varphi \geq 0$ and is not identically 0, its integral on \mathbb{R}^d is > 0 . Up to dividing φ by this number, we may assume that

$$\int_{\mathbb{R}^d} \varphi(x) dx = 1 .$$

Then, by Lemma 2.1.1, we may choose $\delta > 0$ such that $K_{2\delta} \subset \Omega$. Set

$$\chi(x) = \int_{K_\delta} \varphi\left(\frac{x-y}{\delta}\right) \frac{dy}{\delta^d} .$$

By the Leibniz rule of derivation under the \int sign, $\chi \in C^\infty(\mathbb{R}^d)$. Since $\varphi \geq 0$, we have $\chi \geq 0$ and

$$\chi(x) \leq \int_{\mathbb{R}^d} \varphi\left(\frac{x-y}{\delta}\right) \frac{dy}{\delta^d} = \int_{\mathbb{R}^d} \varphi(z) dz = 1 ,$$

after performing the change of variable $y = x - \delta z$. Hence χ is valued in $[0, 1]$. If $x \in K$ and $y \in (K_\delta)^c$, then $|x - y| > \delta$, hence $(x - y)/\delta \notin \text{supp}(\varphi)$ and the integrand cancels at y . Consequently,

$$\forall x \in K , \chi(x) = \int_{\mathbb{R}^d} \varphi\left(\frac{x-y}{\delta}\right) \frac{dy}{\delta^d} = 1 .$$

Finally, if $x \notin K_{2\delta}$, $y \in K_\delta$, $|x - y| > \delta$, and the integrand identically vanishes, so that $\chi(x) = 0$. Hence the support of χ is included in $K_{2\delta}$, which is contained in Ω , so that $\chi \in C_0^\infty(\Omega)$. \square

Remark 2.2.2 Functions χ as in Proposition 2.2.1 are called *plateau functions* on K in Ω . It may happen that we need plateau functions on a compact neighborhood of K in Ω , for instance K_ε for $\varepsilon > 0$ enough, according to Lemma 2.1.1. We shall call these functions *plateau functions on a neighborhood of K* in Ω .

We now come to the construction of partitions of unity, which turns out to be particularly useful in several variables, through the so-called “gluing principle”.

2.2.2 Partitions of unity

Proposition 2.2.3 (Partition of unity) Let K be a compact covered by a finite collection $\Omega_1, \dots, \Omega_n$ of open subsets of Ω . There exist $\chi_1 \in C_0^\infty(\Omega_1), \dots, \chi_n \in C_0^\infty(\Omega_n)$, valued in $[0, 1]$, such that

$$\chi_1 + \dots + \chi_n = 1 \text{ on } K .$$

In particular, if $\varphi \in C_0^\infty(\Omega)$ is supported in $\Omega_1 \cup \dots \cup \Omega_n$, then one can find functions $\varphi_1 \in C_0^\infty(\Omega_1), \dots, \varphi_n \in C_0^\infty(\Omega_n)$, such that

$$\varphi_1 + \dots + \varphi_n = \varphi .$$

Proof.— It is very similar to the one dimensional analogue. First, one proves the

Lemma 2.2.4 (Shrinking Lemma) If $K \subset \Omega_1 \cup \dots \cup \Omega_n$, there exists open subsets U_1, \dots, U_n such that, for every $j \in \{1, \dots, n\}$, $\overline{U_j} \subset \Omega_j$ is compact, and

$$K \subset U_1 \cup \dots \cup U_n .$$

As in the one dimensional case, the proof proceeds by induction on $n \leq 1$. The case $n = 1$ follows from Lemma 2.1.1.

Once the shrinking lemma is proved, set

$$\chi_1 = \psi_1, \chi_2 = \psi_2(1 - \psi_1), \dots, \chi_n = \psi_n(1 - \psi_{n-1}) \dots (1 - \psi_1),$$

where, for every $j \in \{1, \dots, n\}$, ψ_j is a plateau function on $\overline{U_j}$ in Ω_j .

Finally, the last assertion follows by writing $\varphi_j = \chi_j \varphi$, where the χ_j 's are associated to the covering of $\text{supp}(\varphi)$ by $\Omega_1, \dots, \Omega_n$. \square

2.3 Distributions on an open subset of \mathbb{R}^d

2.3.1 Définitions. Exemples

Definition 2.3.1 Let (φ_j) be a sequence of functions in $\mathcal{D}(\Omega)$, and $\varphi \in \mathcal{D}(\Omega)$. We say that (φ_j) tends to φ in $\mathcal{D}(\Omega)$ (or in the $\mathcal{D}(\Omega)$ -sense), when

- i) There exists a compact subset $K \subset \Omega$ such that $\text{supp } \varphi_j \subset K$ for all j .
- ii) For all $\alpha \in \mathbb{N}^d$, $\sup |\partial^\alpha \varphi_j - \partial^\alpha \varphi| \rightarrow 0$ as $j \rightarrow +\infty$.

In that case we may write

$$\varphi = \mathcal{D} - \lim_{j \rightarrow +\infty} \varphi_j.$$

Remark 2.3.2 Notice that, under the conditions of the above definition, we have $\text{supp}(\varphi) \subset K$.

Definition 2.3.3 Let $\Omega \subset \mathbb{R}^d$ an open subset, and T a complex valued linear form on $\mathcal{D}(\Omega)$. One says that T is a distribution on Ω if, for every compact set $K \subset \Omega$,

$$\exists C > 0, \exists m \in \mathbb{N}, \forall \varphi \in \mathcal{C}_0^\infty(\Omega) \text{ with } \text{supp } \varphi \subset K, |T(\varphi)| \leq C \sum_{|\alpha| \leq m} \sup |\partial^\alpha \varphi| = C \|\varphi\|_{\mathcal{C}^m}.$$

We denote by $\mathcal{D}'(\Omega)$ the set of distributions on Ω , and for $T \in \mathcal{D}'(\Omega)$, $\varphi \in \mathcal{D}(\Omega)$, we set $\langle T, \varphi \rangle := T(\varphi)$.

Proposition 2.3.4 A linear form T on $\mathcal{D}(\Omega)$ is a distribution on Ω if and only if $T(\varphi_j) \rightarrow T(\varphi)$ for any sequence (φ_j) of functions in $\mathcal{D}(\Omega)$ that converges to φ in the $\mathcal{D}(\Omega)$ -sense.

The proof is similar to the one dimensional case, as well as the following examples.

Locally integrable functions.

Given $f \in L^1_{loc}(\Omega)$, the formula

$$\langle T_f, \varphi \rangle = \int_{\Omega} f(x) \varphi(x) dx$$

defines a distribution on Ω . Furthermore, the linear mapping

$$f \in L^1_{loc}(\Omega) \mapsto T_f \in \mathcal{D}'(\Omega)$$

is one to one. In the sequel, we shall identify f to T_f .

Dirac masses.

Given $a \in \Omega$, the formula

$$\langle \delta_a, \varphi \rangle = \varphi(a)$$

defines a distribution δ_a on Ω , called the Dirac mass at a . It is not of the form f for $f \in L^1_{loc}(\Omega)$.

As in the previous chapter, one can define finite order distributions, and the corresponding notion of order. In particular, distributions of order $\leq m$ can be extended to continuous linear forms on $\mathcal{C}^m(\Omega)$, non-negative distributions are of order 0 — hence are positive measures, and distributions of order 0 are finite linear combinations of non-negative distributions.

We now come to the important notion of support, for which we shall be a little more specific than we were in Chapter 1.

2.3.2 Restriction and support

Definition 2.3.5 Let $T \in \mathcal{D}'(\Omega)$, and $V \subset \Omega$ an open subset. The restriction of T to V is the distribution $T|_V \in \mathcal{D}'(V)$ defined as

$$\forall \varphi \in \mathcal{D}(V), \quad \langle T|_V, \varphi \rangle = \langle T, \underline{\varphi} \rangle,$$

where $\underline{\varphi}$ denotes the extension of φ by 0 on $\Omega \setminus V$.

We say that T vanishes in V if $T|_V = 0$.

Definition 2.3.6 The support of a distribution $T \in \mathcal{D}'(\Omega)$ is the complement of the union of all the open subsets where T vanishes. We denote it by $\text{supp } T$.

Notice that $\text{supp } T$ is closed, and the following characterizations are convenient.

- $x_0 \notin \text{supp } T$ if and only if there is an open neighborhood V of x_0 such that $T|_V = 0$.
- $x_0 \in \text{supp } T$ if and only if for any open neighborhood V of x_0 , one can find $\varphi \in \mathcal{C}_0^\infty(V)$ such that $\langle T, \varphi \rangle \neq 0$.

As in the one dimensional case, one can characterize distributions supported in one point a , as given by

$$\langle T, \varphi \rangle = \sum_{\alpha} c_{\alpha} \partial^{\alpha} \varphi(a) ,$$

where $(c_{\alpha})_{\alpha \in \mathbb{N}^d}$ is a family of complex numbers which vanish except for a finite set of multi-indices.

Proposition 2.3.7 Let $\varphi \in \mathcal{C}_0^\infty(\Omega)$ and $T \in \mathcal{D}'(\Omega)$. If $\text{supp } \varphi \cap \text{supp } T = \emptyset$, then $\langle T, \varphi \rangle = 0$. In particular, if $\text{supp}(T) = \emptyset$, then $T = 0$.

In view of Lemma 2.2.3, the proof is similar to the one dimensional case. We now emphasize the following corollary, which is very useful.

Corollary 2.3.8 (The gluing principle) Assume

$$\Omega = \bigcup_{j \in J} \Omega_j$$

for some collection $(\Omega_j)_{j \in J}$ of open subsets. Suppose we are given, for every $j \in J$, a distribution T_j on Ω_j , so that we have the following property : for every $j, k \in J$ such that $\Omega_j \cap \Omega_k \neq \emptyset$, then

$$(T_j)|_{\Omega_j \cap \Omega_k} = (T_k)|_{\Omega_j \cap \Omega_k} .$$

Then there exists a unique $T \in \mathcal{D}'(\Omega)$ such that, for every $j \in J$, $T|_{\Omega_j} = T_j$.

Proof.— The uniqueness of T follows from Proposition 2.3.7. Let us prove the existence of T . Given $\varphi \in \mathcal{D}(\Omega)$, we want to define $\langle T, \varphi \rangle$. From Proposition 2.2.3, one can decompose

$$\varphi = \sum_{j \in J} \varphi_j$$

where $\varphi_j \in \mathcal{D}(\Omega_j)$ and $\varphi_j = 0$ except for a finite set of j 's. In such a situation, if T exists, one must have

$$\langle T, \varphi \rangle = \sum_{j \in J} \langle T, \varphi_j \rangle = \sum_{j \in J} \langle T_j, \varphi_j \rangle .$$

Therefore it is natural to establish the following lemma.

Lemma 2.3.9 For every decomposition

$$\varphi = \sum_{j \in J} \varphi_j$$

where $\varphi_j \in \mathcal{D}(\Omega_j)$ and $\varphi_j = 0$ except for a finite set of j 's, the complex number

$$\sum_{j \in J} \langle T_j, \varphi_j \rangle$$

only depends on φ , not on the decomposition.

Let us prove the lemma. Consider the set

$$K = \bigcup_{j \in J} \text{supp}(\varphi_j) .$$

Since $\text{supp}(\varphi_j)$ is compact and is empty except for a finite set of j 's, K is a compact subset of Ω . Apply Proposition 2.2.3 to a finite covering

$$K \subset \bigcup_{r \in R} \Omega_r$$

extracted from the covering of K by the Ω_j 's. The family $(\chi_r)_{r \in R}$ satisfies

$$\text{supp}(\chi_r) \subset \Omega_r , \quad \sum_{r \in R} \chi_r = 1 \text{ on } K .$$

In particular, for every $j \in J$, $\varphi_j = \sum_{r \in R} \chi_r \varphi_j$, therefore

$$\langle T_j, \varphi_j \rangle = \sum_{r \in R} \langle T_j, \chi_r \varphi_j \rangle .$$

For any $r \in R$, we claim that $\langle T_j, \chi_r \varphi_j \rangle = \langle T_r, \chi_r \varphi_j \rangle$. Indeed, either $\Omega_j \cap \Omega_r = \emptyset$, and $\chi_r \varphi_j = 0$, so both sides of this equality cancel. Or $\Omega_j \cap \Omega_r \neq \emptyset$, and, since $\text{supp}(\chi_r \varphi_j) \subset \Omega_j \cap \Omega_r$, the assumption merely says that

$$\langle T_j, \chi_r \varphi_j \rangle = \langle T_r, \chi_r \varphi_j \rangle .$$

Consequently,

$$\begin{aligned} \sum_{j \in J} \langle T_j, \varphi_j \rangle &= \sum_{j \in J} \sum_{r \in R} \langle T_r, \chi_r \varphi_j \rangle = \sum_{r \in R} \langle T_r, \chi_r \left(\sum_{j \in J} \varphi_j \right) \rangle \\ &= \sum_{r \in R} \langle T_r, \chi_r \varphi \rangle . \end{aligned}$$

This proves the lemma. Indeed, given any other decomposition $\varphi = \sum_{j \in J} \tilde{\varphi}_j$, just apply the above construction with

$$K = \bigcup_{j \in J} \text{supp}(\varphi_j) \cup \bigcup_{j \in J} \text{supp}(\tilde{\varphi}_j) ,$$

and we see that

$$\sum_{j \in J} \langle T_j, \varphi_j \rangle = \sum_{r \in R} \langle T_r, \chi_r \varphi \rangle = \sum_{j \in J} \langle T_j, \tilde{\varphi}_j \rangle .$$

Let us complete the proof of Proposition 2.3.8. For every decomposition $\varphi = \sum_{j \in J} \varphi_j$ as in the lemma, we define

$$\langle T, \varphi \rangle = \sum_{j \in J} \langle T_j, \varphi_j \rangle .$$

It is clear that T is a linear form on $\mathcal{D}(\Omega)$. To check it is a distribution, let K a compact subset of Ω and $(\chi_r)_{r \in R}$ be a partition of unity associated to a finite covering of K extracted from $K \subset \cup_{j \in J} \Omega_j$. Then, for every test function φ supported in K , we have, from the lemma,

$$\langle T, \varphi \rangle = \sum_{r \in R} \langle T_r, \chi_r \varphi \rangle .$$

Since $T_r \in \mathcal{D}'(\Omega_r)$, we have

$$|\langle T, \chi_r \varphi \rangle| \leq C_r \|\chi_r \varphi\|_{C^{m_r}} \leq C'_r \|\varphi\|_{C^{m_r}} .$$

Then, with $m = \max_{r \in R} m_r$, we obtain

$$|\langle T, \varphi \rangle| \leq \left(\sum_{r \in R} C'_r \right) \|\varphi\|_{C^m} ,$$

so that $T \in \mathcal{D}'(\Omega)$. Finally, if $\varphi \in \mathcal{D}(\Omega_j)$ for some $j \in J$, we can write the decomposition $\varphi = \sum_{k \in J} \varphi_k$, with $\varphi_j = \varphi$ and $\varphi_k = 0$ if $k \neq j$. Using the lemma, we infer $\langle T, \varphi \rangle = \langle T_j, \varphi \rangle$. In other words, $T|_{\Omega_j} = T_j$. \square

Let us close this paragraph by a number of additional remarks concerning Proposition 2.3.8.

- i) If every T_j is a nonnegative distribution, then so is T . Indeed, the elements χ_r of the partitions of unity can be chosen to be nonnegative.
- ii) If every T_j is a C^m function, then so is T .

2.3.3 Multiplication by a smooth function

Definition 2.3.10 Given $q \in C^\infty(\Omega)$ and $T \in \mathcal{D}'(\Omega)$, we define $qT \in \mathcal{D}'(\Omega)$ by

$$\forall \varphi \in \mathcal{D}(\Omega) , \langle qT, \varphi \rangle = \langle T, q\varphi \rangle .$$

Of course, if $T \in L^1_{loc}(\Omega)$, the definition coincides with the usual product. Let us just mention the following generalization of Proposition 1.5.7 in Chapter 1.

Proposition 2.3.11 Let $T \in \mathcal{D}'(\Omega)$ and $a \in \Omega$ such that

$$\forall j \in \{1, \dots, d\} , (x_j - a_j)T = 0 .$$

Then there exists $c \in \mathbb{C}$ such that $T = c\delta_a$.

Proof.— In view of what we did in Chapter 1, the proof reduces to the following lemma.

Lemma 2.3.12 (Hadamard lemma on an arbitrary open set) If $\varphi \in C_0^\infty(\Omega)$ satisfies $\varphi(a) = 0$, there exists ψ_1, \dots, ψ_d in $C_0^\infty(\Omega)$ such that

$$\varphi(x) = \sum_{j=1}^d (x_j - a_j) \psi_j(x) .$$

Let us prove the lemma. Let $r > 0$ such that $B(a, r) \subset \Omega$, and let χ be a plateau function on $\overline{B}(a, r/2)$, supported in $B(a, r)$. From Corollary 2.1.8 — Hadamard's formula on a convex subset — we can write, for $x \in B(a, r)$,

$$\varphi(x) = \sum_{j=1}^d (x_j - a_j) f_j(x) ,$$

with $f_j \in C^\infty(B(a, r))$ for every j . This implies $\chi\varphi = \sum_{j=1}^d (x_j - a_j) \chi f_j$. On the other hand,

$$(1 - \chi(x))\varphi(x) = \sum_{j=1}^d (x_j - a_j) \frac{(x_j - a_j)(1 - \chi(x))}{|x - a|^2} \varphi(x) .$$

Summing both above identities, the lemma follows with

$$\psi_j(x) = \chi(x) f_j(x) + \frac{(x_j - a_j)(1 - \chi(x))}{|x - a|^2} \varphi(x) .$$

□

2.3.4 Derivation

As in one space dimension, we first consider the case of a C^1 function. In this case, we have the following elementary lemma.

Lemma 2.3.13 Let $f \in C^1(\Omega)$, $\varphi \in C_0^1(\Omega)$, and $j \in \{1, \dots, d\}$.

$$\int_{\Omega} \partial_j f(x) \varphi(x) dx = - \int_{\Omega} f(x) \partial_j \varphi(x) dx .$$

Proof.— Let χ be a plateau function on a neighborhood of the support of φ . It is easy to check that

$$\partial_j(\chi f)\varphi = \partial_j f \varphi , \quad \chi f \partial_j \varphi = f \partial_j \varphi .$$

Furthermore, χf can be extended as a C^1 function on \mathbb{R}^d , vanishing outside of Ω . Therefore we are reduced to prove the lemma with $\Omega = \mathbb{R}^d$. This is an immediate consequence of the Fubini theorem and of integration by parts in the x_j variable. □

On the basis of Lemma 2.3.13, we introduce the following definition.

Definition 2.3.14 Let $T \in \mathcal{D}'(\Omega)$ and $j \in \{1, \dots, d\}$. We define $\partial_j T \in \mathcal{D}'(\Omega)$ by

$$\langle \partial_j T, \varphi \rangle = -\langle T, \partial_j \varphi \rangle, \quad \varphi \in \mathcal{D}(\Omega).$$

Similarly, for every $\alpha \in \mathbb{N}^d$, we define $\partial^\alpha T \in \mathcal{D}'(\Omega)$ by

$$\langle \partial^\alpha T, \varphi \rangle = (-1)^{|\alpha|} \langle T, \partial^\alpha \varphi \rangle, \quad \varphi \in \mathcal{D}(\Omega).$$

At this stage, it is natural to ask for the multidimensional analogue of the identity $H^1 = \delta_0$ proved in Chapter 1. A natural statement would be a formula for $\partial_j(\mathbf{1}_U)$, where U is an open subset of Ω and $\mathbf{1}_U$ is the indicator – or characteristic – function of U , equal to 1 on U and to 0 outside of U . However, open subsets in \mathbb{R}^d are complicated enough so that no such formula exists without additional assumptions on U . A relatively general formula of this kind will be the purpose of the next section, devoted to superficial measures and to the jump formula. At this stage, let us just consider two very simple examples in \mathbb{R}^2 .

i) Half-spaces. Consider $H_a = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 > a\} \subset \mathbb{R}^2$ for some $a \in \mathbb{R}$. Then an elementary calculation gives

$$\langle \partial_1(\mathbf{1}_{H_a}), \varphi \rangle = \int_{\mathbb{R}} \varphi(a, x_2) dx_2, \quad \partial_2(\mathbf{1}_{H_a}) = 0.$$

ii) The unit disc. Consider $D = \{(x_1, x_2) \in \mathbb{R}^2, x_1^2 + x_2^2 < 1\} \subset \mathbb{R}^2$. Then using polar coordinates, one easily checks that

$$\langle \partial_1(\mathbf{1}_D), \varphi \rangle = - \int_0^{2\pi} \cos \theta \varphi(\cos \theta, \sin \theta) d\theta, \quad \langle \partial_2(\mathbf{1}_D), \varphi \rangle = - \int_0^{2\pi} \sin \theta \varphi(\cos \theta, \sin \theta) d\theta.$$

Notice that, in all the above examples, $\partial_j(\mathbf{1}_U)$ is a distribution of order 0, supported by the boundary ∂U of U . This fact will be generalized in the next section.

2.3.5 Convergence

Definition 2.3.15 A sequence (T_n) of elements of $\mathcal{D}'(\Omega)$ converges to $T \in \mathcal{D}'(\Omega)$ if

$$\forall \varphi \in \mathcal{D}(\Omega), \quad \langle T_n, \varphi \rangle \rightarrow \langle T, \varphi \rangle.$$

If $f \in L^1(\mathbb{R}^d)$ satisfies

$$\int_{\mathbb{R}^d} f(x) dx = 1,$$

then

$$\frac{1}{\varepsilon^d} f\left(\frac{x}{\varepsilon}\right) \xrightarrow{\varepsilon \rightarrow 0} \delta_0.$$

This observation leads to the following notion of regularization. Fix $\rho \in C_0^\infty(\mathbb{R}^d)$, supported in the unit ball, such that

$$\int_{\mathbb{R}^d} \rho(x) dx = 1,$$

and consider

$$\rho_\varepsilon(x) = \frac{1}{\varepsilon^d} \rho\left(\frac{x}{\varepsilon}\right).$$

Given an open subset Ω in \mathbb{R}^d , set

$$\Omega^\varepsilon := \{x \in \Omega, d(x, \Omega^c) > \varepsilon\}.$$

If $x \in \Omega^\varepsilon$, the function

$$\rho_\varepsilon(x - \cdot) : y \mapsto \rho_\varepsilon(x - y)$$

is supported in the closed ball of radius ε centered at x , which is included in Ω . Hence this function belongs to $C_0^\infty(\Omega)$, and we may define

$$F^\varepsilon(x) = \langle T, \rho_\varepsilon(x - \cdot) \rangle.$$

Proposition 2.3.16 The function F^ε is smooth on Ω^ε , with

$$\partial^\alpha F^\varepsilon(x) = \langle T, \partial^\alpha \rho_\varepsilon(x - \cdot) \rangle, \quad \alpha \in \mathbb{N}^d,$$

and, for every $\varphi \in \mathcal{D}(\Omega)$,

$$\int_{\Omega} F^\varepsilon(x) \varphi(x) dx \xrightarrow{\varepsilon \rightarrow 0} \langle T, \varphi \rangle.$$

Notice that, given a compact subset K in Ω , the integral $\int_{\Omega} F^\varepsilon(x) \varphi(x) dx$ is well defined for ε small enough and every φ supported in K . The proof of Proposition 2.3.16 is a consequence of the following two results, which will be frequently used throughout the course, and are generalisations to several variables of Propositions 1.9.5 and 1.9.6 from Chapter 1.

Proposition 2.3.17 (Derivation under the bracket) Let $\Omega \subset \mathbb{R}^d$, $Z \subset \mathbb{R}^p$ be open sets, and $T \in \mathcal{D}'(\Omega)$. Let also $\varphi \in C_0^\infty(\Omega \times Z)$. The function

$$G : z \in Z \mapsto \langle T, \varphi(\cdot, z) \rangle$$

is C^∞ , and, for $\alpha \in \mathbb{N}^p$,

$$\partial^\alpha G(z) = \langle T, \partial_z^\alpha \varphi(\cdot, z) \rangle$$

Remark 2.3.18 *i)* We have written $\langle T, \varphi(\cdot, z) \rangle$ in place of $\langle T, \varphi_z \rangle$, where $\varphi_z \in C_0^\infty(\Omega)$ is the function given by $\varphi_z(x) = \varphi(x, z)$.

ii) If $T = f \in L_{loc}^1(\Omega)$, we have $G(z) = \int_{\Omega} f(x) \varphi(x, z) dx$, so that, under the above assumptions, we get $G \in C^\infty(Z)$ and we recover the Leibniz rule of derivation under the \int sign,

$$\partial^\alpha G(z) = \int_{\Omega} f(x) \partial_z^\alpha \varphi(x, z) dx.$$

Proof.— Denote by K the projection of the support of φ onto Ω . Let $z_0 \in Z$ and $x \in \Omega$. For $h \in \mathbb{R}^q$, Taylor's formula at order 1 gives

$$\varphi(x, z_0 + h) = \varphi(x, z_0) + \sum_{j=1}^p \partial_{z_j} \varphi(x, z_0) h_j + r(x, z_0, h),$$

$$\text{with } r(x, z_0, h) = 2 \sum_{|\alpha|=2} \frac{h^\alpha}{\alpha!} \int_0^1 (1-t) \partial_z^\alpha \varphi(x, z_0 + th) dt.$$

Since $x \mapsto r(x, z_0, h)$ is C^∞ with support in K ,

$$|\langle T, r(x, z_0, h) \rangle| \leq C \sum_{|\beta| \leq k} \sup |\partial_x^\beta r(x, z_0, h)|$$

for a constant $C > 0$ and an integer $k \in \mathbb{N}$ independent of z_0 and h . But for $|h| \leq 1$,

$$|\partial_x^\beta r(x, z_0, h)| \leq 2 \sum_{|\alpha| \leq 2} \frac{|h^\alpha|}{\alpha!} \int_0^1 (1-t) |\partial_x^\beta \partial_z^\alpha \varphi(x, z_0 + th)| dt \leq C |h|^2 \sum_{|\alpha| \leq 2} \sup_{K \times \bar{B}(z_0, 1)} |\partial_x^\beta \partial_z^\alpha \varphi(x, z)|,$$

Thus

$$|\langle T, r(x, z_0, h) \rangle| = \mathcal{O}(|h|^2).$$

and,

$$G(z_0 + h) = G(z_0) + \sum_{j=1}^q \langle T, \partial_{z_j} \varphi(x, z_0) \rangle h_j + \mathcal{O}(|h|^2),$$

which shows that G is differentiable at z - in particular G is continuous, and that

$$\partial_j G(z) = \langle T, \partial_{z_j} \varphi(x, z) \rangle.$$

Then one can replace $\varphi(x, z)$ by $\partial_{z_j} \varphi(x, z)$ in the above discussion. We see that for all j , $\partial_j \varphi$ is differentiable, thus in particular continuous. So G is C^1 , and the statement of the proposition is true for any $|\alpha| = 1$. One can easily get the general case by induction. \square

Remark 2.3.19 In view of the proof above, the assumption $\varphi \in C_0^\infty(\Omega \times Z)$ is too strong. What we need is indeed that

$$\text{supp}(\varphi) \subset K \times Z$$

for some compact subset K of Ω .

Proposition 2.3.20 (Integration under the bracket) Let $\Omega \subset \mathbb{R}^d$ be an open set, and $T \in \mathcal{D}'(\Omega)$. Let also $\varphi \in C_0^\infty(\Omega \times \mathbb{R}^p)$. Then

$$\int_{\mathbb{R}^p} \langle T, \varphi(\cdot, z) \rangle dz = \left\langle T, \int_{\mathbb{R}^p} \varphi(\cdot, z) dz \right\rangle$$

Proof.— We start with the case $p = 1$. Let $\varphi \in \mathcal{C}_0^\infty(\Omega \times \mathbb{R})$. We choose $A > 0$ and a compact set $K \subset \Omega$ such that $\text{supp } \varphi \subset K \times [-A, A]$. We denote by $\psi : \Omega \times \mathbb{R} \rightarrow \mathbb{C}$ the function given by

$$\psi(x, z) = \int_{t < z} \varphi(x, t) dt.$$

The function ψ belongs to $\mathcal{C}^\infty(\Omega \times \mathbb{R})$, and for any z , $\text{supp}(\psi)$ is included into $K \times \mathbb{R}$. Therefore Proposition 2.3.17— with Remark 2.3.19— applies. The function

$$G(z) = \langle T, \psi(x, z) \rangle = \left\langle T, \int_{t < z} \varphi(x, t) dt \right\rangle$$

is smooth, and

$$G'(z) = \langle T, \partial_y \psi(x, z) \rangle = \langle T, \varphi(x, z) \rangle.$$

Integrating, we get

$$\left\langle T, \int_{t < z} \varphi(x, t) dt \right\rangle = G(z) = \int_{t < z} G'(t) dt = \int_{t < z} \langle T, \varphi(x, t) \rangle dt .$$

This gives the proposition, taking $z = A$ for example.

For $p > 1$, we proceed by induction on p . Let $\varphi \in \mathcal{C}_0^\infty(\Omega \times \mathbb{R}^p)$. Using the result in the case $p = 1$, we get, for every $z' \in \mathbb{R}^{p-1}$,

$$\left\langle T, \int_{\mathbb{R}} \varphi(x, z', t) dt \right\rangle = \int_{\mathbb{R}} \langle T, \varphi(x, z', t) \rangle dt.$$

It remains to apply the induction assumption to $\tilde{\varphi} \in \mathcal{C}_0^\infty(\Omega \times \mathbb{R}^{p-1})$ defined by

$$\tilde{\varphi}(x, z') = \int_{\mathbb{R}} \varphi(x, z', t) dt .$$

The proof is completed by using the Fubini theorem. □

Finally we come back to the proof of Proposition 2.3.16. Fix $\varphi \in \mathcal{C}_0^\infty(\Omega)$. Applying Proposition 2.3.17 — derivation under the bracket— to

$$\varphi(x) F^\varepsilon(x) = \langle T, \varphi(x) \rho_\varepsilon(x - \cdot) \rangle ,$$

we infer $\varphi F^\varepsilon \in \mathcal{C}^\infty(\Omega)$ and

$$\forall \alpha \in \mathbb{N}^d , \partial^\alpha (\varphi F^\varepsilon)(x) = \langle T, \partial_x^\alpha (\varphi(x) \rho_\varepsilon(x - \cdot)) \rangle .$$

Choosing φ a plateau function near a point $x_0 \in \Omega$, we get the first statement of the proposition,

$$\forall \alpha \in \mathbb{N}^d , \partial^\alpha F^\varepsilon(x_0) = \langle T, \partial^\alpha \rho_\varepsilon(x_0 - \cdot) \rangle .$$

As for the second statement, we apply Proposition 2.3.20 — integration under the bracket — to obtain

$$\int_{\Omega} \varphi(x) F^\varepsilon(x) dx = \langle T, \varphi_\varepsilon \rangle , \quad \varphi_\varepsilon(y) := \int_{\mathbb{R}^d} \varphi(x) \rho_\varepsilon(x - y) dx .$$

Notice that a simple change of variables $x = y + \varepsilon z$ provides

$$\varphi_\varepsilon(y) = \int_{\mathbb{R}^d} \varphi(y + \varepsilon z) \rho(z) dz .$$

Since ρ is supported in the unit ball, we observe that φ_ε is supported in a fixed compact subset of Ω if ε is small enough. Furthermore, the standard derivation under the integral yields $\varphi_\varepsilon \in \mathcal{C}_0^\infty(\Omega)$ and

$$\forall \alpha \in \mathbb{N}^d, \partial^\alpha \varphi_\varepsilon(y) = \int_{\mathbb{R}^d} \partial^\alpha \varphi(y + \varepsilon z) \rho(z) dz .$$

Passing to the limit as ε tends, to 0, $\partial^\alpha \varphi_\varepsilon$ converges uniformly to $\partial^\alpha \varphi$. We conclude that φ_ε converges to φ in $\mathcal{C}_0^\infty(\Omega)$. Consequently,

$$\langle T, \varphi_\varepsilon \rangle \xrightarrow{\varepsilon \rightarrow 0} \langle T, \varphi \rangle .$$

The proof of Proposition 2.3.16 is complete. □

Corollary 2.3.21 Every distribution on Ω is the limit of a sequence of test functions in $\mathcal{D}'(\Omega)$.

Proof.— For every $j \geq 1$, define

$$K_j = \left\{ x \in \Omega : d(x, \Omega^c) \geq \frac{1}{j}, |x| \leq j \right\} .$$

Then K_j is a compact subset of Ω , $K_j \subset \overset{\circ}{K}_{j+1}$ and

$$\bigcup_{j \geq 1} K_j = \Omega .$$

Choose $\chi_j \in \mathcal{C}_0^\infty(\overset{\circ}{K}_{j+1})$ a plateau function near K_j and consider a sequence (ε_j) tending to 0 such that

$$(K_{j+1})_{\varepsilon_j} \subset \Omega .$$

Then define

$$F_j(x) = \chi_j(x) \langle T, \rho_{\varepsilon_j}(x - \cdot) \rangle .$$

By Proposition 2.3.16 and by the above support information, we know that $F_j \in \mathcal{C}_0^\infty(\Omega)$. Furthermore, if $\varphi \in \mathcal{C}_0^\infty(\Omega)$, the support of φ is covered by a finite union of the open subsets $\overset{\circ}{K}_j$'s, hence is contained into some $\overset{\circ}{K}_{j_0}$. Then, for $j \geq j_0$,

$$\int_\omega \varphi(x) F_j(x) dx = \int_\Omega \varphi(x) \langle T, \rho_{\varepsilon_j}(x - \cdot) \rangle dx ,$$

which converges to $\langle T, \varphi \rangle$ by Proposition 2.3.16. Therefore $F_j \rightarrow T$. □

Corollary 2.3.22 Let $T \in \mathcal{D}'(\Omega)$ such that, for every $j \in \{1, \dots, d\}$, $\partial_j T \in \mathcal{C}^0(\Omega)$. Then $T \in \mathcal{C}^1(\Omega)$.

Remark 2.3.23 The proof below is significantly more intricate than its analogue in dimension 1. Indeed, the analogue of the integral formula used in Corollary 1.4.9 in Chapter 1 is much more complicated in several space dimensions, so we prefer to proceed differently. Moreover, let us mention that, on the contrary to the one dimensional case, the assumptions $\partial_j T \in L_{loc}^1(\Omega), j = 1, \dots, d$, do not lead to $T \in \mathcal{C}^0(\Omega)$, but only to $T \in L_{loc}^p(\Omega)$ for $p = \frac{d}{d-1}$ (Sobolev estimates).

Let us prove Corollary 2.3.22. By the gluing principle, it is enough to prove it locally, so we may assume that Ω is a ball B . For every j , denote by $f_j \in C^0(B)$ the continuous function defining $\partial_j T$. As above, we define, on the ball B^ε , the smooth function

$$F^\varepsilon(x) = \langle T, \rho_\varepsilon(x - \cdot) \rangle .$$

Notice that

$$\partial_j F^\varepsilon(x) = \langle T, \partial_{x_j} \rho_\varepsilon(x - \cdot) \rangle = -\langle T, \partial_{y_j} (\rho_\varepsilon(x - \cdot)) \rangle = \langle \partial_j T, \rho_\varepsilon(x - \cdot) \rangle = \int_{\Omega} f_j(y) \rho_\varepsilon(x - y) dy .$$

Fix any ball $B' \subset\subset B$ and $\varepsilon_0 > 0$ such that $B' \subset B^{\varepsilon_0}$. Then the family $(\partial_j F^\varepsilon)_{\varepsilon < \varepsilon_0}$ converges uniformly to f_j on B' as ε tends to 0. Furthermore, picking $\chi \in C_0^\infty(B')$ of integral 1, we have

$$F^\varepsilon(x) - \int_B \chi(y) F^\varepsilon(y) dy = \int_B \chi(y) (F^\varepsilon(x) - F^\varepsilon(y)) dy = \sum_{j=1}^d \int_0^1 \int_B \chi(y) (x_j - y_j) \partial_j F^\varepsilon(y + t(x - y)) dy dt ,$$

and the right hand side converges uniformly on B' . On the other hand, by Proposition 2.3.16,

$$\int_B \chi(y) F^\varepsilon(y) dy \xrightarrow{\varepsilon \rightarrow 0} \langle T, \chi \rangle ,$$

so we infer that F^ε converges uniformly on B' to some function F . Since each $\partial_j F^\varepsilon$ is uniformly convergent on B' , we conclude that $F \in C^1(B')$. But we also know by Proposition 2.3.16 that $T|_{B'} = F$. Since B' is arbitrary, this completes the proof. \square

A very useful property of the convergence of distributions is the following lemma, which is a consequence of the principle of uniform boundedness which is proved at the end of this chapter.

Lemma 2.3.24 (Bicontinuity of the bracket) Let (T_n) be sequence in $\mathcal{D}'(\Omega)$ and (φ_n) be a sequence in $\mathcal{D}(\Omega)$. We assume that $T_n \rightarrow T$ in $\mathcal{D}'(\Omega)$ and $\varphi_n \rightarrow \varphi$ in $\mathcal{D}(\Omega)$. Then

$$\langle T_n, \varphi_n \rangle \rightarrow \langle T, \varphi \rangle .$$

We close this subsection by a useful remark on the gluing principle for convergence of sequences of distributions.

Proposition 2.3.25 Let (T_n) be a sequence of distributions on Ω . Assume

$$\Omega = \bigcup_{j \in J} \Omega_j$$

for some collection $(\Omega_j)_{j \in J}$ of open subsets, and that, for every $j \in J$, $(T_n)|_{\Omega_j}$ converges to some $T^{(j)} \in \mathcal{D}'(\Omega_j)$. Then T_n is convergent in $\mathcal{D}'(\Omega)$.

Proof.— Let K be a compact subset of Ω . From Proposition 2.2.3 let $(\chi_j)_{j \in J}$ be a family of test functions, which are identically 0 except for a finite set of indices j , and such that

$$\text{supp}(\chi_j) \subset \Omega_j , \quad \sum_{j \in J} \chi_j = 1 \text{ on } K .$$

Then, for every test function φ supported in K , we have

$$\langle T_n, \varphi \rangle = \sum_{j \in J} \langle T_n, \chi_j \varphi \rangle \rightarrow \sum_{j \in J} \langle T^{(j)}, \chi_j \varphi \rangle .$$

Since each $T^{(j)}$ is a distribution, the right hand side is estimated by $C \|\varphi\|_{C^m}$ for some $C > 0$ and $m \in \mathbb{N}$ only depending on K . Hence T_n is convergent in $\mathcal{D}'(\Omega)$. \square

2.4 Superficial measures and the jump formula

2.4.1 Motivation

In this section, we come back to the problem of identifying the partial derivatives of the characteristic function of an open set U . It is easy to check that these derivatives are distributions supported by the boundary ∂U of U . Under suitable assumptions on U , we shall show that these distributions are of order 0, and can be expressed in terms of a positive measure supported by ∂U , called the superficial measure of ∂U as an hypersurface. Thus we first need to define the superficial measure on an hypersurface, which is the purpose of the next subsection. An intuitive way to define a superficial measure of a subset of an hypersurface is to thicken this subset with a thickness ε , and to take the limit, as ε tends to 0, of the ratio to ε of the Lebesgue measure of this thickened subset. The next construction tries to make rigorous this intuitive definition.

2.4.2 Preliminaries on non-negative distributions

Definition 2.4.1 We say that $T \in \mathcal{D}'(\Omega)$ is a non-negative distribution when $\langle T, \varphi \rangle \in \mathbb{R}^+$ for any function $\varphi \in \mathcal{C}_0^\infty(\Omega)$ with values in $[0, +\infty)$.

Proposition 2.4.2 If $T \in \mathcal{D}'(\Omega)$ is non-negative, then it is of order 0.

Proof.— Let $K \subset I$ be a compact subset, and $\chi \in \mathcal{C}_0^\infty(I)$ a plateau function above K . For $\varphi \in \mathcal{C}_0^\infty(\Omega)$ supported in K with real values, one has

$$\forall x \in \Omega, -\chi \sup |\varphi| \leq \varphi(x) \leq \chi \sup |\varphi|,$$

thus

$$\langle T, \varphi + \chi \sup |\varphi| \rangle \geq 0 \text{ and } \langle T, \chi \sup |\varphi| - \varphi \rangle \geq 0.$$

This gives

$$|\langle T, \varphi \rangle| \leq \langle T, \chi \sup |\varphi| \rangle.$$

If $\varphi \in \mathcal{C}_0^\infty(\Omega)$ is supported in K and is complex valued, we write $\varphi = \varphi_1 + i\varphi_2$ with φ_1, φ_2 real-valued, and, with what we have seen before,

$$|\langle T, \varphi \rangle| = |\langle T, \varphi_1 + i\varphi_2 \rangle| \leq |\langle T, \varphi_1 \rangle| + |\langle T, \varphi_2 \rangle| \leq C \sup |\varphi_1| + C \sup |\varphi_2| \leq C \sup |\varphi|,$$

and this concludes the proof of the proposition. \square

From Proposition 2.4.2 — and the fact that convenient regularisation processes conserve non-negative functions —, non-negative distributions therefore extends as non-negative linear forms on $\mathcal{C}_0^0(\Omega)$. By the Riesz representation theorem, non-negative linear forms on $\mathcal{C}_0^0(I)$ correspond to positive Borel measures on I which are finite on compact subsets. In other words,

Theorem 2.4.3 For every non-negative distribution T on I there exists such a unique Borel measure μ , finite on compact subsets, such that

$$\forall \varphi \in \mathcal{C}_0^0(\Omega), \langle T, \varphi \rangle = \int_{\Omega} \varphi d\mu .$$

2.4.3 The measure $\delta(f)$ and the superficial measure associated to $\{f = 0\}$.

Let Ω be open subset of \mathbb{R}^d , and let $f : \Omega \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function. We assume the following property,

$$(2.4.1) \quad \forall x \in \Omega, f(x) = 0 \Rightarrow \nabla f(x) \neq 0 .$$

We denote by Σ the set of points $x \in \Omega$ such that $f(x) = 0$.

Theorem 2.4.4 There exists a positive distribution $\delta(f)$, supported by Σ , such that, for every compactly supported function $h \in \mathcal{C}^0(\mathbb{R})$ such that

$$\int_{\mathbb{R}} h(z) dz = 1 ,$$

the family of functions μ_{ε} defined on Ω by

$$\mu_{\varepsilon}(x) := \frac{1}{\varepsilon} h\left(\frac{f(x)}{\varepsilon}\right)$$

converges in $\mathcal{D}'(\Omega)$ to $\delta(f)$ as ε tends to 0. Furthermore, if $g : \Omega \rightarrow \mathbb{R}$ is another \mathcal{C}^1 function which satisfies (2.4.1) and $\Sigma = \{x \in \Omega, g(x) = 0\}$, then

$$(2.4.2) \quad |\nabla g| \delta(g) = |\nabla f| \delta(f) .$$

Proof.— By the gluing principle for convergence of distributions — Proposition 2.3.25—, it is enough to prove that every point a of Ω has an open neighborhood on which μ_{ε} has a limit in the sense of distributions. Moreover, the gluing principle shows that the positivity of this limit near each point a implies that this limit is a positive distribution on Ω — see the remark after Proposition 2.3.8.

Let $a \in \Omega$. If $f(a) \neq 0$, then the continuity of f implies that $|f| \geq c > 0$ on some neighborhood of a , so that, since h is compactly supported in \mathbb{R} , μ_{ε} is identically zero on this neighborhood if ε is small enough. Hence μ_{ε}

tends to 0 in a neighborhood of a .

Now assume $f(a) = 0$. By assumption (2.4.1), $\nabla f(a) \neq 0$, so there exists $j \in \{1, \dots, d\}$ such that $\partial_j f(a) \neq 0$. Let us assume for instance that $j = 1$, and write elements of \mathbb{R}^d as $x = (x_1, y)$, with $x_1 \in \mathbb{R}$ and $y \in \mathbb{R}^{d-1}$. Then, by the implicit function theorem, there exists an open interval $I \subset \mathbb{R}$ and an open subset $W \subset \mathbb{R}^{d-1}$ such that $a = (a_1, b) \in I \times W \subset \Omega$, and a C^1 function $q : W \rightarrow I$ such that

$$\forall (x_1, y) \in I \times W, f(x_1, y) = 0 \iff x_1 = q(y).$$

By the Taylor formula in the x_1 variable, we can write

$$\forall (x_1, y) \in I \times W, f(x_1, y) = f(q(y), y) + (x_1 - q(y)) \int_0^1 \partial_1 f(q(y) + t(x_1 - q(y)), y) dt = m(x_1, y)(x_1 - q(y)),$$

with $m_1(q(y), y) = \partial_1 f(q(y), y) \neq 0$. Given $\varphi \in \mathcal{C}_0^0(I \times W)$, we compute

$$\int_{I \times W} \varphi(x) \mu_\varepsilon(x) dx = \int_W \int_I \varphi(x_1, y) \frac{1}{\varepsilon} h\left(\frac{m(x_1, y)(x_1 - q(y))}{\varepsilon}\right) dx_1 dy.$$

Extending φ by 0 to $\mathbb{R} \times W$, we can write the inner integral as

$$\begin{aligned} \int_{\mathbb{R}} \varphi(x_1, y) \frac{1}{\varepsilon} h\left(\frac{m(x_1, y)(x_1 - q(y))}{\varepsilon}\right) dx_1 &= \int_{\mathbb{R}} \varphi(q(y) + \varepsilon z, y) h(m(q(y) + \varepsilon z, y)z) dz \\ &\rightarrow \left(\int_{\mathbb{R}} h(m(q(y), y)z) dz \right) \varphi(q(y), y) = \frac{\varphi(q(y), y)}{|m(q(y), y)|} \\ &= \frac{\varphi(q(y), y)}{|\partial_1 f(q(y), y)|}. \end{aligned}$$

Passing to the uniform limit under the integral in the y variable, we infer

$$\int_{I \times W} \varphi(x) \mu_\varepsilon(x) dx \rightarrow \int_W \frac{\varphi(q(y), y)}{|\partial_1 f(q(y), y)|} dy.$$

Furthermore, notice that the denominator of the above integrand can be rewritten in terms of $|\nabla f|$ as follows. Taking the derivative with respect to $y \in W$ of the equation $f(q(y), y) = 0$, we obtain, by the chain rule,

$$\nabla_y f(q(y), y) + \partial_1 f(q(y), y) \nabla q(y) = 0,$$

so that

$$\nabla f(q(y), y) = \partial_1 f(q(y), y) \begin{pmatrix} 1 \\ -\nabla q(y) \end{pmatrix}, \quad |\nabla f(q(y), y)| = |\partial_1 f(q(y), y)| (1 + |\nabla q(y)|^2)^{\frac{1}{2}}.$$

Finally, we obtain

$$\int_{I \times W} \varphi(x) \mu_\varepsilon(x) dx \rightarrow \langle \delta(f), \varphi \rangle = \int_W \varphi(q(y), y) \frac{(1 + |\nabla q(y)|^2)^{\frac{1}{2}}}{|\nabla f(q(y), y)|} dy.$$

Thus on W we get a positive distribution $\delta(f)$, hence a distribution of order 0, so we can multiply it by the continuous function $|\nabla f|$, and obtain, if φ is supported into $I \times W$,

$$\langle |\nabla f| \delta(f), \varphi \rangle = \int_W \varphi(q(y), y) (1 + |\nabla q(y)|^2)^{\frac{1}{2}} dy.$$

If g is another \mathcal{C}^1 function which satisfies (2.4.1) and $\Sigma = \{x \in \Omega, g(x) = 0\}$, we take the derivative of $g(q(y), y) = 0$, and obtain

$$\nabla g(q(y), y) = \partial_1 g(q(y), y) \begin{pmatrix} 1 \\ -\nabla q(y) \end{pmatrix}, \quad |\nabla g(q(y), y)| = |\partial_1 g(q(y), y)| (1 + |\nabla q(y)|^2)^{\frac{1}{2}},$$

so that the same computation as above yields

$$\langle |\nabla g| \delta(g), \varphi \rangle = \int_W \varphi(q(y), y) (1 + |\nabla q(y)|^2)^{\frac{1}{2}} dy.$$

In particular, $|\nabla g| \delta(g) = |\nabla f| \delta(f)$ in $I \times W$, and, by the gluing principle,

$$|\nabla g| \delta(g) = |\nabla f| \delta(f)$$

in Ω . □

From this proposition, we infer that the positive distribution $|\nabla f| \delta(f)$ does not depend on the choice of the function f satisfying (2.4.1) and such that $\Sigma = \{f = 0\}$. Therefore this positive distribution is intrinsically associated to Σ .

Definition 2.4.5 The superficial measure of Σ is the positive distribution

$$\sigma = |\nabla f| \delta(f),$$

where f is any \mathcal{C}^1 function satisfying (2.4.1) and such that $\Sigma = \{f = 0\}$.

Let us retain from the above definition and from the proof of Theorem 2.4.4 the following two important facts about the superficial measure on Σ .

i) If $\Sigma = \{f = 0\}$ where f satisfies (2.4.1), then, for every $\varphi \in \mathcal{C}_0^0(\Omega)$,

$$\int_{\Omega} \frac{1}{\varepsilon} h\left(\frac{f(x)}{\varepsilon}\right) \varphi(x) dx \rightarrow \int_{\Omega} \varphi(x) \frac{d\sigma(x)}{|\nabla f(x)|}.$$

ii) If $\Sigma \cap (I \times W) = \{(q(y), y), y \in W\}$ where $q : W \rightarrow I$ is a \mathcal{C}^1 function, then, for every continuous function φ supported into $I \times W$,

$$(2.4.3) \quad \int_{I \times W} \varphi(x) d\sigma(x) = \int_W \varphi(q(y), y) (1 + |\nabla q(y)|^2)^{\frac{1}{2}} dy.$$

Example 2.4.6 (The superficial measure on a sphere) Let $r > 0$. Then the function

$$\begin{aligned} f_r : \mathbb{R}^d \setminus \{0\} &\rightarrow \mathbb{R} \\ x &\mapsto |x| - r \end{aligned}$$

is \mathcal{C}^1 and $f_r(x) = 0$ is the equation of the sphere S_r of radius r centered at 0. Notice that

$$\nabla f_r(x) = \frac{x}{|x|},$$

hence $|\nabla f_r(x)| = 1$. For instance, the intersection of S_r with the open set $]0, +\infty[\times \mathbb{R}^{d-1}$ is given by the equation $x_1 = q(y)$, where

$$q(y) = \sqrt{r^2 - |y|^2}, \quad |y| < r.$$

Notice that

$$\nabla q(y) = \frac{-y}{\sqrt{r^2 - |y|^2}}, \quad 1 + |\nabla q(y)|^2 = \frac{r^2}{r^2 - |y|^2},$$

so that the superficial measure σ_r on S_r is given by $\sigma_r = \delta(f_r)$, and, if φ is compactly supported in $]0, +\infty[\times \mathbb{R}^{d-1}$,

$$\int_{]0, +\infty[\times \mathbb{R}^{d-1}} \varphi(x) d\sigma(x) = \int_{|y| < r} \varphi\left(\sqrt{r^2 - |y|^2}, y\right) \frac{r dy}{\sqrt{r^2 - |y|^2}}.$$

Notice that we have a simple connection between σ_r and σ_1 ,

$$(2.4.4) \quad \forall \varphi \in \mathcal{C}_0^0(\mathbb{R}^d), \quad \int_{\mathbb{R}^d} \varphi(x) d\sigma_r(x) = r^{d-1} \int_{\mathbb{R}^d} \varphi(ry) d\sigma_1(y).$$

Indeed, using the definition,

$$\begin{aligned} \int_{\mathbb{R}^d} \varphi(x) d\sigma_r(x) &= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_{\mathbb{R}^d} h\left(\frac{|x| - r}{\varepsilon}\right) \varphi(x) dx \\ &= \lim_{\varepsilon \rightarrow 0} \frac{r}{\varepsilon} r^{d-1} \int_{\mathbb{R}^d} h\left(\frac{r(|y| - 1)}{\varepsilon}\right) \varphi(ry) dy \\ &= r^{d-1} \int_{\mathbb{R}^d} \varphi(ry) d\sigma_1(y), \end{aligned}$$

using the change of variables $x = ry$ in \mathbb{R}^d and the new small parameter $\varepsilon' = \frac{\varepsilon}{r}$.

Finally, it is easy to check that, if $d = 2$,

$$\int_{\mathbb{R}^2} \varphi(x) d\sigma_r(x) = r \int_0^{2\pi} \varphi(r \cos \theta, r \sin \theta) d\theta.$$

2.4.4 Integration on level sets : the smooth coarea formula

The above definition of the superficial measure easily leads to an important formula of integral calculus. Let $f = \Omega \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that

$$(2.4.5) \quad \forall x \in \Omega, \quad \nabla f(x) \neq 0.$$

It is easy to prove that the set $f(\Omega) := \{f(x), x \in \Omega\}$ is an open subset of \mathbb{R} - hence is an open interval if Ω is connected. Indeed, if $t_0 = f(a)$ with $a \in \Omega$, and if, say, $\partial_1 f(a) \neq 0$, then, on a small neighborhood $I \times W$ of $a = (a_1, b)$, we have $\partial_1 f \neq 0$, hence the function $x_1 \in I \mapsto f(x_1, b)$ is strictly monotone, and consequently its range is an open interval of \mathbb{R} .

For every $t \in f(\Omega)$, denote by Σ_t the set of equation $f(x) = t$, and by σ_t the superficial measure on Σ_t .

Proposition 2.4.7 For every $\varphi \in \mathcal{C}_0^0(\Omega)$,

$$\int_{\Omega} \varphi(x) dx = \int_{f(\Omega)} \left(\int_{\Sigma_t} \varphi(x) \frac{d\sigma_t(x)}{|\nabla f(x)|} \right) dt.$$

Proof.— Fix $h \in \mathcal{C}_0^0(\mathbb{R})$ of integral 1. For every $x \in \Omega$, we observe that, for every $\varepsilon > 0$,

$$1 = \frac{1}{\varepsilon} \int_{\mathbb{R}} h\left(\frac{f(x) - t}{\varepsilon}\right) dt .$$

We plug this identity in the x integral and apply the Fubini theorem,

$$\int_{\Omega} \varphi(x) dx = \int_{\mathbb{R}} \left(\int_{\Omega} \frac{1}{\varepsilon} h\left(\frac{f(x) - t}{\varepsilon}\right) \varphi(x) dx \right) dt .$$

We now pass to the limit inside the inner integral as ε tends to 0. We obtain

$$I_{\varepsilon}(t) = \int_{\Omega} \frac{1}{\varepsilon} h\left(\frac{f(x) - t}{\varepsilon}\right) \varphi(x) dx \rightarrow \begin{cases} 0 & \text{if } t \notin f(\Omega) \\ \int_{\Sigma_t} \varphi(x) \frac{d\sigma_t(x)}{|\nabla f(x)|} & \text{if } t \in f(\Omega) . \end{cases}$$

Then the formula follows from the interversion of the integral in t and of the limit as ε tends to 0. In order to justify this interversion, we apply the dominated convergence theorem. First we notice that, because φ and h are compactly supported, $I_{\varepsilon}(t)$ is supported into a compact subset of \mathbb{R} . So we just have to prove that $I_{\varepsilon}(t)$ is uniformly bounded. By the gluing principle — partitions of unity again —, it is enough to assume that the test function φ is supported in a small open subset V of Ω . Assuming that $\partial_1 f \neq 0$ on V , the implicit function theorem shows that, if $V = I \times W$ is small enough, the equation $f(x_1, y) = t$ in V , for t close to some value t_0 , is equivalent to $x_1 = q(y, t)$, where q is a \mathcal{C}^1 function. Then we can reproduce the proof of Theorem 2.4.4 with the additional parameter t , writing

$$\begin{aligned} I_{\varepsilon}(t) &= \int_W \int_I \frac{1}{\varepsilon} h\left(\frac{m(x_1, y, t)(x_1 - q(y, t))}{\varepsilon}\right) \varphi(x_1, y) dy dx_1 \\ &= \int_W \int_{\mathbb{R}} h(m(q(y, t) + \varepsilon z, y)z) \varphi(q(y, t) + \varepsilon z, y) dz dy , \end{aligned}$$

which is uniformly bounded since φ is compactly supported, and $|m| \geq c > 0$. This completes the proof. \square

Corollary 2.4.8 (Integration on spherical level sets) For every $\varphi \in \mathcal{C}_0^0(\mathbb{R}^d \setminus \{0\})$, we have

$$\int_{\mathbb{R}^d \setminus \{0\}} \varphi(x) dx = \int_0^{\infty} \int_{S_1} \varphi(r\omega) d\sigma_1(\omega) r^{d-1} dr .$$

Proof.— Combine Proposition 2.4.7 applied to $f : x \in \mathbb{R}^d \setminus \{0\} \mapsto |x|$ with the change of variable formula (2.4.4). \square

2.4.5 Superficial measure on a closed hypersurface and the jump formula for a regular open subset

We first need a number of definitions.

Definition 2.4.9 Let Ω be an open subset of \mathbb{R}^d . A closed \mathcal{C}^1 hypersurface of Ω is a closed subset $\Sigma \subset \Omega$ with the following property : every $a \in \Sigma$ has an open neighborhood V in Ω such that there exists a \mathcal{C}^1 function $f : V \rightarrow \mathbb{R}$ with

$$\Sigma \cap V = \{x \in V, f(x) = 0\} , \quad \forall x \in \Sigma \cap V, \nabla f(x) \neq 0 .$$

Essentially this definition is the one we used in the previous subsection, except that one does not impose that the existence of a *global* equation f for Σ . This naturally leads to the following

Proposition 2.4.10 Let Σ be a closed \mathcal{C}^1 hypersurface of Ω . There exists a unique positive distribution σ supported in Σ such that, for every open subset $V \subset \Omega$ and $f \in \mathcal{C}^1(V, \mathbb{R})$ such that

$$\Sigma \cap V = \{x \in V, f(x) = 0\}, \quad \forall x \in \Sigma \cap V, \nabla f(x) \neq 0,$$

we have, in V ,

$$|\nabla f| \delta(f) = \sigma.$$

Furthermore, $\text{supp } \sigma = \Sigma$.

Proof.— Existence and uniqueness of σ follow from Theorem 2.4.4 and from the gluing principle. Indeed, given $a \in \Sigma$, consider $V_a \subset \Omega$ and $f_a : V_a \rightarrow \mathbb{R}$ as in the definition above. We define $\sigma_a \in \mathcal{D}'(V_a)$ as

$$\sigma_a = |\nabla f_a| \delta(f_a).$$

Moreover, we consider $\tilde{\sigma} = 0$ in $\mathcal{D}'(\Omega \setminus \Sigma)$. By Theorem 2.4.4 and the gluing principle applied to the open covering

$$\Omega = (\Omega \setminus \Sigma) \cup \bigcup_{a \in \Sigma} V_a,$$

we infer the existence of $\sigma \in \mathcal{D}'(\Omega)$. Uniqueness also follows from the gluing principle and Theorem 2.4.4. The last statement follows from the local expression of σ given in (2.4.3). Notice that the closedness of Σ is crucial in this proof. \square

Definition 2.4.11 Measure σ is called the superficial measure on the closed hypersurface Σ .

We now come to the important notion of a regular open set.

Definition 2.4.12 Let U be an open subset of Ω . We say that U is a regular open subset of Ω of class \mathcal{C}^1 if every point $a \in \partial U$ has an open neighborhood V in Ω such that there exists a \mathcal{C}^1 function $f : V \rightarrow \mathbb{R}$ with

$$U \cap V = \{x \in V, f(x) > 0\}, \quad \forall x \in \partial U \cap V, \nabla f(x) \neq 0.$$

In such a situation, it is easy to check that ∂U is a closed \mathcal{C}^1 hypersurface of Ω . Indeed, with the notation of the above definition, one verifies that

$$\partial U \cap V = \{x \in V, f(x) = 0\}.$$

However, a regular open set is not only an open set whose boundary is a closed hypersurface. The above definition also imposes that the open subset U is locally *on the same side* of this hypersurface! For instance $U = \mathbb{R}^d \setminus \{x_1 = 0\}$ is not a regular subset of \mathbb{R}^d , though its boundary is the closed hypersurface $\{x_1 = 0\}$.

Definition 2.4.13 Let Ω be an open subset of \mathbb{R}^d .

- If Σ is a closed \mathcal{C}^1 hypersurface of Ω and $a \in \Sigma$, the normal line to Σ at a is the line $\mathbb{R}\nabla f(a)$, where f is any function as in Definition 2.4.9.
- If U is a regular open subset of Ω of class \mathcal{C}^1 , the inward unit normal vector to U at $a \in \partial U$ is the unit vector

$$N^{\text{int}}(a) = \frac{\nabla f(a)}{|\nabla f(a)|},$$

where f is any function as in Definition 2.4.12. The outward unit normal vector to U at $a \in \partial U$ is

$$N^{\text{ext}}(a) = -N^{\text{int}}(a).$$

These definitions make sense because of the independence of the objects with respect to function f . In the case of a closed hypersurface, we already checked, in the proof of Theorem 2.4.4, that the vectors $\nabla f(a)$ are all proportional, so they define the same line. In the case of a regular open subset, there exist only two unit vectors on the normal line to ∂U at $a \in \partial U$, which can be identified as follows. For $\varepsilon > 0$ small enough, $a + \varepsilon N^{\text{int}}(a) \in U$, while $a + \varepsilon N^{\text{ext}}(a) \in (\overline{U})^c$.

Finally, notice that the mapping $N^{\text{int}} : \partial U \rightarrow \mathbb{R}^d$ is continuous. We call it the *inward unit normal vector field* to U . A similar definition holds for the *outward unit normal vector field* to U .

Theorem 2.4.14 (The jump formula) Let U be a regular open subset of Ω of class \mathcal{C}^1 . Denote by σ the superficial measure on the closed hypersurface ∂U , and by $N^{\text{int}} : \partial U \rightarrow \mathbb{R}^d$ the inward unit normal vector field to U . Then, in $\mathcal{D}'(\Omega)$,

$$\forall j \in \{1, \dots, d\}, \partial_j(\mathbf{1}_U) = N_j^{\text{int}} \sigma.$$

Proof.— By the gluing principle, it is enough to prove this identity near every point $a \in \Omega$. If $a \notin \partial U$, then $\mathbf{1}_U$ is a constant function in a neighborhood of a , so that $\partial_j(\mathbf{1}_U) = 0$ in this neighborhood, which is also the case of σ .

Assume $a \in \partial U$, and let $f : V \rightarrow \mathbb{R}$ be a function near a as in Definition 2.4.12. Denote by $\chi : \mathbb{R} \rightarrow \mathbb{R}$ a \mathcal{C}^1 function such that

$$\chi(z) = \begin{cases} 0 & \text{if } z < 0 \\ 1 & \text{if } z \geq 0. \end{cases}$$

Then it is easy to check that

$$\forall x \in V, \mathbf{1}_U(x) = \lim_{\varepsilon \rightarrow 0} \chi\left(\frac{f(x)}{\varepsilon}\right).$$

Furthermore, since the right hand side is uniformly bounded, the dominated convergence theorem implies that this convergence also holds in $\mathcal{D}'(V)$. Therefore, in $\mathcal{D}'(V)$, we have

$$\partial_j(\mathbf{1}_U) = \lim_{\varepsilon \rightarrow 0} \partial_j \left(\chi\left(\frac{f(x)}{\varepsilon}\right) \right) = \lim_{\varepsilon \rightarrow 0} \frac{\partial_j f}{\varepsilon} \chi' \left(\frac{f(x)}{\varepsilon} \right).$$

The function $h = \chi'$ belongs to $\mathcal{C}_0^0(\mathbb{R})$ and satisfies

$$\int_{\mathbb{R}} h(z) dz = \int_0^\infty \chi'(z) dz = 1 .$$

Consequently, for every $\varphi \in \mathcal{C}_0^0(V)$,

$$\int_{\Omega} \varphi(x) \frac{1}{\varepsilon} \chi' \left(\frac{f(x)}{\varepsilon} \right) dx \rightarrow \langle \delta(f), \varphi \rangle ,$$

so that

$$\partial_j(\mathbf{1}_U) = \partial_j f \delta(f) = \frac{\partial_j f}{|\nabla f|} \sigma = N_j^{\text{int}} \sigma .$$

□

Corollary 2.4.15 (The Gauss–Green formula) Let U be regular open subset of class \mathcal{C}^1 in Ω . For every $\varphi \in \mathcal{C}_0^1(\Omega)$,

$$\forall j \in \{1, \dots, d\} , \int_U \partial_j \varphi(x) dx = \int_{\partial U} N_j^{\text{ext}}(x) \varphi(x) d\sigma(x) .$$

Proof.— First assume $\varphi \in \mathcal{C}_0^\infty(\Omega)$. In $\mathcal{D}'(\Omega)$, we apply the jump formula and get

$$\int_U \partial_j \varphi(x) dx = -\langle \partial_j(\mathbf{1}_U), \varphi \rangle = -\langle N_j^{\text{int}} \sigma, \varphi \rangle = \int_{\partial U} N_j^{\text{ext}}(x) \varphi(x) d\sigma(x) .$$

The general case $\varphi \in \mathcal{C}_0^1(\Omega)$ follows by approximation. □

Let us come to the special case where U is a bounded regular open subset \mathbb{R}^d . In this case, we denote by $\mathcal{C}^1(\bar{U})$ the space of restrictions to U of \mathcal{C}^1 functions on \mathbb{R}^d .

Corollary 2.4.16 (The Gauss–Green formula, compact form) Let U be a regular open subset of class \mathcal{C}^1 in \mathbb{R}^d such that \bar{U} is compact . For every $\varphi \in \mathcal{C}^1(\bar{U})$,

$$\forall j \in \{1, \dots, d\} , \int_U \partial_j \varphi(x) dx = \int_{\partial U} N_j^{\text{ext}}(x) \varphi(x) d\sigma(x) .$$

Proof.—

If $\varphi \in \mathcal{C}^1(\bar{U})$, let $\tilde{\varphi} \in \mathcal{C}^1(\tilde{U})$ be an extension of φ to an open neighborhood \tilde{U} of \bar{U} . Let χ be a plateau function on \bar{U} compactly supported in \tilde{U} . The first formula follows from applying the Gauss–Green formula to $\chi \tilde{\varphi} \in \mathcal{C}_0^1(\mathbb{R}^d)$. □

Remark 2.4.17 Both corollaries above combined with the Leibniz formula imply the following multidimensional integration by parts formula.

$$\forall j \in \{1, \dots, d\} , \int_U \partial_j \varphi(x) \psi(x) dx = \int_{\partial U} N_j^{\text{ext}}(x) \varphi(x) \psi(x) d\sigma(x) - \int_U \varphi(x) \partial_j \psi(x) dx .$$

2.5 Sobolev spaces in an open set

We recall that, from now on, we identify a locally integrable function f with the associated distribution T_f .

2.5.1 Definition and general facts

Definition 2.5.1 Let $\Omega \subset \mathbb{R}^d$ be an open set, and $s \in \mathbb{N}$. A distribution $u \in \mathcal{D}'(\Omega)$ belongs to $H^s(\Omega)$ if for all $\alpha \in \mathbb{N}^d$ such that $|\alpha| \leq s$, it holds that $\partial^\alpha u \in L^2(\Omega)$. We denote by $(\cdot | \cdot)_{H^s}$ the sesquilinear form defined on $H^s(\Omega) \times H^s(\Omega)$ by

$$(u|v)_{H^s} = \sum_{|\alpha| \leq s} (\partial^\alpha u | \partial^\alpha v)_{L^2} .$$

We also introduce the associated H^s norm

$$\|u\|_{H^s} = (u|u)_{H^s}^{\frac{1}{2}} = \left(\sum_{|\alpha| \leq s} \|\partial^\alpha u\|_{L^2}^2 \right)^{\frac{1}{2}} ,$$

so that convergence of u_n to u in H^s is equivalent to the convergence of $\partial^\alpha u_n$ to $\partial^\alpha u$ in L^2 for every $|\alpha| \leq s$.

Notice that $H^0(\Omega) = L^2(\Omega)$. The structure of Hilbert space of $L^2(\Omega)$ is transferred to $H^s(\Omega)$, as shown by the next proposition.

Proposition 2.5.2 The sesquilinear form $(\cdot, \cdot)_{H^s}$ is a Hermitian scalar product, which makes $H^s(\Omega)$ a Hilbert space.

Proof.— The only non trivial fact is completeness of $H^s(\Omega)$ with respect to the H^s norm. Let (u_j) be a Cauchy sequence in $H^s(\Omega)$. For all $|\alpha| \leq s$, the sequence $(\partial^\alpha u_j)$ is Cauchy in L^2 , thus converges to a $v_\alpha \in L^2$. In particular, $u_j \rightarrow v_0$ in $\mathcal{D}'(\Omega)$, so that $\partial^\alpha u_j \rightarrow \partial^\alpha v_0 = v_\alpha \in L^2(\Omega)$, and $(u_j) \rightarrow v_0$ in $H^s(\Omega)$. \square

Notice that, for $s \geq 1$, $u \in H^s(\Omega)$ if and only if $u \in H^1(\Omega)$ and, for every $j = 1, \dots, d-1$, $\partial_j u \in H^{s-1}(\Omega)$. By induction on s , this reduces many properties of $H^s(\Omega)$ to the special case $s = 1$, on which we are going to focus in what follows.

2.5.2 Variational formulation of some elliptic problems

Definition 2.5.3 Given an arbitrary open subset Ω of \mathbb{R}^d , we denote by $H_0^1(\Omega)$ the closure of $\mathcal{C}_0^\infty(\Omega)$ in $H^1(\Omega)$.

Note that $H_0^1(\Omega)$ is a closed subspace of the Hilbert space $H^1(\Omega)$, hence it is a Hilbert space with the inner product $(\cdot | \cdot)_{H^1}$ defined in the previous section.

In what follows, we are going to use the Laplacian differential operator, defined on $\mathcal{D}'(\Omega)$ by

$$\Delta T = \sum_{j=1}^d \partial_j^2 T .$$

Theorem 2.5.4 Let Ω be an arbitrary open subset of \mathbb{R}^d . For every $f \in L^2(\Omega)$, there exists $u \in H_0^1(\Omega)$ unique such that

$$(2.5.6) \quad -\Delta u + u = f$$

in $\mathcal{D}'(\Omega)$.

Proof.— Equation (2.5.6) in $\mathcal{D}'(\Omega)$ exactly means

$$\forall \varphi \in \mathcal{C}_0^\infty(\Omega) , \langle -\Delta u + u, \varphi \rangle = \int_{\Omega} f \varphi \, dx .$$

Since we are looking for $u \in H_0^1(\Omega)$, we know in particular that $u \in L^2(\Omega)$ and $\partial_j u \in L^2(\Omega)$ for $j = 1, \dots, d$. Therefore the left hand side reads

$$\langle -\Delta u + u, \varphi \rangle = \sum_{j=1}^d \int_{\Omega} \partial_j u \partial_j \varphi \, dx + \int_{\Omega} u \varphi \, dx = (\varphi | \bar{u})_{H^1} .$$

Summing up, we are looking for some $u \in H_0^1(\Omega)$ such that

$$\forall \varphi \in \mathcal{C}_0^\infty(\Omega) , (\varphi | \bar{u})_{H^1} = \int_{\Omega} f \varphi \, dx .$$

Since both sides of the above equation are linear forms of φ which are continuous for the H^1 norm, the problem is equivalent to finding $u \in H_0^1(\Omega)$ such that

$$\forall v \in H_0^1(\Omega) , (v | \bar{u}) = \int_{\Omega} f v \, dx .$$

By the Riesz representation theorem, the continuous linear form

$$v \in H_0^1(\Omega) \mapsto \int_{\Omega} f v \, dx$$

can be represented by the inner product with a unique element $w \in H_0^1(\Omega)$. Therefore the problem is solved by setting $u = \bar{w}$ — notice that the conjugate of a function in $H_0^1(\Omega)$ is still in $H_0^1(\Omega)$. \square Let us complete this subsection by some remarks about the possible extensions of Theorem 2.5.4

Remark 2.5.5 *i)* The L^2 function f can be replaced more generally by any distribution which extends as a continuous linear form on $H_0^1(\Omega)$. The space of such distributions is denoted by $H^{-1}(\Omega)$.

ii) If $q \in L^\infty(\Omega)$ and there exists $m > 0$ such that

$$q(x) \geq m$$

almost everywhere on Ω , the sesquilinear form

$$(2.5.7) \quad (u|v)_q = \sum_{j=1}^d (\partial_j u | \partial_j v)_{L^2} + \int_{\Omega} q u \bar{v} \, dx$$

is an inner product on $H^1(\Omega)$, inducing a norm which is equivalent to the H^1 norm. Therefore $H^1(\Omega)$ and $H_0^1(\Omega)$ are Hilbert spaces for this new inner product, and one can solve similarly the equation

$$(2.5.8) \quad -\Delta u + qu = f$$

in $\mathcal{D}'(\Omega)$, for every $f \in H^{-1}(\Omega)$.

iii) The above assumption on q can be relaxed for special cases of open sets Ω , in particular if Ω is bounded.

Lemma 2.5.6 Let Ω be a bounded open subset of \mathbb{R}^d . For every $j \in \{1, \dots, d\}$, there exists $C > 0$ such that, for every $\varphi \in H_0^1(\Omega)$,

$$\|\varphi\|_{L^2} \leq C \|\partial_j \varphi\|_{L^2}.$$

By density, it is enough to prove this inequality for $\varphi \in C_0^\infty(\Omega)$. Let us prove Lemma 2.5.6 with $j = 1$, say. We write $x = (x_1, y) \in \mathbb{R} \times \mathbb{R}^d$ and

$$|\varphi(x_1, y)|^2 = - \int_{x_1}^{\infty} \partial_t (|\varphi(t, y)|^2) \, dt = -2 \int_{x_1}^{\infty} \operatorname{Re}(\partial_t \varphi(t, y) \bar{\varphi}(t, y)) \, dt.$$

Using the Cauchy–Schwarz inequality, we infer

$$|\varphi(x_1, y)|^2 \leq 2 \left(\int_{\mathbb{R}} |\partial_t \varphi(t, y)|^2 \, dt \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}} |\varphi(t, y)|^2 \, dt \right)^{\frac{1}{2}}.$$

Assuming $\Omega \subset]a, a + L[\times \mathbb{R}^{d-1}$, we conclude, again by Cauchy–Schwarz, that

$$\int_{\mathbb{R} \times \mathbb{R}^{d-1}} |\varphi(x_1, y)|^2 \, dx_1 \, dy \leq 2L \|\partial_t \varphi\|_{L^2} \|\varphi\|_{L^2}.$$

This completes the proof of the lemma, with $C = 2L$.

In view of Lemma 2.5.6, if Ω is bounded, the sesquilinear form $(\cdot | \cdot)_q$ defined in (2.5.7) is still an inner product on $H_0^1(\Omega)$, with a norm equivalent to the H^1 norm, if $q \in L^\infty(\Omega)$ is just non–negative — including q identically 0. Hence, under this more general assumption, it is still possible to solve equation (2.5.8) on Ω bounded. In particular, if Ω is bounded, for every $f \in H^{-1}(\Omega)$, there exists $u \in H_0^1(\Omega)$ unique such that

$$-\Delta u = f$$

in $\mathcal{D}'(\Omega)$. We will come back to this equation at the end of this paragraph, in the particular case where Ω is regular. We will see that, in this case, the subspace $H_0^1(\Omega)$ of $H^1(\Omega)$ can be described in a more explicit way.

2.5.3 Approximation by smooth functions

Let us start with the special case $\Omega = \mathbb{R}^d$.

Proposition 2.5.7 $C_0^\infty(\mathbb{R}^d)$ is dense into $H^1(\mathbb{R}^d)$. In other words, $H_0^1(\mathbb{R}^d) = H^1(\mathbb{R}^d)$.

Proof.— We proceed by cut off and regularisation. Let $\chi \in C_0^\infty(\mathbb{R}^d)$ such that $\chi(x) = 1$ on the unit ball. For $n \geq 1$, set

$$\chi_n(x) = \chi\left(\frac{x}{n}\right).$$

Given $v \in L^2(\mathbb{R}^d)$, we already know that $\chi_n v$ tends to v in L^2 as $n \rightarrow \infty$. Let $u \in H^1(\mathbb{R}^d)$. Let us prove that $\chi_n u \rightarrow u$ in H^1 as $n \rightarrow \infty$. It is equivalent to establish that $\chi_n u \rightarrow u$ in L^2 , which is already known, and that for every $j = 1, \dots, d$, $\partial_j(\chi_n u) \rightarrow \partial_j u$. From the Leibniz formula for distributions,

$$\partial_j(\chi_n u) = \chi_n \partial_j u + (\partial_j \chi_n) u,$$

so we just have to prove that $(\partial_j \chi_n) u \rightarrow 0$ in L^2 . This follows from $u \in L^2$ and

$$\|\partial_j \chi_n\|_\infty = O\left(\frac{1}{n}\right).$$

As a second step, we prove that, if $u \in H^1(\mathbb{R}^d)$ is compactly supported, then u can be approximated by a sequence of compactly supported smooth functions. Let $\rho \in C_0^\infty(\mathbb{R}^d)$, supported in the unit ball B , and such that

$$\int_{\mathbb{R}^d} \rho(z) dz = 1.$$

Consider, for $\varepsilon > 0$,

$$u_\varepsilon(x) = \rho_\varepsilon * u(x) = \int_{\mathbb{R}^d} \frac{1}{\varepsilon^d} \rho\left(\frac{x-y}{\varepsilon}\right) u(y) dy.$$

Then $u_\varepsilon \in C_0^\infty(\mathbb{R}^d)$, with $\text{supp}(u_\varepsilon) \subset \text{supp}(u) + \varepsilon B$. We already know that $u_\varepsilon \rightarrow u$ in L^2 as $\varepsilon \rightarrow 0$. Furthermore, for $j = 1, \dots, d$,

$$\partial_j u_\varepsilon(x) = \int_{\mathbb{R}^d} \partial_j \rho_\varepsilon(x-y) u(y) dy = - \int_{\mathbb{R}^d} \partial_{y_j} [\rho_\varepsilon(x-y)] u(y) dy = \rho_\varepsilon * \partial_j u(x).$$

Therefore $\partial_j u_\varepsilon \rightarrow \partial_j u$ as $\varepsilon \rightarrow 0$. Summing up, $u_\varepsilon \rightarrow u$ in H^1 as $\varepsilon \rightarrow 0$, and the proof is complete. \square Notice that the second step of the above proof implies the following useful result.

Proposition 2.5.8 Every compactly supported element u of $H^1(\Omega)$ is the limit of a sequence of $C_0^\infty(\Omega)$ supported in an arbitrarily small neighbourhood of $\text{supp}(u)$.

Next we deal with the main result of this subsection, which concerns the special case of a bounded regular open subset of \mathbb{R}^d .

Proposition 2.5.9 Let Ω be a bounded regular open subset of \mathbb{R}^d with a \mathcal{C}^∞ boundary. Then $\mathcal{C}^\infty(\overline{\Omega})$ is dense in $H^1(\Omega)$.

Proof.— Given $a \in \partial\Omega$, there exists an open neighbourhood V_a of a such that $\Omega \cap V_a$ is the strict epigraph of a smooth function. Since the V_a cover the compact subset $\partial\Omega$ of \mathbb{R}^d , we may extract a finite covering

$$\partial\Omega \subset \bigcup_{k=1}^N V_k$$

and we consider a partition of unity χ_0, \dots, χ_N associated to the covering

$$\overline{\Omega} \subset \Omega \cup \bigcup_{k=1}^N V_k,$$

so that $\chi_0 \in \mathcal{C}_0^\infty(\Omega)$, $\chi_k \in \mathcal{C}_0^\infty(V_k)$ for $k = 1, \dots, N$, and

$$\chi_0 + \chi_1 + \dots + \chi_N = 1$$

on $\overline{\Omega}$. In particular, on Ω , we have

$$u = \chi_0 u + \sum_{k=1}^N \chi_k u.$$

We may apply Proposition 2.5.8 to $\chi_0 u$. Therefore it remains to prove that, for every $k = 1, \dots, N$, $\chi_k u$ can be approximated in $H^1(\Omega)$ by a sequence of $\mathcal{C}^\infty(\overline{\Omega})$.

We drop the index k for simplicity, and we consider an open subset $V = I \times W$ of $\mathbb{R}^d = \mathbb{R}_{x_1} \times \mathbb{R}_y^{d-1}$ and a smooth function $q : W \rightarrow \mathbb{R}$ such that

$$V \cap \Omega = \{(x_1, y) \in I \times W, x_1 > q(y)\},$$

and $\chi \in \mathcal{C}_0^\infty(V)$. For $(z, y) \in]0, \infty[\times W$, we set

$$v(z, y) = (\chi u)(z + q(y), y).$$

Notice that, for some compact subset K of W and some $a > 0$,

$$\text{supp}(v) \subset]0, a] \times K.$$

We need a couple of results concerning this function v .

Lemma 2.5.10 The function v belongs to $H^1(]0, \infty[\times W)$, with

$$\partial_z v(z, y) = \partial_z (\chi u)(z + q(y), y), \quad \partial_{y_j} v(z, y) = \partial_{y_j} (\chi u)(z + q(y), y) + \partial_{y_j} q(y) \partial_z (\chi u)(z + q(y), y).$$

Let us prove Lemma 2.5.10. First of all, $v \in L^2(]0, \infty[\times W)$, since, by the change of variables formula,

$$\int_W \int_0^\infty |v(z, y)|^2 dz dy = \int_W \int_{q(y)}^\infty |(\chi u)(x_1, y)|^2 dx_1 dy < +\infty.$$

Let $\varphi \in C_0^\infty(]0, \infty[\times W)$. Let us calculate, using the fact that $\chi u \in H^1(\Omega)$,

$$\begin{aligned} - \int_W \int_0^\infty v(z, y) \partial_z \varphi(z, y) dz dy &= - \int_W \int_{q(y)}^\infty (\chi u)(x_1, y) \partial_z \varphi(x_1 - q(y), y) dx_1 dy \\ &= - \int_\Omega (\chi u)(x_1, y) \frac{\partial}{\partial x_1} [\varphi(x_1 - q(y), y)] dx_1 dy \\ &= \int_\Omega \frac{\partial}{\partial x_1} [(\chi u)(x_1, y)] \varphi(x_1 - q(y), y) dx_1 dy \\ &= \int_W \int_0^\infty \partial_{x_1} (\chi u)(z + q(y), y) \varphi(z, y) dz dy . \end{aligned}$$

Similarly,

$$\begin{aligned} - \int_W \int_0^\infty v(z, y) \partial_{y_j} \varphi(z, y) dz dy &= - \int_W \int_{q(y)}^\infty (\chi u)(x_1, y) \partial_{y_j} \varphi(x_1 - q(y), y) dx_1 dy \\ &= - \int_\Omega (\chi u)(x_1, y) \frac{\partial}{\partial y_j} [\varphi(x_1 - q(y), y)] dx_1 dy \\ &\quad - \int_\Omega (\chi u)(x_1, y) \partial_{y_j} q(y) \frac{\partial}{\partial x_1} [\varphi(x_1 - q(y), y)] dx_1 dy \\ &= \int_\Omega [\partial_{y_j} + \partial_{y_j} q(y) \partial_{x_1}] (\chi u)(x_1, y) \varphi(x_1 - q(y), y) dx_1 dy \\ &= \int_W \int_0^\infty [\partial_{y_j} + \partial_{y_j} q(y) \partial_{x_1}] (\chi u)(z + q(y), y) \varphi(z, y) dz dy . \end{aligned}$$

Since the functions

$$\begin{aligned} (z, y) \in]0, \infty[\times W &\mapsto \partial_{x_1} (\chi u)(z + q(y), y) \\ (z, y) \in]0, \infty[\times W &\mapsto [\partial_{y_j} + \partial_{y_j} q(y) \partial_{x_1}] (\chi u)(z + q(y), y) \end{aligned}$$

belong to $L^2(]0, \infty[\times W)$ by change of variables again, this completes the proof of Lemma 2.5.10.

Lemma 2.5.11 Given $v \in H^1(]0, \infty[\times W)$, define $\tilde{v} \in L^2(\mathbb{R} \times W)$ by

$$\tilde{v}(z, y) = \begin{cases} v(z, y) & \text{if } z > 0 \\ v(-z, y) & \text{if } z < 0 \end{cases}$$

Then $\tilde{v} \in H^1(\mathbb{R} \times W)$.

Let us prove Lemma 2.5.11. First we observe that

$$\|\tilde{v}\|_{L^2(\mathbb{R} \times W)}^2 = 2\|v\|_{L^2(]0, \infty[\times W)}^2 < \infty$$

so $\tilde{v} \in L^2(\mathbb{R} \times W)$ indeed. Let $\varphi \in C_0^\infty(\mathbb{R} \times W)$. Let us calculate

$$\begin{aligned} - \int_{\mathbb{R}} \int_W \tilde{v}(z, y) \partial_{y_j} \varphi(z, y) dz dy &= - \int_0^{+\infty} \int_W v(z, y) \partial_{y_j} \varphi(z, y) dz dy - \int_{-\infty}^0 \int_W v(-z, y) \partial_{y_j} \varphi(z, y) dz dy \\ &= - \int_0^\infty \int_W v(z, y) [\partial_{y_j} \varphi(z, y) + \partial_{y_j} \varphi(-z, y)] dz dy \end{aligned}$$

In order to use the information $v \in H^1(]0, \infty[\times W)$, we need to reduce to test functions compactly supported in $]0, \infty[\times W$. We introduce $\chi \in C^\infty(\mathbb{R})$ such that $\chi(t) = 0$ for $t \leq \frac{1}{2}$ and $\chi(t) = 1$ for $t \geq 1$, and we set, for $\varepsilon > 0$,

$$\chi_\varepsilon(z) = \chi\left(\frac{z}{\varepsilon}\right).$$

The $\chi_\varepsilon(z) \rightarrow \mathbf{1}_{]0, \infty[}(z)$ as $\varepsilon \rightarrow 0$, and

$$\begin{aligned} - \int_{\mathbb{R}} \int_W \tilde{v}(z, y) \partial_{y_j} \varphi(z, y) dz dy &= - \lim_{\varepsilon \rightarrow 0} \int_0^\infty \int_W v(z, y) \partial_{y_j} [\chi_\varepsilon(z) (\varphi(z, y) + \varphi(-z, y))] dz dy \\ &= \lim_{\varepsilon \rightarrow 0} \int_0^\infty \int_W \partial_{y_j} v(z, y) \chi_\varepsilon(z) (\varphi(z, y) + \varphi(-z, y)) dz dy \\ &= \int_0^\infty \int_W \partial_{y_j} v(z, y) (\varphi(z, y) + \varphi(-z, y)) dz dy \\ &= \int_{\mathbb{R}} \int_W f_j(z, y) \varphi(z, y) dz dy, \end{aligned}$$

where

$$f_j(z, y) := \begin{cases} \partial_{y_j} v(z, y) & \text{if } z > 0 \\ \partial_{y_j} v(-z, y) & \text{if } z < 0 \end{cases}$$

defines an L^2 function on $\mathbb{R} \times W$. Similarly,

$$\begin{aligned} - \int_{\mathbb{R}} \int_W \tilde{v}(z, y) \partial_z \varphi(z, y) dz dy &= - \int_0^{+\infty} \int_W v(z, y) \partial_z \varphi(z, y) dz dy - \int_{-\infty}^0 \int_W v(-z, y) \partial_z \varphi(z, y) dz dy \\ &= - \int_0^\infty \int_W v(z, y) \partial_z [\varphi(z, y) - \varphi(-z, y)] dz dy \\ &= - \lim_{\varepsilon \rightarrow 0} \int_0^\infty \int_W v(z, y) \partial_z [\chi_\varepsilon(z) (\varphi(z, y) - \varphi(-z, y))] dz dy \\ &\quad + \lim_{\varepsilon \rightarrow 0} \int_0^\infty \int_W v(z, y) \chi'_\varepsilon(z) (\varphi(z, y) - \varphi(-z, y)) dz dy \\ &= \lim_{\varepsilon \rightarrow 0} \int_0^\infty \int_W \partial_z v(z, y) \chi_\varepsilon(z) (\varphi(z, y) - \varphi(-z, y)) dz dy \\ &\quad + \lim_{\varepsilon \rightarrow 0} O\left(\frac{1}{\varepsilon}\right) \int_{\frac{\varepsilon}{2}}^\varepsilon |v(z, y)| O(z) dz dy \\ &= \int_0^\infty \int_W \partial_z v(z, y) (\varphi(z, y) - \varphi(-z, y)) dz dy \\ &= \int_{\mathbb{R}} \int_W g(z, y) \varphi(z, y) dz dy, \end{aligned}$$

where where

$$g(z, y) := \begin{cases} \partial_z v(z, y) & \text{if } z > 0 \\ -\partial_z v(-z, y) & \text{if } z < 0 \end{cases}$$

defines an L^2 function on $\mathbb{R} \times W$. Finally, we get $\tilde{v} \in H^1(\mathbb{R} \times W)$ with

$$\partial_{y_j} \tilde{v} = f_j, \quad \partial_z \tilde{v} = g.$$

This completes the proof of Lemma 2.5.11.

Let us complete the proof of Proposition 2.5.9. Applying Lemma 2.5.11 to

$$v(z, y) = (\chi u)(z + q(y), y),$$

we observe that $\tilde{v} \in H^1(\mathbb{R} \times W)$ and $\text{supp}(\tilde{v}) \subset [-a, a] \times K$, a compact subset of $\mathbb{R} \times W$. Now we appeal to Proposition 2.5.8 and obtain that there exist $\tilde{v}_\varepsilon \in C_0^\infty(\mathbb{R} \times W)$ such that $\text{supp}(\tilde{v}_\varepsilon) \subset [-(a + \varepsilon, a + \varepsilon) \times K_\varepsilon$ and $\tilde{v}_\varepsilon \rightarrow \tilde{v}$ in $H^1(\mathbb{R} \times W)$. Then the restriction v_ε of \tilde{v}_ε to $]0, +\infty[\times W$ converge to v in $H^1(]0, \infty[\times W)$, and

$$u_\varepsilon(x_1, y) = v_\varepsilon(x_1 - q(y), y)$$

is a family of functions in $C^\infty(\overline{\Omega})$ which, in view of the formulae for derivatives established in Lemma 2.5.10, converges to χu in $H^1(\Omega)$ as $\varepsilon \rightarrow 0$. \square

2.5.4 The trace theorem

Theorem 2.5.12 Let Ω be a bounded regular open subset of \mathbb{R}^d , with a C^∞ boundary. Denote by σ the superficial measure on $\partial\Omega$ and by N^{ext} the exterior unit normal on $\partial\Omega$. There exists a unique linear mapping

$$\gamma_0 : H^1(\Omega) \rightarrow L^2(\partial\Omega, \sigma)$$

such that, for every $\varphi \in C^\infty(\overline{\Omega})$,

$$\gamma_0 \varphi = \varphi|_{\partial\Omega}.$$

Furthermore, γ_0 satisfies the following identity,

$$(2.5.9) \quad \forall j \in \{1, \dots, d\}, \forall u, v \in H^1(\Omega), \int_{\Omega} u \partial_j v \, dx = \int_{\partial\Omega} \gamma_0 u \gamma_0 v N_j^{ext} \, d\sigma - \int_{\Omega} \partial_j u v \, dx.$$

Furthermore, $\text{Ker } \gamma_0$ is the closure $H_0^1(\Omega)$ of $C_0^\infty(\Omega)$ in $H^1(\Omega)$.

Proof.— From Corollary 2.4.16 and Remark 2.4.17, we have, for every $\varphi, \psi \in C^\infty(\overline{\Omega})$,

$$(2.5.10) \quad \forall j \in \{1, \dots, d\}, \int_{\Omega} \varphi \partial_j \psi \, dx = \int_{\partial\Omega} \varphi \psi N_j^{ext} \, d\sigma - \int_{\Omega} \partial_j \varphi \psi \, dx.$$

Consider the mapping

$$\gamma_0 : \varphi \in C^\infty(\overline{\Omega}) \mapsto \varphi|_{\partial\Omega} \in L^2(\partial\Omega, \sigma).$$

We claim that this mapping is continuous if $C^\infty(\overline{\Omega})$ is endowed with the H^1 norm. Indeed, this is equivalent to proving the estimate

$$(2.5.11) \quad \|\varphi\|_{L^2(\partial\Omega, \sigma)} \leq C \|\varphi\|_{H^1(\Omega)}.$$

Using a partition of unity associated to a finite open covering of the compact set $\partial\Omega$, it is enough to prove the above inequality when φ is supported in the intersection of Ω with a small neighbourhood of some point of $\partial\Omega$. On such a neighbourhood, we may assume that, for some j ,

$$|N_j^{ext}| \geq c$$

for some $c > 0$. Then (2.5.11) follows from (2.5.10) with $\psi = \overline{\varphi}$ and the Cauchy–Schwarz inequality.

At this stage, we appeal to Proposition 2.5.9. Since γ_0 is a continuous linear mapping from a dense subspace of $H^1(\Omega)$ into the Banach space $L^2(\partial\Omega, \sigma)$, it admits a unique linear continuous extension from $H^1(\Omega)$ to

$L^2(\partial\Omega, \sigma)$, still denoted by γ_0 . Since both sides of (2.5.9) are continuous bilinear maps on $H^1(\Omega) \times H^1(\Omega)$ and coincide on the dense subspace $\mathcal{C}^\infty(\overline{\Omega}) \times \mathcal{C}^\infty(\overline{\Omega})$ in view of (2.5.10), we infer that (2.5.9) holds.

Finally, let us characterise the kernel of γ_0 . If $\varphi \in \mathcal{C}_0^\infty(\Omega)$, we have $\gamma_0\varphi = \varphi|_{\partial\Omega} = 0$, therefore $\text{Ker } \gamma_0$ contains the closure $H_0^1(\Omega)$ of $\mathcal{C}_0^\infty(\Omega)$ in $H^1(\Omega)$. Conversely, let us assume that $u \in \text{Ker } \gamma_0$. For every $f \in L^2(\Omega)$, denote by $\underline{f} \in L^2(\mathbb{R}^d)$ the extension of f by 0 to \mathbb{R}^d . From identity (2.5.9) with $v = \varphi|_\Omega, \varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$, we infer, in $\mathcal{D}'(\mathbb{R}^d)$,

$$\partial_j(\underline{u}) = \underline{\partial_j u} + \gamma_0 u N_j^{int} \sigma = \underline{\partial_j u}.$$

Therefore $\underline{u} \in H^1(\mathbb{R}^d)$. Using the same partition of unity (χ_k) as in the proof of Proposition 2.5.9, we are reduced to prove that, for every $k \geq 1$,

$$\chi_k u \in H_0^1(\Omega).$$

Notice that $\underline{\chi_k u} = \chi_k \underline{u} \in H^1(\mathbb{R}^d)$. Let us drop again the index k and consider

$$v(z, y) = (\chi u)(z + q(y), y), \quad (z, y) \in]0, \infty[\times W.$$

Then

$$\underline{\chi u}(z + q(y), y) = \underline{v}(z, y),$$

where \underline{v} denotes the extension of v from $]0, \infty[\times W$ to $\mathbb{R} \times W$ by 0, and so we infer, by the same argument as in Lemma 2.5.10, that $\underline{v} \in H^1(\mathbb{R} \times W)$. Then we consider, for $\varepsilon > 0$,

$$v_\varepsilon(z, y) = \underline{v}(z - \varepsilon, y).$$

Of course $v_\varepsilon \in H^1(\mathbb{R} \times W)$ and $\text{supp}(v_\varepsilon) \subset [\varepsilon, a + \varepsilon] \times W$, a compact subset of $]0, \infty[\times W$. Furthermore, the continuous action of translations on L^2 implies that

$$\|v_\varepsilon - v\|_{H^1}^2 = \int_0^\infty \int_W \left(|v(z - \varepsilon, y) - v(z, y)|^2 + \sum_j |\partial_j v(z - \varepsilon, y) - \partial_j v(z, y)|^2 \right) dz dy$$

tends to 0 as $\varepsilon \rightarrow 0$. We conclude that $v_\varepsilon \rightarrow v$ in $H^1(]0, \infty[\times W)$, and, again by Lemma 2.5.10, that the family u_ε defined by

$$u_\varepsilon(x_1, y) = v_\varepsilon(x_1 - q(y), y)$$

converges to χu in $H^1(\Omega)$. Since, for every $\varepsilon > 0$, u_ε is supported into a compact subset of Ω , Proposition 2.5.8 implies that $v_\varepsilon \in H_0^1(\Omega)$. Consequently, $\chi u \in H_0^1(\Omega)$. \square

2.5.5 Solving the Dirichlet problem

Theorem 2.5.13 Let Ω be a bounded regular open subset of \mathbb{R}^d with a \mathcal{C}^∞ boundary. For every $f \in L^2(\Omega)$, there exists a unique $u \in H^1(\Omega)$ such that

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ \gamma_0 u = 0 & \text{in } \partial\Omega. \end{cases}$$

Proof.— This is an immediate consequence of Remark 2.5.5 (iii) and of Theorem 2.5.12. \square

Remark 2.5.14 One can prove in fact that, under the assumptions of Theorem 2.5.13, $u \in H^2(\Omega)$, so that the mapping

$$\Delta : H^2(\Omega) \cap H_0^1(\Omega) \rightarrow L^2(\Omega)$$

is an isomorphism if Ω is a regular bounded open subset. Furthermore, more regularity on f implies more regularity on u , typically : if $f \in H^m(\Omega)$ for some $m \in \mathbb{N}$, then $u \in H^{m+2}(\Omega)$; if $f \in C^\infty(\overline{\Omega})$, then $u \in C^\infty(\overline{\Omega})$. The latter statement is a consequence of the previous one and of

$$\bigcap_{m \in \mathbb{N}} H^m(\Omega) = C^\infty(\overline{\Omega}) .$$

2.6 The uniform boundedness principle

In this section, we prove an important result about families of distributions, which implies Lemma 2.3.24.

Theorem 2.6.1 Let (T_n) be a family of distributions on Ω such that, for every $\varphi \in \mathcal{D}(\Omega)$,

$$\sup_n |\langle T_n, \varphi \rangle| < +\infty .$$

Then, for every compact subset K of Ω , there exist $C > 0$ and $m \in \mathbb{N}$ such that, for every $\varphi \in \mathcal{D}(\Omega)$ with $\text{supp } \varphi \subset K$,

$$\sup_n |\langle T_n, \varphi \rangle| \leq C \|\varphi\|_{C^m} .$$

In other words, if for instance a sequence T_n is such that $\langle T_n, \varphi \rangle$ has a limit for every $\varphi \in \mathcal{D}(\Omega)$, one gets a uniform estimate on the action of the sequence T_n . Let us show how this implies Lemma 2.3.24. If $T_n \rightarrow T$, then the assumption of Theorem 2.6.1 is fulfilled. If $\varphi_n \rightarrow \varphi$ in $\mathcal{D}(\Omega)$, let K be a compact subset which contains the support of φ_n for every n , and hence the support of φ . By Theorem 2.6.1, we have, for some C, m independent of n ,

$$|\langle T_n, \varphi_n - \varphi \rangle| \leq C \|\varphi_n - \varphi\|_{C^m} \rightarrow 0 .$$

Then

$$\langle T_n, \varphi_n \rangle = \langle T_n, \varphi_n - \varphi \rangle + \langle T_n, \varphi \rangle \rightarrow \langle T, \varphi \rangle .$$

This completes the proof of Lemma 2.3.24.

In the rest of this section, we give a proof of Theorem 2.6.1 in three steps. For every compact subset K of Ω , we denote by \mathcal{D}_K the subspace of $\mathcal{D}(\Omega)$ of test functions φ such $\text{supp}(\varphi) \subset K$.

2.6.1 Step1. Realizing \mathcal{D}_K as a complete metric space

Lemma 2.6.2 There exists a distance function d on \mathcal{D}_K having the following properties.

i) If $\varphi_n, \varphi \in \mathcal{D}_K$ then $\varphi_n \rightarrow \varphi$ in \mathcal{D} if and only if $d(\varphi_n, \varphi) \rightarrow 0$.

- ii) The metric space (\mathcal{D}_K, d) is complete.
- iii) For every $\varphi, \psi, \theta \in \mathcal{D}_K$, $d(\varphi + \theta, \psi + \theta) = d(\varphi, \psi)$.
- iv) For every $\varepsilon > 0$, there exist $m \in \mathbb{N}$ and $r > 0$ such that the set $\{\varphi \in \mathcal{D}_K : \|\varphi\|_m \leq r\}$ is contained into the closed ball for d centered at 0 and of radius ε .

Proof.— For $\varphi, \psi \in \mathcal{D}_K$, we set

$$d(\varphi, \psi) = \sum_{m=0}^{\infty} \min(2^{-m}, \|\varphi - \psi\|_{C^m}) .$$

It is clear that d takes values in $[0, \infty[$, and that $d(\varphi, \psi) = 0 \iff \varphi = \psi$. The symmetry $d(\varphi, \psi) = d(\psi, \varphi)$ is trivial, as well item iii) of the Theorem. Finally, the triangle inequality is a consequence of the elementary inequality

$$\min(a, x + y) \leq \min(a, x) + \min(a, y) , \quad a \geq 0, x \geq 0, y \geq 0 .$$

Let us prove item i). If $\varphi_n, \varphi \in \mathcal{D}_K$, the statement $\varphi_n \rightarrow \varphi$ in \mathcal{D} is equivalent to

$$(2.6.12) \quad \forall m \in \mathbb{N}, \|\varphi_n - \varphi\|_{C^m} \rightarrow 0 .$$

Since $d(\varphi_n, \varphi) \geq \min(2^{-m}, \|\varphi_n - \varphi\|_{C^m})$, it is clear that (2.6.12) is implied by $d(\varphi_n, \varphi) \rightarrow 0$. Conversely, (2.6.12) implies that every term of the series defining $d(\varphi_n, \varphi)$ tends to 0. Since this series is normally convergent, this implies $d(\varphi_n, \varphi) \rightarrow 0$.

Let us prove item ii). Let (φ_n) be a Cauchy sequence in the metric space (\mathcal{D}_K, d) ,

$$\forall \varepsilon > 0, \exists N(\varepsilon) \in \mathbb{N} : \forall n, p \geq N(\varepsilon), d(\varphi_n, \varphi_p) \leq \varepsilon .$$

Let $m \in \mathbb{N}$ and $\varepsilon > 0$ be such that $\varepsilon < 2^{-m}$. Since $d(\varphi_n, \varphi_p) \geq \min(2^{-m}, \|\varphi_n - \varphi_p\|_{C^m})$, we conclude that, for $n, p \geq N(\varepsilon)$, $\|\varphi_n - \varphi_p\|_{C^m} \leq \varepsilon$. In other words, (φ_n) is a Cauchy sequence in the Banach space \mathcal{C}_K^m of C^m functions supported in K . Hence there exists $\varphi^{[m]} \in \mathcal{C}_K^m$ such that $\|\varphi_n - \varphi^{[m]}\|_{C^m} \rightarrow 0$. Since this is true for every $m \in \mathbb{N}$, and since convergence in \mathcal{C}_K^{m+1} implies convergence in \mathcal{C}_K^m , we conclude that

$$\varphi^{[m]} = \varphi \in \bigcap_{m \in \mathbb{N}} \mathcal{C}_K^m = \mathcal{D}_K ,$$

and that $\varphi_n \rightarrow \varphi$ in \mathcal{D} , which, by item i), means $d(\varphi_n, \varphi) \rightarrow 0$. Hence (\mathcal{D}_K, d) is a complete metric space.

Finally, let us prove item iv). Let $m \in \mathbb{N}$ be such that $2^{-m} \leq \frac{\varepsilon}{2}$, and let $r > 0$ be such that $(m+1)r \leq \frac{\varepsilon}{2}$. If $\|\varphi\|_{C^m} \leq r$, we have

$$\begin{aligned} d(\varphi, 0) &= \sum_{q=0}^{\infty} \min(2^{-q}, \|\varphi\|_{C^q}) \\ &\leq \sum_{q=0}^m \|\varphi\|_{C^q} + \sum_{q=m+1}^{\infty} 2^{-q} \\ &\leq (m+1)r + 2^{-m} \leq \varepsilon . \end{aligned}$$

This completes the proof. □

2.6.2 Step 2. The Baire lemma

Lemma 2.6.3 Let (\mathcal{E}, d) be a complete metric space. Every countable intersection of open dense subsets of \mathcal{E} is dense in \mathcal{E} . Every countable union of closed subsets of \mathcal{E} with empty interior has empty interior.

Proof.—The second statement is equivalent to the first one, since the complement of an open set is a closed set, and the complement of a dense set has an empty interior.

Let us prove the first statement. Denote by $B(\varphi, r)$ and $B_f(\varphi, r)$ respectively the open and the closed balls of radius r centered at $\varphi \in \mathcal{E}$. Let $(\mathcal{O}_k)_{k \in \mathbb{N}}$ be a sequence of dense open subsets of \mathcal{E} . Fix $\varphi_0 \in \mathcal{E}$ and $\varepsilon > 0$. We want to prove that

$$B(\varphi_0, \varepsilon) \cap \bigcap_{k \in \mathbb{N}} \mathcal{O}_k \neq \emptyset.$$

Since \mathcal{O}_0 is dense, $B(\varphi_0, \varepsilon) \cap \mathcal{O}_0$ contains some element φ_1 . Since $B(\varphi_0, \varepsilon) \cap \mathcal{O}_0$ is open, there exists $r_1 > 0$ such that

$$B_f(\varphi_1, r_1) \subset B(\varphi_0, \varepsilon) \cap \mathcal{O}_0,$$

moreover we may assume that $r_1 \leq \frac{\varepsilon}{2}$.

Since \mathcal{O}_1 is dense, $B(\varphi_1, r_1) \cap \mathcal{O}_1$ contains some element φ_2 . Since $B(\varphi_1, r_1) \cap \mathcal{O}_1$ is open, there exists $r_2 > 0$ such that

$$B_f(\varphi_2, r_2) \subset B(\varphi_1, r_1) \cap \mathcal{O}_1,$$

moreover we may assume that $r_2 \leq \frac{\varepsilon}{2^2}$.

Continuing that way, we define a sequence $(\varphi_k)_{k \geq 1}$ of elements of \mathcal{E} and a sequence $(r_k)_{k \geq 1}$ of positive numbers such that

$$\forall k \geq 1 \quad B_f(\varphi_{k+1}, r_{k+1}) \subset B(\varphi_k, r_k) \cap \mathcal{O}_k, \quad r_k \leq \frac{\varepsilon}{2^k}.$$

In particular, the sequence of closed balls $(B_f(\varphi_k, r_k))_{k \geq 1}$ is decreasing, with a radius tending to 0. Moreover,

$$B_f(\varphi_k, r_k) \subset \mathcal{O}_k \cap \cdots \cap \mathcal{O}_0 \cap B(\varphi_0, \varepsilon).$$

Hence we are reduced to prove that

$$\bigcap_{k=1}^{\infty} B_f(\varphi_k, r_k) \neq \emptyset.$$

We observe that $(\varphi_k)_{k \geq 1}$ is a Cauchy sequence in \mathcal{E} . Indeed, if $\ell \geq 0$, $\varphi_{k+\ell} \in B_f(\varphi_k, r_k)$, hence

$$d(\varphi_{k+\ell}, \varphi_k) \leq r_k,$$

which tends to 0 as k tends to ∞ . Since \mathcal{E} is a complete metric space, φ_k has a limit $\varphi \in \mathcal{E}$. Passing to the limit as ℓ tends to infinity in the above inequality, we get, for every $k \geq 1$,

$$d(\varphi, \varphi_k) \leq r_k,$$

so that

$$\varphi \in \bigcap_{k=1}^{\infty} B_f(\varphi_k, r_k).$$

This completes the proof. □

2.6.3 Step 3. Proof of Theorem 2.6.1

Let us use the notation of Theorem 2.6.1. We set, for every $k \in \mathbb{N}$,

$$\mathcal{F}_k = \{\varphi \in \mathcal{D}_K : \forall n, |\langle T_n, \varphi \rangle| \leq k\}.$$

Since each T_n is continuous on \mathcal{D}_K , \mathcal{F}_k is a closed subset of \mathcal{D}_K . Furthermore, the assumption of Theorem 2.6.1 precisely means that

$$\bigcup_{k \in \mathbb{N}} \mathcal{F}_k = \mathcal{D}_K.$$

Let endow \mathcal{D}_K with a distance function as in Lemma 2.6.2. By Baire's lemma, we infer that there exists k_0 such that \mathcal{F}_{k_0} has a nonempty interior, which means that there exists some $\varphi_0 \in \mathcal{D}_K$ and $\varepsilon > 0$ such that

$$B_f(\varphi_0, \varepsilon) \subset \mathcal{F}_{k_0}.$$

By item iii) of Lemma 2.6.2, we know that

$$B_f(0, \varepsilon) = -\varphi_0 + B_f(\varphi_0, \varepsilon),$$

so that, for every $\psi \in B_f(0, \varepsilon)$, for every n ,

$$|\langle T_n, \psi \rangle| \leq |\langle T_n, -\varphi_0 \rangle| + k_0 \leq A,$$

for some $A > 0$. Using item iv) of Lemma 2.6.2, we infer that there exists $m \in \mathbb{N}$ and $r > 0$ such that every $\psi \in \mathcal{D}_K$ such that $\|\psi\|_{C^m} \leq r$ belongs to $B_f(0, \varepsilon)$, hence satisfies

$$\sup_n |\langle T_n, \psi \rangle| \leq A.$$

Given $\varphi \in \mathcal{D}_K \setminus \{0\}$, we may apply this fact to

$$\psi = r \frac{\varphi}{\|\varphi\|_{C^m}},$$

and we conclude that

$$\sup_n |\langle T_n, \varphi \rangle| \leq \frac{A}{r} \|\varphi\|_{C^m}.$$

The proof is complete.

Remark 2.6.4 The above theorem is an adaptation of the Banach–Steinhaus theorem to the case of vector spaces admitting a distance enjoying Lemma 2.6.2. Such spaces are called Fréchet spaces.

Chapter 3

The Fourier Transformation

The Fourier transform of a function $f \in L^1(\mathbb{R}^d)$ is the function $\mathcal{F}(f) \in L^\infty(\mathbb{R}^d)$ given by

$$\mathcal{F}(f)(\xi) = \int_{\mathbb{R}^d} e^{-ix \cdot \xi} f(x) dx, \text{ with } \|\mathcal{F}(f)\|_{L^\infty} \leq \|f\|_{L^1}.$$

The important role played by the Fourier transform in PDE's theory is mainly due to the fact that, when these objects are well-defined,

$$\mathcal{F}(\partial_j f)(\xi) = i\xi_j \mathcal{F}(f)(\xi).$$

Otherwise stated, \mathcal{F} transforms the action of a differential operator with constant coefficients to that of the product by a polynomial. This would be worthless without an inversion formula giving back the function f in terms of $\mathcal{F}(f)$, as

$$f(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \mathcal{F}(f)(\xi) d\xi,$$

Unfortunately, this formula only makes sense when $\mathcal{F}(f) \in L^1(\mathbb{R}^d)$, and this is not the case in general for $f \in L^1$. In this chapter, we are going to introduce a subspace of distributions on \mathbb{R}^d , which contains $L^1(\mathbb{R}^d)$, and a Fourier transformation on this space which satisfies the two above identity.

3.1 The Schwartz space

3.1.1 Definitions and examples

Definition 3.1.1 A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is said to be rapidly decreasing if, as $|x| \rightarrow \infty$,

$$\forall \alpha \in \mathbb{N}^d, x^\alpha f(x) \rightarrow 0.$$

Notice that, if f is rapidly decreasing and continuous, all the functions $x^\alpha f$ are bounded on \mathbb{R}^d .

Definition 3.1.2 We denote $\mathcal{S}(\mathbb{R}^d)$ the set of functions $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$ which are rapidly decreasing as well as all their derivatives. In other words,

$$\forall (\alpha, \beta) \in \mathbb{N}^d, x^\alpha \partial^\beta \varphi(x) \rightarrow 0$$

as $|x| \rightarrow \infty$. The set $\mathcal{S}(\mathbb{R}^d)$ is a vector space; it is called the Schwartz space.

Example 3.1.3 i) $\mathcal{C}_0^\infty(\mathbb{R}^d) \subset \mathcal{S}(\mathbb{R}^d)$.

ii) For $z \in \mathbb{C}$ such that $\operatorname{Re} z > 0$, the function $\varphi(x) = e^{-z|x|^2}$ belongs to $\mathcal{S}(\mathbb{R}^d)$.

iii) If $\varphi_1, \varphi_2 \in \mathcal{S}(\mathbb{R}^d)$, then $\varphi_1 \varphi_2 \in \mathcal{S}(\mathbb{R}^d)$.

iv) No rational function (even smooth ones) belongs to $\mathcal{S}(\mathbb{R}^d)$.

The topology on $\mathcal{S}(\mathbb{R}^d)$ we will work with is that given by the family of norms $(N_p)_{p \in \mathbb{N}}$ given by

$$N_p(\varphi) = \sup_{|\alpha|, |\beta| \leq p} |x^\alpha \partial^\beta \varphi(x)|.$$

It is clear that for $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$, we have the equivalence

$$\varphi \in \mathcal{S}(\mathbb{R}^d) \iff \forall p \in \mathbb{N}, N_p(\varphi) < +\infty.$$

Proposition 3.1.4 If $\varphi \in \mathcal{S}(\mathbb{R}^d)$ then $x^\alpha \partial^\beta \varphi \in \mathcal{S}(\mathbb{R}^d)$ for every $\alpha, \beta \in \mathbb{N}^d$.

Proof.— This follows immediately from the fact that

$$(3.1.1) \quad N_p(x^\alpha \partial^\beta \varphi) = \sup_{|\lambda|, |\mu| \leq p} |x^\lambda \partial^\mu (x^\alpha \partial^\beta \varphi(x))| \leq C_{p, \alpha, \beta} N_{p+q}(\varphi)$$

when $|\alpha|, |\beta| \leq q$. □

3.1.2 Convergence in $\mathcal{S}(\mathbb{R}^d)$ and density results

Definition 3.1.5 Let (φ_n) be a sequence of functions in $\mathcal{S}(\mathbb{R}^d)$. One says that (φ_n) converges to φ in $\mathcal{S}(\mathbb{R}^d)$ when, for every $\alpha, \beta \in \mathbb{N}^d$, $x^\alpha \partial^\beta \varphi_n(x) \rightarrow x^\alpha \partial^\beta \varphi(x)$ uniformly in $x \in \mathbb{R}^d$. Equivalently, for all $p \in \mathbb{N}$,

$$N_p(\varphi_n - \varphi) \rightarrow 0 \text{ as } j \rightarrow +\infty.$$

Remark 3.1.6 Let $\alpha \in \mathbb{N}^d$. It follows from (3.1.1) that, if (φ_n) converges to φ in $\mathcal{S}(\mathbb{R}^d)$, then $(x^\alpha \varphi_n)$ converges to $x^\alpha \varphi$ and $(\partial^\alpha \varphi_n)$ converges to $(\partial^\alpha \varphi)$ in $\mathcal{S}(\mathbb{R}^d)$. Otherwise stated, multiplication by a polynomial and derivation are continuous operations in $\mathcal{S}(\mathbb{R}^d)$.

We already know that $\mathcal{S} \subset L^\infty$, with $\|\varphi\|_{L^\infty} = N_0(\varphi)$. The next proposition states a similar property for L^1 .

Proposition 3.1.7 $\mathcal{S}(\mathbb{R}^d) \subset L^1(\mathbb{R}^d)$, and there exists $C_d > 0$ such that

$$\forall \varphi \in \mathcal{S}(\mathbb{R}^d), \|\varphi\|_{L^1} \leq C_d N_{d+1}(\varphi).$$

Proof.— We have

$$(1 + |x|^2)^{d+1} = (1 + x_1^2 + \dots + x_d^2)^{d+1} = \sum_{|\alpha| \leq d+1} c_{\alpha,d} (x^\alpha)^2,$$

hence

$$(1 + |x|^2)^{d+1} |\varphi(x)|^2 \leq B_d N_{d+1}(\varphi)^2.$$

Consequently,

$$\int_{\mathbb{R}^d} |\varphi(x)| dx \leq \sqrt{B_d} N_{d+1}(\varphi) \int_{\mathbb{R}^d} \frac{dx}{(1 + |x|^2)^{(d+1)/2}} \leq C_d N_{d+1}(\varphi),$$

since $x \mapsto (1 + |x|^2)^{-(d+1)/2}$ is integrable on \mathbb{R}^d . □

Remark 3.1.8 For every $q \in \mathbb{N}$, one can establish similarly

$$\forall \varphi \in \mathcal{S}(\mathbb{R}^d), \forall x \in \mathbb{R}^d, (1 + |x|^q) |\varphi(x)| \leq C_{q,d} N_q(\varphi).$$

Since $\mathcal{C}_0^\infty(\mathbb{R}^d)$ is dense in the $L^p(\mathbb{R}^d)$ spaces for $p \in [1, +\infty[$, we get the

Corollary 3.1.9 The set $\mathcal{S}(\mathbb{R}^d)$ is dense in all the $L^p(\mathbb{R}^d)$ for $p \in [1, +\infty[$.

We also have the important

Proposition 3.1.10 The space $\mathcal{C}_0^\infty(\mathbb{R}^d)$ is dense in $\mathcal{S}(\mathbb{R}^d)$: for any $\varphi \in \mathcal{S}(\mathbb{R}^d)$, there is a sequence (φ_n) of functions in $\mathcal{C}_0^\infty(\mathbb{R}^d)$ which converges to φ in $\mathcal{S}(\mathbb{R}^d)$.

Proof.— Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$, and let $\chi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$ be a plateau function over $B(0, 1)$. We set $\varphi_n(x) = \varphi(x)\chi(x/n)$. The functions φ_n are smooth with compact support, and equal φ in $B(0, n)$. By the Leibniz formula, we obtain

$$\partial^\beta(\varphi - \varphi_n)(x) = \partial^\beta \varphi(x)(1 - \chi(x/n)) + \sum_{|\gamma| \geq 1, \gamma \leq \beta} C_{\beta}^{\gamma} \frac{1}{n^{|\gamma|}} \partial^{\beta-\gamma} \varphi(x) (\partial^\gamma \chi)\left(\frac{x}{n}\right).$$

Thus, as $n \rightarrow +\infty$,

$$\|x^\alpha \partial^\beta(\varphi - \varphi_n)(x)\|_\infty \leq \sup_{|x| \geq n} |x^\alpha \partial^\beta \varphi(x)| + \frac{C}{n} \sum_{\gamma \leq \beta} \|x^\alpha \partial^{\beta-\gamma} \varphi\|_\infty \rightarrow 0.$$

□

3.2 The Fourier transformation in $\mathcal{S}(\mathbb{R}^d)$

3.2.1 Definition and first properties

For $\varphi \in \mathcal{S}(\mathbb{R}^d)$ we have $\varphi \in L^1(\mathbb{R}^d)$, so that $\mathcal{F}(\varphi)$ is well defined and belongs to $L^\infty(\mathbb{R}^d)$.

Definition 3.2.1 For $\varphi \in \mathcal{S}(\mathbb{R}^d)$, we denote $\hat{\varphi}$, $\mathcal{F}(\varphi)$ or even $\mathcal{F}_{x \rightarrow \xi}(\varphi(x))$ the function in $L^\infty(\mathbb{R}^d)$ given by

$$\hat{\varphi}(\xi) = \mathcal{F}(\varphi)(\xi) = \mathcal{F}_{x \rightarrow \xi}(\varphi(x)) = \int_{\mathbb{R}^d} e^{-ix \cdot \xi} \varphi(x) dx.$$

The linear map $\mathcal{F} : \varphi \mapsto \hat{\varphi}$ is called the Fourier transformation.

Here follows some of the properties of the Fourier transform on $\mathcal{S}(\mathbb{R}^d)$ that we have asked for in the introduction.

Proposition 3.2.2 Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$. Then $\hat{\varphi} \in \mathcal{S}(\mathbb{R}^d)$. More precisely,

- i) For all $j \in \{1, \dots, n\}$ $\mathcal{F}_{x \rightarrow \xi}(x_j \varphi(x)) = i \partial_j \mathcal{F}(\varphi)(\xi)$.
- ii) For all $j \in \{1, \dots, n\}$, we have $\mathcal{F}(\partial_j \varphi)(\xi) = i \xi_j \mathcal{F}(\varphi)(\xi)$.
- iii) For $a \in \mathbb{R}^d$, $\mathcal{F}_{x \rightarrow \xi}(\varphi(x - a)) = e^{-ia \cdot \xi} \mathcal{F}(\varphi)(\xi)$.
- iv) For $a \in \mathbb{R}^d$, $\mathcal{F}_{x \rightarrow \xi}(e^{ia \cdot x} \varphi(x)) = \mathcal{F}(\varphi)(\xi - a)$.
- v) For all integer $p \geq 0$, $N_p(\hat{\varphi}) \leq C_{p,d} N_{p+d+1}(\varphi)$.

Proof.— i) The function $(x, \xi) \mapsto e^{-ix \cdot \xi} \varphi(x)$ is \mathcal{C}^1 on \mathbb{R}^d , et

$$|\partial_{\xi_j}(e^{-ix \cdot \xi} \varphi(x))| = |-ix_j e^{-ix \cdot \xi} \varphi(x)| = |x_j \varphi(x)| \in L^1(\mathbb{R}^d).$$

By Lebesgue theorem, we see that $\mathcal{F}(\varphi)$ is \mathcal{C}^1 and

$$\partial_{\xi_j} \mathcal{F}(\varphi)(\xi) = \int_{\mathbb{R}^d} -ix_j e^{-ix \cdot \xi} \varphi(x) dx = \mathcal{F}(-ix_j \varphi(x)).$$

ii) Let us first write the proof for $j = 1$. Integrating by parts, we get

$$\int_{\mathbb{R}} \partial_1 \varphi(x) e^{-ix \cdot \xi} dx_1 = i \xi_1 \int_{\mathbb{R}} \varphi(x) e^{-ix \cdot \xi} dx_1$$

Now we integrate with respect to the variable x' . By Fubini, since $\varphi, \partial_1 \varphi \in \mathcal{S}(\mathbb{R}^d) \subset L^1(\mathbb{R}^d)$, we have

$$\int_{\mathbb{R}^d} \partial_1 \varphi(x) e^{-ix \cdot \xi} dx = i \xi_1 \int_{\mathbb{R}^d} \varphi(x) e^{-ix \cdot \xi} dx.$$

To get (iv), we only have to write

$$\mathcal{F}_{x \rightarrow \xi}(e^{ia \cdot x} \varphi(x)) = \int_{\mathbb{R}^d} e^{-ix \cdot \xi} e^{ia \cdot x} \varphi(x) dx = \int_{\mathbb{R}^d} e^{-i(\xi-a) \cdot x} \varphi(x) dx = \mathcal{F}(\varphi)(\xi - a).$$

Eventually, performing a change of variable, we have

$$\mathcal{F}_{x \rightarrow \xi}(\varphi(x - a)) = \int_{\mathbb{R}^d} e^{-ix \cdot \xi} \varphi(x - a) dx = \int_{\mathbb{R}^d} e^{-i(x+a) \cdot \xi} \varphi(x) dx = e^{-ia \cdot \xi} \mathcal{F}(\varphi)(\xi),$$

and this is property (iii).

Iterating properties $i)$, $ii)$, we obtain that $\hat{\varphi} \in \mathcal{S}(\mathbb{R}^d)$, and, using Proposition 3.1.7, that

$$\begin{aligned} N_p(\hat{\varphi}) &= \sup_{|\alpha|, |\beta| \leq p} \|\xi^\alpha \partial^\beta \hat{\varphi}\|_{L^\infty} = \sup_{|\alpha|, |\beta| \leq p} \|\mathcal{F}(\partial^\alpha(x^\beta \varphi))\|_{L^\infty} \\ &\leq \sup_{|\alpha|, |\beta| \leq p} \|\partial^\alpha(x^\beta \varphi)\|_{L^1} \leq B_d \sup_{|\alpha|, |\beta| \leq p} N_{d+1}(\partial^\alpha(x^\beta \varphi)) \\ &\leq C_{p,d} N_{p+d+1}(\varphi). \end{aligned}$$

□

Because of the presence of a factor $i = \sqrt{-1}$ in (i) and (ii), it is sometimes convenient to use the notation

$$D_j = \frac{1}{i} \partial_j.$$

Then, for example, (ii) becomes $\mathcal{F}(D_j \varphi) = \xi_j \mathcal{F}(\varphi)$, and (i) is $D_j \mathcal{F}(\varphi) = -\mathcal{F}(x_j \varphi)$. Summing up, we have

$$\begin{cases} \widehat{D_j \varphi} = \xi_j \hat{\varphi}, \\ \widehat{x_j \varphi} = -D_j \hat{\varphi}. \end{cases}$$

Notice also that, by (i), $\mathcal{F}(\varphi)$ is \mathcal{C}^∞ since $x^\alpha \varphi \in \mathcal{S}$ for all $\alpha \in \mathbb{N}^d$.

3.2.2 Gaussians

For $\lambda > 0$, we set

$$G_\lambda(x) = e^{-\lambda \frac{|x|^2}{2}}, \quad x \in \mathbb{R}^d.$$

Proposition 3.2.3

$$\hat{G}_\lambda = \left(\frac{2\pi}{\lambda} \right)^{\frac{d}{2}} G_{\frac{1}{\lambda}}.$$

Proof.— Since $G_\lambda(x_1, \dots, x_d) = G_\lambda(x_1) \dots G_\lambda(x_d)$, it is enough to prove the case $d = 1$, which we now assume. Then we observe that

$$G'_\lambda(x) = -\lambda x G_\lambda(x),$$

hence by properties $i)$ and $ii)$ of Proposition 3.2.2, we infer

$$i\xi \hat{G}_\lambda(\xi) = -i\lambda \frac{d}{d\xi} \hat{G}_\lambda(\xi).$$

Solving this differential equation leads to $\hat{G}_\lambda(\xi) = \hat{G}_\lambda(0) e^{-\frac{\xi^2}{2\lambda}}$, and the proof is completed by the Gauss integral identity

$$\hat{G}_\lambda(0) = \int_{\mathbb{R}} e^{-\lambda \frac{x^2}{2}} dx = \sqrt{\frac{2\pi}{\lambda}}.$$

□

Exercise 3.2.4 Compute $\mathcal{F}_{x \rightarrow \xi}(e^{-\lambda x^2})$ by calculating $\int_{\text{Im } \zeta = \xi} e^{-\lambda \zeta^2} d\zeta$ using the Cauchy integral formula for holomorphic functions.

3.2.3 The Inversion formula

Theorem 3.2.5 Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$. Then

$$\varphi(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \hat{\varphi}(\xi) d\xi.$$

Proof.— We first observe that it is enough to prove this identity for $x = 0$. Indeed, in view of property (iii) of Proposition 3.2.2, the general statement follows from applying the formula

$$\psi(0) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \hat{\psi}(\xi) d\xi$$

to the function ψ_x defined by $\psi_x(y) = \varphi(x + y)$.

Consider the linear form

$$L : \psi \in \mathcal{S}(\mathbb{R}^d) \mapsto \int_{\mathbb{R}^d} \hat{\psi}(\xi) d\xi$$

We claim that, if $\psi(0) = 0$, then $L(\psi) = 0$. Indeed, applying the Hadamard lemma 2.3.12 and its proof, we infer

$$\psi(x) = \sum_{j=1}^d x_j \psi_j(x),$$

where $\psi_1, \dots, \psi_d \in \mathcal{S}(\mathbb{R}^d)$. Taking the Fourier transform of both sides and applying property (i) of Proposition 3.2.2, we infer

$$\hat{\psi}(\xi) = \sum_{j=1}^d i \partial_j \hat{\psi}_j(\xi).$$

Integrating both sides on \mathbb{R}^d , we conclude

$$\int_{\mathbb{R}^d} \hat{\psi}(\xi) d\xi = 0.$$

Hence we have proved that the kernel of the linear form $\psi \mapsto \psi(0)$ on $\mathcal{S}(\mathbb{R}^d)$ is contained in the kernel of L . Consequently, there exists $c \in \mathbb{C}$ such that, for every $\psi \in \mathcal{S}(\mathbb{R}^d)$,

$$L(\psi) = c\psi(0).$$

Applying this identity to $\psi = G_1$, we infer, in view of Proposition 3.2.3,

$$c = (2\pi)^d .$$

□

In order to reformulate this important theorem, we introduce the symmetry $\sigma : \mathcal{S} \rightarrow \mathcal{S}$ defined by

$$\sigma\varphi(x) = \check{\varphi}(x) := \varphi(-x) ,$$

and observe that

$$\sigma \circ \mathcal{F} = \mathcal{F} \circ \sigma =: \check{\mathcal{F}} .$$

Corollary 3.2.6 The Fourier transformation is an isomorphism on the vector space $\mathcal{S}(\mathbb{R}^d)$. Its inverse \mathcal{F}^{-1} is given by

$$\mathcal{F}^{-1} = (2\pi)^{-d} \check{\mathcal{F}} .$$

Equivalently, we have

$$(3.2.2) \quad \mathcal{F} \circ \mathcal{F} = (2\pi)^d \sigma .$$

Remark 3.2.7 As an immediate consequence of (3.2.2), $\mathcal{F}^4 = (\mathcal{F} \circ \mathcal{F})^2 = (2\pi)^{2d} I$.

3.2.4 The “change of head” lemma

The following proposition is elementary but crucial for the whole chapter.

Proposition 3.2.8 Let $\varphi, \psi \in \mathcal{S}(\mathbb{R}^d)$. Then

$$\int_{\mathbb{R}^d} \hat{\varphi}(\xi) \psi(\xi) d\xi = \int_{\mathbb{R}^d} \varphi(x) \hat{\psi}(x) dx .$$

Proof.— By Fubini’s theorem,

$$\begin{aligned} \int_{\mathbb{R}^d} \hat{\varphi}(\xi) \psi(\xi) d\xi &= \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} e^{-ix \cdot \xi} \varphi(x) dx \right) \psi(\xi) d\xi \\ &= \int_{\mathbb{R}^d} \varphi(x) \left(\int_{\mathbb{R}^d} e^{-ix \cdot \xi} \psi(\xi) d\xi \right) dx = \int_{\mathbb{R}^d} \varphi(x) \hat{\psi}(x) dx . \end{aligned}$$

□

We now come to some important identities which easily follow from the inversion formula.

3.2.5 The Plancherel identity

Proposition 3.2.9 (Plancherel formula) Let φ and ψ be two functions in $\mathcal{S}(\mathbb{R}^d)$. Then

$$\int_{\mathbb{R}^d} \varphi(x) \overline{\psi(x)} dx = (2\pi)^{-d} \int_{\mathbb{R}^d} \hat{\varphi}(\xi) \overline{\hat{\psi}(\xi)} d\xi$$

Proof.— We first observe the following identity,

$$(3.2.3) \quad \overline{\hat{\varphi}} = \sigma(\hat{\overline{\varphi}}),$$

which is merely a reformulation of

$$\overline{\hat{\varphi}}(\xi) = \int_{\mathbb{R}^d} e^{ix \cdot \xi} \overline{\varphi(x)} dx.$$

Then, applying Proposition 3.2.8, we obtain

$$\int_{\mathbb{R}^d} \hat{\varphi} \overline{\hat{\psi}} = \int_{\mathbb{R}^d} \varphi \overline{\hat{\hat{\psi}}}.$$

But, combining identity (3.2.3) with identity (3.2.2),

$$\hat{\hat{\psi}} = \sigma \overline{\hat{\psi}} = (2\pi)^d \sigma \overline{\overline{\hat{\psi}}} = (2\pi)^d \overline{\hat{\psi}}.$$

This completes the proof. □

In terms of the Hermitian scalar product in $L^2(\mathbb{R}^d, \mathbb{C})$, the above identity can be written as

$$(\varphi, \psi)_{L^2} = \frac{1}{(2\pi)^d} (\hat{\varphi}, \hat{\psi})_{L^2}.$$

In particular, for $\psi = \varphi$, we obtain the famous Plancherel formula in $\mathcal{S}(\mathbb{R}^d)$:

Corollary 3.2.10 (Plancherel) For any $\varphi \in \mathcal{S}(\mathbb{R}^d)$, it holds that

$$\|\varphi\|_{L^2(\mathbb{R}^d)} = (2\pi)^{-d/2} \|\hat{\varphi}\|_{L^2(\mathbb{R}^d)}.$$

3.3 The space $\mathcal{S}'(\mathbb{R}^d)$ of tempered distributions

3.3.1 Definition, examples

Definition 3.3.1 A distribution $T \in \mathcal{D}'(\mathbb{R}^d)$ is said to be (a) tempered (distribution) when there exists $C > 0$ and $p \in \mathbb{N}$ such that, for all test function $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$,

$$|\langle T, \varphi \rangle| \leq CN_p(\varphi).$$

Notation. We denote by $\mathcal{S}'(\mathbb{R}^d)$ the space of tempered distributions on \mathbb{R}^d .

Example 3.3.2 If $T \in \mathcal{E}'(\mathbb{R}^d)$, there is $C > 0$, $m \in \mathbb{N}$, such that for any $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$,

$$|\langle T, \varphi \rangle| \leq C \sum_{|\alpha| \leq m} \sup |\partial^\alpha \varphi| \leq CN_p(\varphi).$$

Thus $\mathcal{E}'(\mathbb{R}^d) \subset \mathcal{S}'(\mathbb{R}^d)$, that is compactly supported distributions are tempered.

Example 3.3.3 For $p \in [1, +\infty]$, we have $L^p(\mathbb{R}^d) \subset \mathcal{S}'(\mathbb{R}^d)$. Indeed, if $f \in L^p(\mathbb{R}^d)$ and $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$, and for $q \in [1, +\infty]$ such that $1/p + 1/q = 1$, we have

$$|\langle T_f, \varphi \rangle| \leq \left| \int f(x)\varphi(x)dx \right| \leq \|f\|_{L^p} \|\varphi\|_{L^q} \leq C \|f\|_{L^p} N_{n+1}(\varphi),$$

thanks to Proposition 3.2.2. In particular, in view of Proposition 3.1.7, $\mathcal{S}(\mathbb{R}^d) \subset \mathcal{S}'(\mathbb{R}^d)$.

Example 3.3.4 Let $f \in L^1_{loc}(\mathbb{R}^d)$ be such that, for some $C > 0$ and $p \in \mathbb{N}$,

$$\forall R > 0, \int_{|x| \leq R} |f(x)| dx \leq C(1 + R)^p.$$

. Then we claim that f is a tempered distribution. Indeed, if $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$, we have

$$\begin{aligned} \left| \int_{\mathbb{R}^d} f\varphi \right| &\leq \int_{\mathbb{R}^d} |f(x)||\varphi(x)| dx = \int_{|x| < 1} |f(x)||\varphi(x)| dx + \sum_{k=0}^{\infty} \int_{2^k \leq |x| < 2^{k+1}} |f(x)||\varphi(x)| dx \\ &\leq 2^p CN_0(\varphi) + C \sum_{k=0}^{\infty} (1 + 2^{k+1})^p \sup_{2^k \leq |x| < 2^{k+1}} |\varphi(x)| \\ &\leq C' \left(N_0(\varphi) + \sum_{k=0}^{\infty} 2^{-k} \sup_{2^k \leq |x| < 2^{k+1}} |x|^{p+1} |\varphi(x)| \right) \\ &\leq C'' N_{p+1}(\varphi). \end{aligned}$$

Example 3.3.5 The previous example admits the following partial converse. If $f \in L^1_{loc}(\mathbb{R}^d)$ has nonnegative values and is a tempered distribution, then, for some $C > 0$ and $p \in \mathbb{N}$,

$$\forall R > 0, \int_{|x| \leq R} f(x) dx \leq C(1 + R)^p.$$

Indeed, if f is tempered, we have, for some $A > 0$ et $p \in \mathbb{N}$,

$$\forall \varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d), \left| \int_{\mathbb{R}^d} f(x)\varphi(x) dx \right| \leq AN_p(\varphi).$$

Let $\chi \geq 0$ be a plateau function over the unit ball centered at 0, supported into the ball of radius 2 centered at 0. Then

$$0 \leq \int_{|x| \leq R} f(x) dx \leq \int_{\mathbb{R}^d} f(x)\chi\left(\frac{x}{R}\right) dx \leq AN_p\left(\chi\left(\frac{\cdot}{R}\right)\right) \leq C(1 + R)^p.$$

In particular, e^x is not tempered on \mathbb{R} , since, for every $p \in \mathbb{N}$,

$$R^{-p} \int_0^R e^x dx \xrightarrow{R \rightarrow \infty} +\infty.$$

We conclude this first paragraph with a simple observation.

Proposition 3.3.6 If $T \in \mathcal{S}'(\mathbb{R}^d)$, then $x^\alpha \partial^\beta T \in \mathcal{S}'(\mathbb{R}^d)$ for all $\alpha, \beta \in \mathbb{N}^d$.

Proof.— It is sufficient to show that $x_j T$ and $\partial_j T$ are tempered distributions when T is. Since $\varphi \in \mathcal{S}(\mathbb{R}^d)$, $\langle x_j T, \varphi \rangle = \langle T, x_j \varphi \rangle$ and $\langle \partial_j T, \varphi \rangle = -\langle T, \partial_j \varphi \rangle$, the proposition follows directly from (3.1.1). \square

Exercise 3.3.7 Show that the function $x \mapsto e^x e^{ie^x}$ is not bounded by a polynomial, but belongs to $\mathcal{S}'(\mathbb{R})$. Hint: it is the derivative of a tempered distribution.

The multiplication of a tempered distribution by a smooth function does not always yield a tempered distribution. However, it is the case when the function has moderate growth, in the following sense.

Definition 3.3.8 A function $f \in \mathcal{C}^\infty(\mathbb{R}^d)$ has moderate growth when for any $\beta \in \mathbb{N}^d$, there is $C_\beta > 0$ and $m_\beta \in \mathbb{N}$ such that

$$|\partial^\beta f(x)| \leq C_\beta (1 + |x|)^{m_\beta}.$$

We denote $\mathcal{O}_M(\mathbb{R}^d)$ the set of such functions.

Proposition 3.3.9 If $T \in \mathcal{S}'(\mathbb{R}^d)$ and $f \in \mathcal{O}_M(\mathbb{R}^d)$, then $fT \in \mathcal{S}'(\mathbb{R}^d)$.

Proof.— Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$, and $\alpha, \beta \in \mathbb{N}^d$. The Leibniz formula gives

$$|x^\alpha \partial^\beta (f\varphi)| \leq \sum_{\gamma \leq \alpha} \binom{\beta}{\gamma} |x^\alpha \partial^\gamma f| |\partial^{\beta-\gamma} \varphi| \leq C_\gamma \sum_{\gamma \leq \alpha} \binom{\beta}{\gamma} (1 + |x|)^{m_\gamma} |\partial^{\beta-\gamma} \varphi|.$$

Then, as in (3.1.1), we obtain

$$N_p(f\varphi) \leq CN_{p+M}(\varphi),$$

where $M = \max_{|\gamma| \leq p} m_\gamma$. Thus, for $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$, we obtain

$$|\langle fT, \varphi \rangle| = |\langle T, f\varphi \rangle| \leq CN_p(f\varphi) \leq C'N_{p+M}(\varphi),$$

and this means that fT is a tempered distribution. \square

Exercise 3.3.10 Show that $\text{vp}(1/x) \in \mathcal{S}'(\mathbb{R})$.

3.3.2 The fundamental characterization of \mathcal{S}'

Proposition 3.3.11 If T is a tempered distribution, then T extends in a unique way as a linear form \tilde{T} on $\mathcal{S}(\mathbb{R}^d)$ continuous in the following sense: if $\varphi_n \rightarrow \varphi$ in $\mathcal{S}(\mathbb{R}^d)$, then $\langle \tilde{T}, \varphi_n \rangle \rightarrow \langle \tilde{T}, \varphi \rangle$.

Proof.— Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$. There is a sequence (φ_j) in $\mathcal{C}_0^\infty(\mathbb{R}^d)$ such that $\varphi_j \rightarrow \varphi$ in $\mathcal{S}(\mathbb{R}^d)$. The sequence $(\langle T, \varphi_j \rangle)_j$ is a Cauchy sequence in \mathbb{C} since T is tempered:

$$(3.3.4) \quad |\langle T, \varphi_j - \varphi_k \rangle| \leq CN_p(\varphi_j - \varphi_k).$$

Its limit does not depend on the choice of the sequence (φ_j) , since when $\varphi_j, \psi_j \rightarrow \varphi$, we have

$$|\langle T, \varphi_j - \psi_j \rangle| \leq CN_p(\varphi_j - \psi_j) \rightarrow 0.$$

Thus we can let \tilde{T} be the linear form on $\mathcal{S}(\mathbb{R}^d)$ given by

$$\langle \tilde{T}, \varphi \rangle = \lim_{j \rightarrow +\infty} \langle T, \varphi_j \rangle,$$

where (φ_j) is any sequence in $\mathcal{C}_0^\infty(\mathbb{R}^d)$ which converges to φ . Since N_p is a continuous function on \mathcal{S} , passing to the limit in

$$|\langle T, \varphi_j \rangle| \leq CN_p(\varphi_j),$$

we get

$$|\langle \tilde{T}, \varphi \rangle| \leq CN_p(\varphi).$$

for any $\varphi \in \mathcal{S}(\mathbb{R}^d)$. So \tilde{T} is continuous.

Finally, if T_1 is another continuous extension of T to $\mathcal{S}(\mathbb{R}^d)$, we should have

$$\langle T_1, \varphi \rangle = \langle T_1, \varphi - \varphi_j \rangle + \langle T_1, \varphi_j \rangle \rightarrow 0 + \langle \tilde{T}, \varphi \rangle,$$

so that $T_1 = \tilde{T}$. □

Remark 3.3.12 For simplicity, we shall drop the *tilde* in the sequel, and set T in place of \tilde{T} . Notice that, conversely, every continuous linear form on $\mathcal{S}(\mathbb{R}^d)$ satisfies a bound of the form

$$\forall \varphi \in \mathcal{S}, |\langle T, \varphi \rangle| \leq CN_p(\varphi)$$

hence its restriction to $\mathcal{C}_0^\infty(\mathbb{R}^d)$ is a tempered distribution. Summing up, tempered distributions identify to continuous linear forms on the Schwartz space \mathcal{S} . This explains the notation \mathcal{S}' . We shall use this identification systematically.

3.3.3 Convergence in $\mathcal{S}'(\mathbb{R}^d)$

Definition 3.3.13 Let (T_n) be a sequence of tempered distributions. One says that (T_n) tends to T in $\mathcal{S}'(\mathbb{R}^d)$ when for any function $\varphi \in \mathcal{S}(\mathbb{R}^d)$, it holds that $\langle T_n, \varphi \rangle \rightarrow \langle T, \varphi \rangle$.

As it is the case in $\mathcal{D}'(\mathbb{R}^d)$, this notion of convergence, a weak one, implies a stronger one. We admit the following result.

Proposition 3.3.14 If $T_n \rightarrow T$ in \mathcal{S}' , there exists $C > 0$ and $p \in \mathbb{N}$ such that

$$\forall \varphi \in \mathcal{S}, \forall n, |\langle T_n, \varphi \rangle| \leq C N_p(\varphi).$$

As we did for the convergence in \mathcal{D}' , the above proposition relies on a uniform boundedness principle, which follows from identifying the convergence on \mathcal{S} as the one on a complete metric space.

Remark 3.3.15 When $T_n \rightarrow T$ in $\mathcal{S}'(\mathbb{R})$, it is true that $T_n \rightarrow T$ in $\mathcal{D}'(\mathbb{R})$ since $\mathcal{C}_0^\infty(\mathbb{R}^d) \subset \mathcal{S}(\mathbb{R}^d)$. The converse is not true in general, as shown by the following example: for any sequence (a_n) of complex numbers, the sequence $(a_n \delta_n)$ tends to 0 in $\mathcal{D}'(\mathbb{R})$. However it only converges (necessarily to 0) in $\mathcal{S}'(\mathbb{R}^d)$ if (a_n) has moderate growth, i.e. there is $C > 0, p \in \mathbb{N}$ such that

$$\forall n, |a_n| \leq C(1+n)^p.$$

Remark 3.3.16 If $f_n \rightarrow f$ in $L^p(\mathbb{R}^d)$, then $f_n \rightarrow f$ in $\mathcal{S}'(\mathbb{R}^d)$. If $T_n \rightarrow T$ in $\mathcal{S}'(\mathbb{R}^d)$, then $fT_n \rightarrow fT$ in $\mathcal{S}'(\mathbb{R}^d)$ for all $f \in \mathcal{O}_M(\mathbb{R}^d)$.

3.4 The Fourier Transformation in $\mathcal{S}'(\mathbb{R}^d)$

3.4.1 Definition

Let $T \in \mathcal{S}'(\mathbb{R}^d)$ be a tempered distribution. The linear form on $\mathcal{S}(\mathbb{R}^d)$ given by $\varphi \mapsto \langle T, \hat{\varphi} \rangle$ is a tempered distribution since there exist $C > 0$ and $p \in \mathbb{N}$ such that

$$|\langle T, \hat{\varphi} \rangle| \leq C N_p(\hat{\varphi}) \leq C' N_{p+d+1}(\varphi),$$

thanks to Proposition 3.2.6.

Definition 3.4.1 For $T \in \mathcal{S}'(\mathbb{R}^d)$, we denote by $\hat{T} = \mathcal{F}(T)$ the tempered distribution given by

$$\langle \hat{T}, \varphi \rangle = \langle T, \hat{\varphi} \rangle, \varphi \in \mathcal{S}(\mathbb{R}^d).$$

Example 3.4.2 i) For $f \in L^1$, we have by Fubini's theorem,

$$\begin{aligned} \langle \hat{f}, \varphi \rangle &= \int_{\mathbb{R}^d} f(x) \hat{\varphi}(x) dx = \int_{\mathbb{R}^d} f(x) \left(\int_{\mathbb{R}^d} \varphi(\xi) e^{-ix \cdot \xi} d\xi \right) dx \\ &= \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} f(x) e^{-ix \cdot \xi} dx \right) \varphi(\xi) d\xi = \int_{\mathbb{R}^d} \hat{f}(\xi) \varphi(\xi) d\xi, \end{aligned}$$

so that $\widehat{\hat{f}} = T_{\hat{f}}$. The Fourier transformation on \mathcal{S}' restricts to the classical Fourier transformation on L^1 (see also Proposition 3.4.5 below).

ii) For $\varphi \in \mathcal{S}(\mathbb{R}^d)$, $\langle \hat{\delta}_0, \varphi \rangle = \int \varphi(x) dx$, thus $\hat{\delta}_0 = 1$.

Proposition 3.4.3 The Fourier Transform \mathcal{F} is an isomorphism on $\mathcal{S}'(\mathbb{R}^d)$. Its inverse is $\mathcal{F}^{-1} = (2\pi)^{-d}\check{\mathcal{F}}$. Moreover \mathcal{F} and \mathcal{F}^{-1} are continuous on $\mathcal{S}'(\mathbb{R}^d)$, in the following sense: if $T_n \rightarrow T \in \mathcal{S}'(\mathbb{R}^d)$, then $\mathcal{F}(T_n) \rightarrow \mathcal{F}(T)$ in $\mathcal{S}'(\mathbb{R}^d)$.

These results follow immediately from the above definition and Proposition 3.2.6. The same way, transferring to $\mathcal{S}'(\mathbb{R}^d)$ the properties of the Fourier transform on $\mathcal{S}(\mathbb{R}^d)$, we obtain easily the following identities in $\mathcal{S}'(\mathbb{R}^d)$:

$$\mathcal{F}(D_j T) = \xi_j \hat{T}, \quad \mathcal{F}(x_j T) = -D_j \hat{T}, \quad \mathcal{F}(\tau_a T) = e^{-ia \cdot \xi} \mathcal{F}(T), \quad \mathcal{F}(e^{ia \cdot x} T) = \tau_a \mathcal{F}(T).$$

Example 3.4.4 $\hat{1} = \mathcal{F} \circ \mathcal{F}(\delta_0) = (2\pi)^d \check{\delta}_0 = (2\pi)^d \delta_0$.

3.4.2 The Fourier Transformation on L^1 and L^2

Here we briefly sum up the main properties of the Fourier transform of a tempered distribution given by an L^1 or an L^2 function.

Proposition 3.4.5 If $T = f \in L^1(\mathbb{R}^d)$, then $\mathcal{F}(T) = \hat{f}$. More precisely

- i) $\mathcal{F}(T)$ is the continuous function given by $\mathcal{F}(T)(\xi) = \int e^{-ix \cdot \xi} f(x) dx$, and $\mathcal{F}(T)(\xi) \rightarrow 0$ when $|\xi| \rightarrow +\infty$.
- ii) If moreover $\mathcal{F}(T)$ belongs to $L^1(\mathbb{R}^d)$, then $\mathcal{F}^{-1}(\mathcal{F}(T)) = T$ almost everywhere.

Proof.— The fact that \hat{f} is a continuous function follows easily from Lebesgue theorem of continuity for integral with parameters, and the fact that \hat{f} goes to 0 at infinity is called the Riemann-Lebesgue lemma. For $\varphi \in \mathcal{S}(\mathbb{R}^d)$ we have

$$\langle \hat{T}, \varphi \rangle = \langle T, \hat{\varphi} \rangle = \int f(\xi) \hat{\varphi}(\xi) d\xi = \int f(\xi) \left(\int e^{-ix \cdot \xi} \varphi(x) dx \right) d\xi.$$

Since the function $(x, \xi) \mapsto f(\xi) e^{-ix \cdot \xi} \varphi(x)$ belongs to $L^1(\mathbb{R}_x^d \times \mathbb{R}_\xi^d)$, we have

$$\langle \hat{T}, \varphi \rangle = \int \varphi(x) \left(\int e^{-ix \cdot \xi} f(\xi) d\xi \right) dx = \langle \hat{f}, \varphi \rangle,$$

and this ends the proof of (i). We also know that $\mathcal{F}^{-1}(\hat{f}) = f$ in $\mathcal{S}'(\mathbb{R}^d)$. If $\hat{f} \in L^1(\mathbb{R}^d)$, we thus have $\mathcal{F}^{-1}(\hat{f}) = f$ in $\mathcal{D}'(\mathbb{R}^d)$, and this implies that $\mathcal{F}^{-1}(\hat{f}) = f$ almost everywhere. \square

Proposition 3.4.6 The map $T \in \mathcal{S}'(\mathbb{R}^d) \mapsto (2\pi)^{-d/2} \mathcal{F}(T) \in \mathcal{S}'(\mathbb{R}^d)$ induces a bijective isometry on $L^2(\mathbb{R}^d)$.

Proof.— Let $f \in L^2(\mathbb{R}^d)$. For every $\varphi \in \mathcal{S}(\mathbb{R}^d)$, we have

$$\langle \hat{f}, \varphi \rangle = \int_{\mathbb{R}^d} f(x) \hat{\varphi}(x) dx$$

and therefore, by the Cauchy–Schwarz inequality and the Plancherel identity (Corollary 3.2.10),

$$|\langle \hat{f}, \varphi \rangle| \leq \|f\|_{L^2} \|\hat{\varphi}\|_{L^2} = (2\pi)^{\frac{d}{2}} \|f\|_{L^2} \|\varphi\|_{L^2}.$$

By the Riesz representation theorem, this implies that $\hat{\in} L^2(\mathbb{R}^d)$ and

$$\|\hat{f}\|_{L^2} \leq (2\pi)^{\frac{d}{2}} \|f\|_{L^2}.$$

Applying this inequality to \hat{f} and using the inversion formula from Proposition 3.4.3, we infer

$$(2\pi)^d \|f\|_{L^2} = \|\hat{\hat{f}}\|_{L^2} \leq (2\pi)^{\frac{d}{2}} \|\hat{f}\|_{L^2}$$

so that, eventually,

$$\|f\|_{L^2} = (2\pi)^{-\frac{d}{2}} \|\hat{f}\|_{L^2}.$$

Since it is clear from Proposition 3.4.3 that \mathcal{F} is bijective on L^2 , the proof is complete. \square

Remark 3.4.7 There are functions f in $L^2(\mathbb{R}^d)$ such that $x \mapsto e^{-ix \cdot \xi} f(x)$ is not integrable whatever the value of ξ is (for example, for $d = 1$, $f(x) = (1 + |x|)^{-3/4}$). Nevertheless, for $R > 0$, the function g_R given by

$$g_R(\xi) = \int_{|x| < R} e^{-ix \cdot \xi} f(x) dx$$

tends to \hat{f} in $L^2(\mathbb{R}^d)$ thanks to the Proposition 3.4.6, since $f 1_{|x| < R} \rightarrow f$ in $L^2(\mathbb{R}^d)$. Thus, for $f \in L^2(\mathbb{R}^d)$,

$$\hat{f}(\xi) = \lim_{R \rightarrow +\infty} \int_{|x| < R} e^{-ix \cdot \xi} f(x) dx$$

in the L^2 sense.

3.4.3 The Fourier Transform of compactly supported distributions

The Fourier transformation exchanges the speed of decay at infinity of a function with the regularity of its image, as shown by example by the following inequality

$$\|D^\alpha \hat{\varphi}\|_{L^\infty} = \|\mathcal{F}(x^\alpha \varphi)\|_{L^\infty} \leq \|x^\alpha \varphi\|_{L^1}, \quad \varphi \in \mathcal{S}(\mathbb{R}^d).$$

The best speed of decay at infinity for a function is achieved for compactly supported ones ; the result of this subsection explores this phenomenon more precisely.

Proposition 3.4.8 If $T \in \mathcal{D}'(\mathbb{R}^d)$ is compactly supported, its Fourier transform $\mathcal{F}(T)$ is the smooth function on \mathbb{R}^d given by

$$\mathcal{F}(T)(\xi) = \langle T_x, e^{-ix \cdot \xi} \rangle.$$

Moreover, there is an integer $m \in \mathbb{N}$ such that, for all $\alpha \in \mathbb{N}^d$, there is a $C_\alpha > 0$ satisfying

$$|\partial^\alpha \mathcal{F}(T)(\xi)| \leq C_\alpha (1 + |\xi|)^m.$$

Proof.— Differentiating under the bracket, we see that the function $v(\xi)$ given by

$$v(\xi) = \langle T_x, e^{-ix \cdot \xi} \rangle$$

is a C^∞ function (as a matter of fact, one should write $v(\xi) = \langle T_x, \chi(x)e^{-ix \cdot \xi} \rangle$, for some plateau function $\chi \in C_0^\infty(\mathbb{R}^d)$ above the support of T , and then apply the theorem). We also have

$$\partial^\alpha v(\xi) = \langle T_x, (-ix)^\alpha e^{-ix \cdot \xi} \rangle,$$

where, for some constant $C > 0$, an integer $m \in \mathbb{N}$ and a compact set K that depend only on T ,

$$|\partial^\alpha v(\xi)| \leq C \sum_{|\beta| \leq m} \sup_{x \in K} |\partial_x^\beta ((-ix)^\alpha e^{-ix \cdot \xi})| \leq C_\alpha (1 + |\xi|)^m$$

Last, we have $\hat{T} = v$. Indeed, for $\varphi \in C_0^\infty(\mathbb{R}^d)$, thanks to the result about integrating under the bracket,

$$\langle \hat{T}, \varphi \rangle = \langle T_x, \chi(x) \int e^{-ix \cdot \xi} \varphi(\xi) d\xi \rangle = \int \langle T_x, \chi(x) e^{-ix \cdot \xi} \rangle \varphi(\xi) d\xi,$$

where $\hat{T}(\xi) = \langle T_x, \chi(x) e^{-ix \cdot \xi} \rangle$. □

3.4.4 Fourier characterization of Sobolev spaces on \mathbb{R}^d

We recall that Sobolev spaces $H^s(\Omega)$ were introduced in Chapter 2 for $s \in \mathbb{N}$ and Ω an open subset of \mathbb{R}^d . In this subsection, we use the Fourier transform to provide more information about these spaces when $\Omega = \mathbb{R}^d$.

For $\xi \in \mathbb{R}^d$, we set $\langle \xi \rangle = \sqrt{1 + |\xi|^2}$. The function $\xi \mapsto \langle \xi \rangle$ is smooth on \mathbb{R}^d , and there is a constant $C > 0$ such that, for $|\xi| \geq 1$,

$$\frac{1}{C} |\xi| \leq \langle \xi \rangle \leq C |\xi|.$$

Thus, $\langle \xi \rangle$ is a regularized version of $|\xi|$, in the sense that it has the same behavior at infinity. Furthermore, since this function is smooth and since all its derivatives have growth at most polynomial at infinity, it acts on $\mathcal{S}'(\mathbb{R}^d)$ by multiplication.

Proposition 3.4.9 Let $s \in \mathbb{N}$ and $u \in \mathcal{S}'(\mathbb{R}^d)$. Then $u \in H^s(\mathbb{R}^d)$ if and only if $\langle \xi \rangle^s \hat{u} \in L^2(\mathbb{R}^d)$. Furthermore, there exists $C_s > 0$ such that

$$\forall u \in H^s(\mathbb{R}^d), \quad C_s^{-1} \|\langle \xi \rangle^s \hat{u}\|_{L^2} \leq \|u\|_{H^s} \leq C_s \|\langle \xi \rangle^s \hat{u}\|_{L^2}.$$

Proof.— If $u \in H^s(\mathbb{R}^d)$, then $\partial^\alpha u \in L^2(\mathbb{R}^d)$ for $|\alpha| \leq s$. By the Plancherel theorem, this means

$$\xi^\alpha \hat{u} \in L^2(\mathbb{R}^d), \quad |\alpha| \leq s.$$

Now observe that

$$\langle \xi \rangle^{2s} = (1 + \xi_1^2 + \cdots + \xi_d^2)^s = \sum_{|\alpha| \leq s} c_\alpha \xi^{2\alpha}$$

for some positive constants c_α . Consequently,

$$\langle \xi \rangle^{2s} |\hat{u}(\xi)|^2 = \sum_{|\alpha| \leq s} c_\alpha \xi^{2\alpha} |\hat{u}(\xi)|^2 \in L^1(\mathbb{R}^d),$$

whence $\langle \xi \rangle^s \hat{u} \in L^2(\mathbb{R}^d)$.

Conversely, if $\langle \xi \rangle^s \hat{u} \in L^2(\mathbb{R}^d)$, then, for $|\alpha| \leq s$, $\xi^\alpha \langle \xi \rangle^{-s}$ is bounded on \mathbb{R}^d and we have

$$\xi^\alpha \hat{u} = \xi^\alpha \langle \xi \rangle^{-s} \langle \xi \rangle^s \hat{u} \in L^2(\mathbb{R}^d),$$

which precisely means that $\partial^\alpha u \in L^2(\mathbb{R}^d)$.

The inequality follows from these considerations and from the Plancherel formula. \square

Proposition 3.4.9 suggests to extend the definition of $H^s(\mathbb{R}^d)$ to every real number s as follows.

Definition 3.4.10 Let $s \in \mathbb{R}$. A tempered distribution $u \in \mathcal{S}'(\mathbb{R}^d)$ belongs to $H^s(\mathbb{R}^d)$ if $\langle \xi \rangle^s \hat{u} \in L^2(\mathbb{R}^d)$.

Example 3.4.11 *i)* $\delta_0 \in H^s(\mathbb{R}^d)$ if and only if $s < \frac{-d}{2}$. Indeed $\hat{\delta}_0 = 1$, so that $\langle \xi \rangle^s \hat{\delta}_0 \in L^2(\mathbb{R}^d)$ if and only if $2s < -d$.

ii) Constant functions do not belong to $H^s(\mathbb{R}^d)$, since $\hat{1} = (2\pi)^d \delta_0$ is not in L^1_{loc} .

Here follows another illustration of the fact that H^s contains elements that are more and more singular as s decreases.

Proposition 3.4.12 Let $T \in \mathcal{E}'(\mathbb{R}^d)$ be a compactly supported distribution, with order $m \geq 0$. Then $T \in H^s(\mathbb{R}^d)$ for any $s < -m - \frac{d}{2}$.

Proof.— For $T \in \mathcal{E}'(\mathbb{R}^d)$, we know that $\hat{T} \in \mathcal{C}^\infty$. Moreover

$$|\langle \xi \rangle^s \hat{T}(\xi)| = |\langle \xi \rangle^s \langle T_x, e^{-ix \cdot \xi} \rangle| \leq C \langle \xi \rangle^s \sum_{|\alpha| \leq m} \sup |\partial_x^\alpha (e^{-ix \cdot \xi})| \leq C \langle \xi \rangle^{s+m}$$

Thus $T \in H^s(\mathbb{R}^d)$ when $2(s+m) < -d$, as stated. \square

We close this subsection by proving a connection between H^s regularity and \mathcal{C}^k regularity.

We denote by $\mathcal{C}^k_{\rightarrow 0}(\mathbb{R}^d)$ the space of \mathcal{C}^k functions on \mathbb{R}^d that tends to 0 at infinity, as well as all their derivatives of order $\leq k$.

Proposition 3.4.13 (Sobolev embedding) If $s > \frac{d}{2} + k$, then every element of $H^s(\mathbb{R}^d)$ belongs to $\mathcal{C}^k_{\rightarrow 0}(\mathbb{R}^d)$, and the embedding $H^s(\mathbb{R}^d) \rightarrow \mathcal{C}^k_{\rightarrow 0}(\mathbb{R}^d)$ is continuous.

Proof.— Let $u \in H^s(\mathbb{R}^d)$. For $\alpha \in \mathbb{N}^d$ such that $|\alpha| \leq k$, we have $\xi^\alpha \hat{u} \in L^1$. Indeed,

$$|\xi^\alpha \hat{u}(\xi)| \leq \frac{|\xi|^{|\alpha|}}{\langle \xi \rangle^s} \langle \xi \rangle^s |\hat{u}(\xi)| \leq \langle \xi \rangle^{k-s} \langle \xi \rangle^s |\hat{u}(\xi)|,$$

and $\langle \xi \rangle^{k-s} \in L^2(\mathbb{R}^d)$ since $-2(k-s) > d$. By the Cauchy-Schwarz inequality, we thus get

$$(3.4.5) \quad \|\xi^\alpha \hat{u}\|_{L^1} \leq C_{s,n} \|u\|_{H^s}.$$

Therefore $\partial^\alpha u = \mathcal{F}^{-1}((i\xi)^\alpha \hat{u}) \in \mathcal{C}_{\rightarrow 0}^0$ by Proposition 3.4.5, and the fact that the identity from $H^s(\mathbb{R}^d)$ to $\mathcal{C}_{\rightarrow 0}^k(\mathbb{R}^d)$ is continuous is just a way to read the inequalities

$$\forall |\alpha| \leq k, \|\partial^\alpha u\|_{L^\infty} \leq \|\xi^\alpha \hat{u}\|_{L^1} \leq C_{s,n} \|u\|_{H^s}.$$

□ Another interesting consequence of Proposition 3.4.9 is that we recover Theorem 2.5.4 on \mathbb{R}^d in a very general way, with optimal regularity of the solution.

Proposition 3.4.14 For every $s \in \mathbb{R}^d$, the mapping $u \in H^s(\mathbb{R}^s) \mapsto u - \Delta u \in H^{s-2}(\mathbb{R}^d)$ is an isomorphism.

Proof.— Indeed, this mapping is conjugated through Fourier transform to the mapping

$$\hat{u} \mapsto \langle \xi \rangle^2 \hat{u}$$

so that the statement becomes trivial in view of Proposition 3.4.9 or the subsequent definition of $H^s(\mathbb{R}^d)$. □

Combining the above three propositions, we obtain the following extension of Theorem 1.9.16 to several dimensions.

Theorem 3.4.15 For every distribution T on \mathbb{R}^d with compact support, there exist $p \in \mathbb{N}$ and a family $(f_\alpha)_{|\alpha| \leq p}$ of continuous functions on \mathbb{R}^d such that

$$T = \sum_{|\alpha| \leq p} \partial^\alpha (f_\alpha).$$

Proof.— From Proposition 3.4.12, we know that, for some $s \in \mathbb{R}$, $T \in H^s$. Let q be a positive integer such that

$$s + 2q > \frac{d}{2}.$$

By Proposition 3.4.14 applied q times, there exists $g \in H^{s+2q}$ unique such that

$$(I - \Delta)^q g = T.$$

By Proposition 3.4.13, we know that g is continuous. It remains to write

$$T = (I - \Delta)^q g = \left(1 - \sum_{j=1}^d \partial_j^2 \right)^q g = \sum_{|\beta| \leq q} c_\beta \partial^{2\beta} g$$

and the proof is complete. □

Finally, we introduce spaces of distributions in an arbitrary open set of \mathbb{R}^d which will be very useful in studying local regularity of solutions of partial differential equations.

Definition 3.4.16 Let Ω be an open subset of \mathbb{R}^d and let $s \in \mathbb{R}$. We denote by $H_{\text{loc}}^s(\Omega)$ the space of distributions $T \in \mathcal{D}'(\Omega)$ such that, for any $\chi \in \mathcal{C}_0^\infty(\Omega)$, the extension to \mathbb{R}^d by 0 of the compactly supported distribution χu belongs to $H^s(\mathbb{R}^d)$.

We notice the following important regularity theorem.

Proposition 3.4.17

$$\bigcap_{s \in \mathbb{R}} H_{\text{loc}}^s(\Omega) = \mathcal{C}^\infty(\Omega) .$$

Proof.— If $u \in \mathcal{C}^\infty(\Omega)$, then, for any $\chi \in \mathcal{C}_0^\infty(\Omega)$, $\chi u \in \mathcal{C}_0^\infty(\Omega)$, so that its extension to \mathbb{R}^d by 0 is such that $\widehat{\chi u}$ belongs to $\mathcal{S}(\mathbb{R}^d)$, and consequently $\chi u \in H^s$ for every s .

Conversely, if $u \in H_{\text{loc}}^s(\Omega)$ for every $s \in \mathbb{R}$, then, any $\chi \in \mathcal{C}_0^\infty(\Omega)$, $\chi u \in H^s(\mathbb{R}^d)$ for every s , hence, from Proposition 3.4.13, $\chi u \in \mathcal{C}_{\rightarrow 0}^k(\mathbb{R}^d)$ for every k , in particular $\chi u \in \mathcal{C}^\infty(\mathbb{R}^d)$. Since this holds for any $\chi \in \mathcal{C}_0^\infty(\Omega)$, this completes the proof. \square

3.5 Some applications

3.5.1 Partial differential equations with constant coefficients

Let $p \in \mathbb{C}[X_1, \dots, X_d]$ be a polynomial of n variables with complex coefficients,

$$p(X) = \sum_{|\alpha| \leq m} a_\alpha X^\alpha, \quad X \in \mathbb{R}^d,$$

with $a_\alpha \in \mathbb{C}$. The integer $m \in \mathbb{N}$ is the degree of P . Given an open subset Ω of \mathbb{R}^d , we denote by $p(\partial)$ the operator on $\mathcal{D}'(\Omega)$ given by

$$\mathcal{D}'(\Omega) \ni T \mapsto p(\partial)T = \sum_{|\alpha| \leq m} a_\alpha \partial^\alpha T \in \mathcal{D}'(\Omega).$$

Those operators are called linear partial differential operators with constant coefficients. The equation

$$p(\partial)u = f,$$

where $f \in \mathcal{D}'(\Omega)$ is given, and $u \in \mathcal{D}'(\Omega)$ is the unknown, is called a Partial Differential Equation (PDE) of order m with constant coefficients. When $f \neq 0$, it is said to be inhomogeneous, or with source term f .

Remark 3.5.1 For $d = 1$, the equation $p(\partial)u = f$ is a linear differential equation of order m with constant coefficients, that can be explicitly solved. For $d > 1$, the situation is drastically different and it may even be very difficult to show existence of solutions.

Example 3.5.2 If $p(X_1, \dots, X_d) = X_1^2 + \dots + X_d^2$, then

$$p(\partial)T = \partial_1^2 T + \dots + \partial_d^2 T = \Delta T .$$

We already encountered operator Δ which is called the Laplacean, and equation $\Delta u = f$ is called the Laplace (or Poisson) equation. A specificity of Δ is that it is invariant by the rotations of \mathbb{R}^d . Moreover, one can show that every linear partial differential operator with constant coefficients on \mathbb{R}^d which is invariant by rotations, is of the form

$$P = Q(\Delta)$$

where Q is a polynomial of one variable. This invariance property explains why the Laplacean appears so frequently in many areas of Mathematical Physics.

The following proposition is an easy consequence of properties of the Fourier transform on $\mathcal{S}'(\mathbb{R}^d)$

Proposition 3.5.3 Let $p \in \mathbb{C}[X_1, \dots, X_d]$ and $P = p(\partial)$ be the corresponding differential operator. For every $u \in \mathcal{S}'(\mathbb{R}^d)$, we have

$$\widehat{Pu} = p(i\xi)\hat{u} .$$

The importance of the above proposition appears through the multiplicity of its consequences. We shall draw some of them in the next subsections.

3.5.2 Local results

Let $P = p(\partial)$ be a linear partial differential operators with constant coefficients. In this section, we study some properties of P on $\mathcal{D}'(\Omega)$, where Ω is an open subset of \mathbb{R}^d . We start with a very natural statement.

Proposition 3.5.4 If P is a linear partial differential operators with constant coefficients of order $\leq m$ and $s \in \mathbb{R}$, then $P : H_{\text{loc}}^s(\Omega) \rightarrow H_{\text{loc}}^{s-m}(\Omega)$.

Proof.— It is based on the following lemma.

Lemma 3.5.5 If P is a linear partial differential operators with constant coefficients of order $\leq m$ and $\varphi \in \mathcal{C}_0^\infty(\Omega)$, there exists functions $\varphi_\beta \in \mathcal{C}_0^\infty(\Omega)$, $|\beta| \leq m - 1$, such that, for every $u \in \mathcal{D}'(\Omega)$,

$$P(\varphi u) = \varphi Pu + \sum_{|\beta| \leq m-1} \partial^\beta (\varphi_\beta u) .$$

Let us prove this lemma. By linearity, it is enough to prove it for $P = \partial^\alpha$. We proceed by induction on $|\alpha|$. If $\alpha = 0$, the formula trivially holds. Assume it is true for $|\alpha| \leq m - 1$, and let us prove it for $|\alpha| = m$. By the Leibniz formula,

$$\partial^\alpha (\varphi u) = \varphi \partial^\alpha u + \sum_{\beta \leq \alpha, \beta \neq \alpha} \binom{\alpha}{\beta} \partial^{\alpha-\beta} \varphi \partial^\beta u .$$

Using the induction hypothesis, we may write each term in the second part of the right hand side as a sum of terms $\partial^\gamma(\psi_\gamma u)$, with $|\gamma| \leq m - 1$. This completes the proof of Lemma 3.5.5.

Coming back to the proof of Proposition 3.5.4, we use Lemma 3.5.5 to write

$$P(\varphi u) = \varphi Pu + \sum_{|\beta| \leq m-1} \partial^\beta(\varphi_\beta u).$$

If $u \in H_{\text{loc}}^s(\Omega)$, then $\varphi u, \varphi_\beta u \in H^s(\mathbb{R}^d)$ and thus

$$P(\varphi u) \in H^{s-m}(\mathbb{R}^d), \partial^\beta(\varphi_\beta u) \in H^{s-m+1}(\mathbb{R}^d).$$

This leads to $\varphi Pu \in H^{s-m}(\mathbb{R}^d)$, so that $Pu \in H_{\text{loc}}^{s-m}(\mathbb{R}^d)$. \square

A classical problem is the local regularity of solutions $u \in \mathcal{D}'(\Omega)$ of equation $p(\partial)u = f$, knowing the local regularity of f . It is possible to give a general answer to this question under some assumption on p , which we now define.

Definition 3.5.6 Operator $p(\partial) = \sum_{|\alpha| \leq m} a_\alpha \partial^\alpha$ is called elliptic of order m if the highest homogeneous part $p_m = \sum_{|\alpha|=m} a_\alpha X^\alpha$ of polynomial p satisfies

$$\forall \xi \in \mathbb{R}^d \setminus \{0\}, p_m(\xi) \neq 0.$$

For instance, the Laplace operator is elliptic, while the operator $\partial_1^2 - \partial_2^2$ is not.

Theorem 3.5.7 Let P be a linear partial differential operators with constant coefficients, elliptic of order m , and let Ω be an open subset of \mathbb{R}^d . If $u \in \mathcal{D}'(\Omega)$ satisfies $Pu \in H_{\text{loc}}^s(\Omega)$ for some $s \in \mathbb{R}$, then $u \in H_{\text{loc}}^{s+m}(\Omega)$. In particular, if $Pu \in C^\infty(\Omega)$, then $u \in C^\infty(\Omega)$.

Proof.— Let us first prove the result under the additional assumption that $u \in H_{\text{loc}}^\sigma(\Omega)$ for some $\sigma \in \mathbb{R}$. Since p is elliptic and since the unit sphere is compact, there exists $c > 0$ such that, for every $\xi \in \mathbb{R}^d$ with $|\xi| = 1$,

$$|p_m(i\xi)| \geq c.$$

By homogeneity, this implies

$$\forall \xi \in \mathbb{R}^d, |p_m(\xi)| \geq c|\xi|^m.$$

Since

$$p(i\xi) - i^m p_m(i\xi) = O(|\xi|^{m-1})$$

as $|\xi| \rightarrow \infty$, we infer that there exists $R > 0$ such that, for every $\xi \in \mathbb{R}^d$ with $|\xi| \geq R$,

$$(3.5.6) \quad |p(i\xi)| \geq \frac{c}{2} |\xi|^m.$$

Let $\chi \in C_0^\infty(\mathbb{R}^d)$ be a plateau function on the ball $B(0, R)$. The function

$$q(\xi) = \frac{1 - \chi(\xi)}{p(i\xi)}$$

is C^∞ on \mathbb{R}^d and satisfies, due to estimate (3.5.6),

$$|q(\xi)| = O(\langle \xi \rangle^{-m}).$$

Therefore we can define an operator $Q : H^\tau(\mathbb{R}^d) \rightarrow H^{\tau+m}(\mathbb{R}^d)$ for every $\tau \in \mathbb{R}$ by

$$\widehat{Q(T)} = q\widehat{T}.$$

Now let $u \in H_{\text{loc}}^\sigma(\Omega)$ such that $Pu \in H_{\text{loc}}^s(\Omega)$. For every $\varphi \in C_0^\infty(\Omega)$, Lemma 3.5.5 yields

$$P(\varphi u) = \varphi Pu + \sum_{|\alpha| \leq m-1} \partial^\alpha(\varphi_\alpha u),$$

where $\varphi_\alpha \in C_0^\infty(\Omega)$. Consequently,

$$QP(\varphi u) = Q(\varphi Pu) + \sum_{|\alpha| \leq m-1} Q(\partial^\alpha(\varphi_\alpha u)).$$

In the right hand side, the first term belongs to $H^{s+m}(\mathbb{R}^d)$ while the other terms belong to $H^{\sigma+1}(\mathbb{R}^d)$. By Proposition 3.5.3, $QP(\varphi u) = \varphi u - R(\varphi u)$, where

$$\widehat{Rv} = \chi\widehat{v},$$

so that $R(\varphi u) \in H^\tau(\mathbb{R}^d)$ for every $\tau \in \mathbb{R}$. We infer that $\varphi u \in H^{\min(s+m, \sigma+1)}(\mathbb{R}^d)$, and so

$$u \in H_{\text{loc}}^{\min(s+m, \sigma+1)}(\Omega).$$

If $\sigma + 1 < s + m$, we iterate this result replacing σ by $\sigma + 1$, and, after a sufficient number of iterations, we conclude

$$u \in H_{\text{loc}}^{s+m}(\Omega).$$

Now let $u \in \mathcal{D}'(\Omega)$ such that $Pu \in H_{\text{loc}}^s(\Omega)$. Let V be an open subset of Ω such that $\overline{V} \subset \Omega$ is compact. If $\psi \in C_0^\infty(\Omega)$ is a plateau function on \overline{V} , Proposition 3.4.12 implies that there exists $\sigma \in \mathbb{R}$ such that $\psi u \in H^\sigma(\mathbb{R}^d)$. The following lemma implies that

$$u|_V \in H_{\text{loc}}^\sigma(V).$$

Lemma 3.5.8 If $\sigma \in \mathbb{R}$, $v \in H^\sigma(\mathbb{R}^d)$ and $\varphi \in C_0^\infty(\mathbb{R}^d)$, then $\varphi v \in H^\sigma(\mathbb{R}^d)$.

Assuming this lemma, we can apply to $u|_V$ the first part of the proof and conclude that $u \in H_{\text{loc}}^{s+m}(V)$. Since V is an arbitrary open subset of Ω such that $\overline{V} \subset \Omega$ is compact, we conclude that $u \in H_{\text{loc}}^{s+m}(\Omega)$.

Let us prove Lemma 3.5.8. We have, from Proposition 3.4.8,

$$\widehat{\varphi v}(\xi) = \langle v, \varphi e^{-ix \cdot \xi} \rangle.$$

Using the inversion formula and the properties of the Fourier transform on \mathcal{S} ,

$$\langle v, \varphi e^{-ix \cdot \xi} \rangle = \langle \hat{v}, \hat{\varphi}(\xi - \cdot) \rangle = \int_{\mathbb{R}^d} \hat{v}(\eta) \hat{\varphi}(\xi - \eta) d\eta .$$

This leads to

$$\langle \xi \rangle^\sigma \widehat{\varphi v}(\xi) = \langle \xi \rangle^\sigma \int_{\mathbb{R}^d} \hat{v}(\eta) \hat{\varphi}(\xi - \eta) d\eta = \langle \xi \rangle^\sigma \int_{\mathbb{R}^d} \hat{v}(\xi - \eta) \hat{\varphi}(\eta) d\eta,$$

and we have to prove that the right hand side is square integrable in ξ on \mathbb{R}^d . We decompose the integral into two parts, corresponding to the domains $|\eta| \leq |\xi|/2$ and $|\eta| > |\xi|/2$. Using the rapid decay of $\hat{\varphi}$, the second integral is rapidly decreasing in ξ . As for the first integral, we apply the Cauchy–Schwarz inequality,

$$\langle \xi \rangle^{2\sigma} \left| \int_{|\eta| \leq |\xi|/2} \hat{v}(\xi - \eta) \hat{\varphi}(\eta) d\eta \right|^2 \leq \left(\int_{\mathbb{R}^d} |\hat{\varphi}(\eta)| d\eta \right) \left(\int_{|\eta| \leq |\xi|/2} \langle \xi \rangle^{2\sigma} |\hat{v}(\xi - \eta)|^2 |\hat{\varphi}(\eta)| d\eta \right) .$$

Since $|\eta| \leq |\xi|/2$, we have $\langle \xi \rangle \leq A \langle \xi - \eta \rangle$ for some constant $A > 0$, and therefore the right hand side is bounded by

$$B \int_{\mathbb{R}^d} \langle \xi - \eta \rangle^{2\sigma} |\hat{v}(\xi - \eta)|^2 |\hat{\varphi}(\eta)| d\eta ,$$

whose integral with respect to ξ is finite by Fubini's theorem. This completes the proof. \square

3.5.3 Global results

We start with solutions of homogeneous equations.

Corollary 3.5.9 With the notation of Proposition 3.5.3, we have

- i) If $p(i\xi) \neq 0$ for every $\xi \in \mathbb{R}^d$, then the only tempered solution u of $Pu = 0$ is $u = 0$.
- ii) If $p(i\xi) \neq 0$ for every $\xi \in \mathbb{R}^d \setminus \{0\}$, then every tempered solution of $Pu = 0$ is a polynomial function.

Proof.— If $u \in \mathcal{S}'$ satisfies $Pu = 0$, then $p(i\xi)\hat{u} = 0$, which implies that the distribution \hat{u} is supported in the set

$$\{\xi \in \mathbb{R}^d, p(i\xi) = 0\}$$

If this set is empty, this implies $\hat{u} = 0$, hence $u = 0$, whence (i).

If this set is $\{0\}$, this implies, from the structure of the distributions supported at one point only, that \hat{u} is a finite linear combination of derivatives of δ_0 . Applying \mathcal{F}^{-1} , we conclude that u is a polynomial function, which is (ii). \square

Remark 3.5.10 A consequence of Corollary 3.5.9 is that, under assumption ii), bounded solutions of $Pu = 0$ on \mathbb{R}^d are constants. Indeed, since $L^\infty(\mathbb{R}^d) \subset \mathcal{S}'(\mathbb{R}^d)$, this follows from the elementary fact claiming that a polynomial function which is bounded on \mathbb{R}^d is constant. This is trivial if $d = 1$, but less trivial if $d \geq 2$. Let us sketch a proof of it. Let $p \in \mathbb{C}[X_1, \dots, X_d]$ be such that function $x \in \mathbb{R}^d \mapsto p(x) \in \mathbb{C}$ is bounded. For every $y \in \mathbb{R}^d$, consider

$$p_y(t) = p(ty) , t \in \mathbb{R} .$$

Function p_y is a one variable polynomial function, which is bounded, hence it is a constant. Writing

$$p(X) = \sum_{|\alpha| \leq m} a_\alpha X^\alpha,$$

this implies, for every $r = 1, \dots, m$, for every $y \in \mathbb{R}^d$,

$$\sum_{|\alpha|=r} a_\alpha y^\alpha = 0.$$

Taking the ∂^α derivative of the left hand side, we infer $a_\alpha = 0$.

Example 3.5.11 ▪ Case (i) is fulfilled by $P = \Delta + \lambda$ if $\lambda \in \mathbb{C} \setminus \mathbb{R}_+$.

- Case (ii) is fulfilled by $P = \Delta$ on \mathbb{R}^d . Consequently, tempered harmonic functions on \mathbb{R}^d are polynomial functions. In particular, we infer the strong Liouville theorem : any bounded harmonic function on \mathbb{R}^d is a constant.

Case (ii) is also fulfilled by $P = \partial_1 + i\partial_2$ on \mathbb{R}^2 . In other words, the only tempered entire functions on \mathbb{C} are polynomial, and again (Liouville), we recover that the only bounded entire functions are constant.

The second application concerns fundamental solutions.

Definition 3.5.12 Let $P = p(\partial)$ be a linear partial differential operators with constant coefficients. One says that $E \in \mathcal{D}'(\mathbb{R}^d)$ is a fundamental solution of P when $PE = \delta_0$.

Notice that in Physics, fundamental solutions are often called Green functions. The importance of fundamental solutions is provided by the following remark.

Proposition 3.5.13 If P has a fundamental solution $E \in \mathcal{D}'(\mathbb{R}^d)$, then for all $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$, the equation $Pu = \varphi$ has a solution given by $u = E * \varphi$.

Proof.— In chapter 2, we saw that $\partial_j(T * \varphi) = T * \partial_j \varphi = \partial_j T * \varphi$, hence

$$P(E * \varphi) = (PE) * \varphi = \delta * \varphi = \varphi,$$

so that $u = E * \varphi$ is a solution of $Pu = \varphi$. □

Using a convenient definition of convolution of distributions, it is in fact possible to extend the above proposition to any right hand side which is a compactly supported distribution. We shall see an example of this in Theorem 3.5.17 below.

B. Malgrange and L. Ehrenpreis have proved, independently in 1954/1955, that any non trivial linear partial differential operator with constant coefficients has a fundamental solution. It is fact possible to prove that one can choose this distribution to be tempered. The proofs of these results are beyond the scope of these lectures. In what follows, we rather study the important case of the Laplace operator.

Proposition 3.5.14 Let $d \geq 3$, and let $E_d \in \mathcal{D}'(\mathbb{R}^d)$ be the $L^1_{\text{loc}}(\mathbb{R}^d)$ function given by

$$E_d(x) = \frac{1}{(d-2)\sigma(S^{d-1})|x|^{d-2}}.$$

Then

$$-\Delta E_d = \delta_0.$$

Proof.— We are going to use Fourier transformation. Indeed, by Proposition 3.5.3, in $\mathcal{S}'(\mathbb{R}^d)$, the equation

$$-\Delta E = \delta_0$$

is equivalent to

$$|\xi|^2 \hat{E} = 1.$$

Since $d \geq 3$, the function $1/|\xi|^2$ is locally integrable in \mathbb{R}^d . More precisely, by decomposing it as the sum of the contributions for $|\xi| \leq 1$ and for $|\xi| > 1$, one shows that this function is the sum of an L^1 function and of an L^∞ function. As a consequence, $1/|\xi|^2$ belongs to $\mathcal{S}'(\mathbb{R}^d)$, and therefore

$$E_d := \mathcal{F}^{-1} \left(\frac{1}{|\xi|^2} \right)$$

is a tempered fundamental solution of $-\Delta$. It remains to calculate E_d . Since $-\Delta E_d = 0$ on $\mathbb{R}^d \setminus \{0\}$, we know from Theorem 3.5.7 that E_d is a C^∞ function on $\mathbb{R}^d \setminus \{0\}$. We claim that this function is of the form

$$E_d(x) = \frac{c_d}{|x|^{d-2}}$$

for some constant c_d . This is equivalent to the following statement : for every rotation R of \mathbb{R}^d , for every $\lambda > 0$, for every $x \in \mathbb{R}^d \setminus \{0\}$,

$$E_d(\lambda R x) = \lambda^{2-d} E_d(x),$$

or equivalently, for every $\varphi \in C_0^\infty(\mathbb{R}^d \setminus \{0\})$,

$$(3.5.7) \quad \int_{\mathbb{R}^d} E_d(\lambda R x) \varphi(x) dx = \lambda^{2-d} \int_{\mathbb{R}^d} E_d(x) \varphi(x) dx.$$

Let $Ax := \lambda R x$ and $\varphi_A(x) := |\det A|^{-1} \varphi(A^{-1}x)$. We have

$$\begin{aligned} \int_{\mathbb{R}^d} E_d(Ax) \varphi(x) dx &= \int_{\mathbb{R}^d} E_d(x) \varphi_A(x) dx \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} \frac{\widehat{\varphi}_A(\xi)}{|\xi|^2} d\xi \\ \int_{\mathbb{R}^d} E_d(x) \varphi(x) dx &= (2\pi)^{-d} \int_{\mathbb{R}^d} \frac{\widehat{\varphi}(\xi)}{|\xi|^2} d\xi. \end{aligned}$$

Furthermore,

$$\widehat{\varphi}_A(\xi) = \int_{\mathbb{R}^d} |\det A|^{-1} \varphi(A^{-1}x) e^{-ix \cdot \xi} dx = \int_{\mathbb{R}^d} \varphi(x) e^{-iAx \cdot \xi} dx = \widehat{\varphi}({}^t A \xi).$$

Consequently,

$$\int_{\mathbb{R}^d} \frac{\widehat{\varphi}_A(\xi)}{|\xi|^2} d\xi = \int_{\mathbb{R}^d} \frac{\varphi({}^t A\xi)}{|\xi|^2} d\xi = |\det A|^{-1} \int_{\mathbb{R}^d} \frac{\varphi(\xi)}{|{}^t A^{-1}\xi|^2} d\xi.$$

Since ${}^t A^{-1} = \lambda^{-1}R$, we conclude

$$\int_{\mathbb{R}^d} \frac{\widehat{\varphi}_A(\xi)}{|\xi|^2} d\xi = \lambda^{2-d} \int_{\mathbb{R}^d} \frac{\widehat{\varphi}(\xi)}{|\xi|^2} d\xi$$

and identity (3.5.7) is proved.

In order to calculate c_d , it is enough to check against the Gaussian function G_1 ,

$$c_d \int_{\mathbb{R}^d} \frac{G_1(x)}{|x|^{d-2}} dx = (2\pi)^{-d} \int_{\mathbb{R}^d} \frac{\widehat{G}_1(\xi)}{|\xi|^2} d\xi.$$

Since $\widehat{G}_1(\xi) = (2\pi)^{d/2}G_1(\xi)$, we infer

$$c_d \int_{\mathbb{R}^d} \frac{e^{-|x|^2/2}}{|x|^{d-2}} dx = (2\pi)^{-d/2} \int_{\mathbb{R}^d} \frac{e^{-|\xi|^2/2}}{|\xi|^2} d\xi.$$

Passing in spherical coordinates, this reads

$$c_d \int_0^\infty r e^{-r^2/2} dr = (2\pi)^{-d/2} \int_0^\infty r^{d-3} e^{-r^2/2} dr,$$

or

$$c_d = (2\pi)^{-d/2} 2^{(d-4)/2} \int_0^\infty t^{(d-4)/2} e^{-t} dt = \frac{1}{4} \pi^{-d/2} \Gamma\left(\frac{d}{2} - 1\right)$$

Recall that, by applying the integration formula on spherical level sets from Corollary 2.4.8 to a Gaussian function, one gets

$$\sigma(S^{d-1}) = \frac{2\pi^{d/2}}{\Gamma(d/2)} = \frac{4\pi^{d/2}}{(d-2)\Gamma(d/2-1)}$$

so that

$$c_d = \frac{1}{(d-2)\sigma(S^{d-1})}.$$

□

Remark 3.5.15 The value of c_d can also be determined by applying the Gauss–Green formula to the integral of $|x|^{2-d}\Delta\varphi(x)$ outside a ball of radius ε .

Remark 3.5.16 It is possible to extend Proposition 3.5.14 to $d = 1, 2$. For $d = 1$, it is easy to check that

$$E_1(x) = -\frac{1}{2}|x|$$

is a fundamental solution of $-\partial^2$. The case $d = 2$ is more delicate. The above approach by Fourier analysis has to be slightly modified, because $1/|\xi|^2$ is not locally integrable in \mathbb{R}^2 . A good way to do this is to consider the operator

$$P = \partial_1 + i\partial_2.$$

Since $1/(\xi_1 + i\xi_2) \in L^1_{\text{loc}}(\mathbb{R}^2)$, the distribution $E \in \mathcal{S}'(\mathbb{R}^2)$ such that

$$\widehat{E} = \frac{1}{i(\xi_1 + i\xi_2)}$$

satisfies $PE = \delta_0$. Furthermore, from Theorem 3.5.7, E is a holomorphic function outside $\{0\}$, and the above proof shows that it is homogeneous of degree -1 . This finally leads to

$$E(x) = \frac{1}{2\pi(x_1 + ix_2)}$$

and, observing that

$$E = (\partial_1 - i\partial_2)[(4\pi)^{-1} \log(x_1^2 + x_2^2)]$$

in $\mathcal{D}'(\mathbb{R}^2)$, we conclude that

$$E_2 = -\frac{1}{2\pi} \log|x|$$

is a fundamental solution of $-\Delta$ on \mathbb{R}^2 .

Our next and final step is to study equation $-\Delta u = f$ when f is an arbitrary compactly supported distribution on \mathbb{R}^2 . Notice that, from Theorem 3.5.7, every solution u of this equation is a C^∞ function on the complement of $\text{supp}(f)$. For simplicity, we shall restrict ourselves to the case $d = 3$, for which this equation is the Poisson equation of electrostatics.

Theorem 3.5.17 For every compactly supported distribution f on \mathbb{R}^3 , there exists a unique distribution u on \mathbb{R}^3 satisfying $-\Delta u = f$ and $u(x) \rightarrow 0$ as $x \rightarrow \infty$. Furthermore, for $x \notin \text{supp}(f)$, $u(x)$ is given by

$$(3.5.8) \quad u(x) = \left\langle f, \frac{\chi}{4\pi|x - \cdot|} \right\rangle,$$

for every $\chi \in \mathcal{C}_0^\infty(\mathbb{R}^3)$ equal to 1 in a neighbourhood of $\text{supp}(f)$ and equal to 0 near x .

Proof.— The uniqueness of u is immediate in view of the Liouville theorem for harmonic functions – see Example 3.5.11. For the existence, we look for u in $\mathcal{S}'(\mathbb{R}^3)$, and Proposition 3.5.3 leads to equation

$$|\xi|^2 \hat{u} = \hat{f}.$$

Since \hat{f} is smooth on \mathbb{R}^3 with moderate growth and $1/|\xi|^2$ is the sum of an L^1 function near $\xi = 0$ and of an L^∞ function near infinity, we can define u by

$$\hat{u}(\xi) = \frac{\hat{f}(\xi)}{|\xi|^2}.$$

Our task is now to prove identity (3.5.8), since it implies that $u(x) \rightarrow 0$ as $x \rightarrow \infty$. As a first step, we are going to prove this identity if $f = \varphi \in \mathcal{C}_0^\infty(\mathbb{R}^3)$. In this case, denote by g the inverse Fourier transform of

$$\frac{\hat{\varphi}}{|\xi|^2}.$$

Since \hat{g} is integrable on \mathbb{R}^3 and rapidly decaying at infinity, we know that g is smooth and tends to 0 at infinity. Consider the following smooth function,

$$E_3 * \varphi(x) = \int_{\mathbb{R}^3} \frac{\varphi(y)}{4\pi|x - y|} dy.$$

Then

$$-\Delta(E_3 * \varphi) = \varphi = -\Delta g$$

and it is clear on the above expression that $E_3 * \varphi(x) \rightarrow 0$ as $x \rightarrow \infty$. Therefore, by the Liouville theorem, $g = E_3 * \varphi$.

Let us come to the general case of a compactly supported distribution f . Let $\rho \in \mathcal{C}^\infty(\mathbb{R}^3)$ with integral 1, and, for every $\varepsilon > 0$,

$$\rho_\varepsilon(x) = \frac{1}{\varepsilon^3} \rho\left(\frac{x}{\varepsilon}\right).$$

Then $\rho_\varepsilon * f = f_\varepsilon \in \mathcal{C}_0^\infty(\mathbb{R}^3)$ and converges to f in $\mathcal{D}'(\mathbb{R}^3)$ as $\varepsilon \rightarrow 0$. In fact, as we already observed in similar cases, f_ε is supported into a small neighbourhood of $\text{supp}(f)$, so that $\hat{f}_\varepsilon(\xi) \rightarrow \hat{f}(\xi)$ with a uniform moderate bound $(1 + |\xi|)^m$ for some m . Let $u_\varepsilon = E_3 * f_\varepsilon$. We know from the above identity that

$$\hat{u}_\varepsilon = \frac{\hat{f}_\varepsilon}{|\xi|^2}$$

and the right hand side converges to \hat{u} in $\mathcal{S}'(\mathbb{R}^3)$. Hence u_ε converges to u in $\mathcal{S}'(\mathbb{R}^3)$, and for every test function $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^3)$,

$$\begin{aligned} \langle u, \varphi \rangle &= \lim_{\varepsilon \rightarrow 0} \langle u_\varepsilon, \varphi \rangle = \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^3} E_3 * f_\varepsilon(x) \varphi(x) dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} E_3(x-y) f_\varepsilon(y) \varphi(x) dx dy \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^3} f_\varepsilon(y) E_3 * \varphi(y) dy \\ &= \langle f, \chi(E_3 * \varphi) \rangle \end{aligned}$$

for every $\chi \in \mathcal{C}_0^\infty(\mathbb{R}^3)$ equal to 1 in a neighbourhood of $\text{supp}(f)$. Notice that

$$\chi(y) E_3 * \varphi(y) = \int_{\mathbb{R}^3} \frac{\chi(y)}{4\pi|x-y|} \varphi(x) dx.$$

If $\text{supp}(\varphi) \cap \text{supp}(\chi) = \emptyset$, we can apply Proposition 2.3.20 of integration under the bracket to obtain

$$\langle u, \varphi \rangle = \langle f, \chi(E_3 * \varphi) \rangle = \int_{\mathbb{R}^3} \varphi(x) \left\langle f, \frac{\chi}{4\pi|x-\cdot|} \right\rangle dx$$

and this yields formula (3.5.8). □

Remark 3.5.18 Notice that formula (3.5.8) leads to a complete expansion of $u(x)$ as $x \rightarrow \infty$. In particular,

$$u(x) = \frac{q}{4\pi|x|} + O\left(\frac{1}{|x|^2}\right)$$

where $q = \langle f, \chi \rangle$ for every $\chi \in \mathcal{C}_0^\infty(\mathbb{R}^3)$ equal to 1 in a neighbourhood of $\text{supp}(f)$. This expansion supports the well known fact in Physics that any charge distribution f with nonzero total charge ($q = \langle f, 1 \rangle \neq 0$) can be seen at infinity as a point distribution with charge q .

We conclude this subsection by a remark about the generalization of the above approach to partial differential equations with non constant coefficients.

Remark 3.5.19 There are several directions where Fourier transformation can be used to study solutions of partial differential equations with non constant coefficients.

- i) The first and most direct topic concerns equations with affine coefficients. Indeed, the Fourier transform converts such equations into first order linear equations with polynomial coefficients, which can be solved in general. A famous example in dimension $d = 1$ is the Airy differential equation,

$$u''(x) = xu(x)$$

on \mathbb{R} . It is well known from the classical theory of differential equations that the space of solutions of this equation is contained into $\mathcal{C}^\infty(\mathbb{R})$ and is two-dimensional. Let us look for solutions u which are moreover tempered. Then the equation satisfied by \hat{u} is

$$(i\xi)^2 \hat{u} = i \frac{d}{d\xi} \hat{u},$$

or

$$\frac{d\hat{u}}{d\xi} = i\xi^2 \hat{u}.$$

The solutions in $\mathcal{D}'(\mathbb{R})$ are given by

$$\hat{u}(\xi) = c e^{i\frac{\xi^3}{3}}, \quad c \in \mathbb{C},$$

which are indeed tempered distributions, since in $L^\infty(\mathbb{R})$. Consequently, the space of tempered solutions of the Airy differential equation is one dimensional, generated by the Airy function,

$$Ai(x) = \mathcal{F}^{-1} \left(e^{i\frac{\xi^3}{3}} \right).$$

- ii) The second and much more general topic concerns the regularity theory of distribution solutions to partial differential equations with arbitrary smooth coefficients in arbitrary open subsets of \mathbb{R}^d , using cutoff functions in a smart way. This is the starting point of the theory of pseudo-differential operators, which is beyond the scope of this course.

3.6 Fourier series

In this subsection, we show how the theory of Fourier series can be considered as a special case of the Fourier transformation on tempered distributions. We start with a very natural definition. Recall that, for $a \in \mathbb{R}^d$ and $T \in \mathcal{D}'(\mathbb{R}^d)$, we have defined $\tau_a(T) \in \mathcal{D}'(\mathbb{R}^d)$ by

$$\forall \varphi \in \mathcal{D}(\mathbb{R}^d), \quad \langle \tau_a(T), \varphi \rangle = \langle T, \tau_{-a}(\varphi) \rangle,$$

and $\tau_b(\varphi)(x) := \varphi(x - b)$.

Definition 3.6.1 A distribution $T \in \mathcal{D}'(\mathbb{R}^d)$ is said to be \mathbb{Z}^d -periodic if

$$\forall \gamma \in \mathbb{Z}^d, \quad \tau_\gamma(T) = T.$$

Notice that derivatives of \mathbb{Z}^d -periodic distributions are \mathbb{Z}^d -periodic, as well as their product with $C^\infty \mathbb{Z}^d$ -periodic functions. A typical example of a \mathbb{Z}^d -periodic distribution is given by distributions defined by \mathbb{Z}^d -periodic locally integrable functions. Another example is

$$T = \sum_{\gamma \in \mathbb{Z}^d} \delta_\gamma .$$

The following proposition explains how to construct \mathbb{Z}^d -periodic locally integrable functions from L^1 functions on \mathbb{R}^d .

Proposition 3.6.2 Let $f \in L^1(\mathbb{R}^d)$. For almost every $x \in \mathbb{R}^d$, we have

$$\sum_{\gamma \in \mathbb{Z}^d} |f(x - \gamma)| < \infty ,$$

so that the formula

$$f^\sharp(x) := \sum_{\gamma \in \mathbb{Z}^d} f(x - \gamma)$$

defines a \mathbb{Z}^d -periodic locally integrable function on \mathbb{R}^d . Furthermore, if Q denotes the cube $[0, 1]^d$, the following identity holds,

$$(3.6.9) \quad \int_Q f^\sharp(x) dx = \int_{\mathbb{R}^d} f(x) dx .$$

If $\varphi \in \mathcal{S}(\mathbb{R}^d)$, then $\varphi^\sharp \in C^\infty(\mathbb{R}^d)$, $\partial^\alpha \varphi^\sharp = (\partial^\alpha \varphi)^\sharp$ and

$$(3.6.10) \quad \|\partial^\alpha \varphi^\sharp\|_{L^\infty} \leq C_d N_{d+1+|\alpha|}(\varphi) .$$

Furthermore, there exists $\chi \in C_0^\infty(\mathbb{R}^d)$ such that $\chi^\sharp = 1$.

Proof.— By the monotone convergence theorem,

$$\int_Q \sum_{\gamma \in \mathbb{Z}^d} |f(x - \gamma)| dx = \sum_{\gamma \in \mathbb{Z}^d} \int_Q |f(x - \gamma)| dx = \sum_{\gamma \in \mathbb{Z}^d} \int_{\gamma+Q} |f(y)| dy .$$

Since

$$\bigcup_{\gamma \in \mathbb{Z}^d} (\gamma + Q) = \mathbb{R}^d$$

and since the cubes $\gamma + Q$ are pairwise disjoint, we conclude

$$\sum_{\gamma \in \mathbb{Z}^d} \int_{\gamma+Q} |f(y)| dy = \int_{\mathbb{R}^d} |f(y)| dy < \infty .$$

Consequently, for almost every $x \in \mathbb{R}^d$,

$$\sum_{\gamma \in \mathbb{Z}^d} |f(x - \gamma)| < \infty ,$$

and so we can define $f^\sharp(x)$. Notice that, for every $\gamma_0 \in \mathbb{Z}^d$,

$$f^\sharp(x - \gamma_0) = \sum_{\gamma \in \mathbb{Z}^d} f(x - \gamma_0 - \gamma) = \sum_{\beta \in \mathbb{Z}^d} f(x - \beta) = f^\sharp(x).$$

Furthermore, by the above calculation,

$$\int_Q |f^\sharp(x)| dx \leq \int_{\mathbb{R}^d} |f(x)| dx < \infty,$$

so f^\sharp is a \mathbb{Z}^d -periodic locally integrable function on \mathbb{R}^d . Identity (3.6.9) follows from integrating the series defining f^\sharp , and from using again the disjoint covering of \mathbb{R}^d by the cubes $\gamma + Q$.

Let us suppose $\varphi \in \mathcal{S}(\mathbb{R}^d)$. Recall the estimate of the proof of Proposition 3.1.7.

$$|\varphi(x)| \leq \frac{A_d}{(1 + |x|^2)^{\frac{d+1}{2}}} N_{d+1}(\varphi).$$

Consequently, for every $\gamma \in \mathbb{Z}^d$, we have

$$\sup_{x \in \gamma + Q} |\varphi(x)| \leq \frac{B_d}{(1 + |\gamma|^2)^{\frac{d+1}{2}}} N_{d+1}(\varphi).$$

Using a comparison between the sum over \mathbb{Z}^d and the integral over \mathbb{R}^d , one can easily show that

$$\sum_{\gamma \in \mathbb{Z}^d} \frac{1}{(1 + |\gamma|^2)^{\frac{d+1}{2}}} < \infty.$$

Hence the series defining $\varphi^\sharp(x)$ is uniformly convergent on compact subset of \mathbb{R}^d , and this also true for the series defining $(\partial^\alpha \varphi)^\sharp$. Consequently, φ^\sharp is smooth, $\partial^\alpha \varphi^\sharp = (\partial^\alpha \varphi)^\sharp$ and (3.6.10) follows. Finally, choose $\chi_1 \in \mathcal{C}_0^\infty(\mathbb{R}^d)$ such that $0 \leq \chi_1(x) \leq 1$ and $\chi_1(x) = 1$ on Q . Then $\chi^\sharp \in \mathcal{C}^\infty(\mathbb{R}^d)$ is \mathbb{Z}^d -periodic, and $\chi_1^\sharp \geq 1$. Therefore

$$\chi := \frac{\chi_1}{\chi_1^\sharp} \in \mathcal{C}_0^\infty(\mathbb{R}^d)$$

satisfies $\chi^\sharp = 1$. □

Given $k \in (2\pi\mathbb{Z})^d$, notice that

$$e_k(x) := e^{ik \cdot x}$$

defines a $\mathcal{C}^\infty \mathbb{Z}^d$ -periodic function. We say that a sequence $(c_k)_{k \in (2\pi\mathbb{Z})^d}$ of complex numbers has moderate growth if there exists $M, C > 0$ such that

$$\forall k \in (2\pi\mathbb{Z})^d, |c_k| \leq C(1 + |k|)^M.$$

If (c_k) is such a sequence, the series

$$\sum_{k \in (2\pi\mathbb{Z})^d} c_k e_k$$

defines a tempered distribution. Indeed, if $\varphi \in \mathcal{S}(\mathbb{R}^d)$, $\hat{\varphi}(k) = O((1 + |k|)^{-p}) N_{p+d+1}(\varphi)$ for every p , so the series

$$\sum_{k \in (2\pi\mathbb{Z})^d} c_k \hat{\varphi}(-k)$$

is absolutely summable, and we can define a tempered distribution by

$$\forall \varphi \in \mathcal{S}(\mathbb{R}^d), \left\langle \sum_{k \in (2\pi\mathbb{Z})^d} c_k e_k, \varphi \right\rangle = \sum_{k \in (2\pi\mathbb{Z})^d} c_k \hat{\varphi}(-k).$$

The next theorem claims that every \mathbb{Z}^d -periodic distribution is of this form.

Theorem 3.6.3 Every \mathbb{Z}^d -periodic distribution T can be written uniquely

$$(3.6.11) \quad T = \sum_{k \in (2\pi\mathbb{Z})^d} c_k(T) e_k, \quad e_k(x) = e^{ik \cdot x},$$

where $(c_k(T))_{k \in (2\pi\mathbb{Z})^d}$ is a sequence of complex numbers with moderate growth. Furthermore, for every $\chi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$ such that $\chi^\sharp = 1$, we have

$$\forall k \in (2\pi\mathbb{Z})^d, \quad c_k(T) = \langle T, \chi e_{-k} \rangle.$$

Proof.— First, we prove that every periodic distribution T is tempered. Indeed, if $\varphi, \psi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$, let us check the formula

$$(3.6.12) \quad \langle T, \varphi \psi^\sharp \rangle = \langle T, \varphi^\sharp \psi \rangle.$$

Since the series defining φ^\sharp converges in \mathcal{C}^∞ , we have

$$\langle T, \varphi \psi^\sharp \rangle = \sum_{\gamma \in \mathbb{Z}^d} \langle T, \varphi \tau_\gamma \psi \rangle = \sum_{\gamma \in \mathbb{Z}^d} \langle T, \tau_{-\gamma} \varphi \psi \rangle = \langle T, \varphi^\sharp \psi \rangle.$$

Then, introducing $\chi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$ such that $\chi^\sharp = 1$, we have, for every $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$,

$$\langle T, \varphi \rangle = \langle T, \varphi \chi^\sharp \rangle = \langle T, \varphi^\sharp \chi \rangle.$$

Since T is a distribution, we infer, for some $C > 0$ and some integer m ,

$$|\langle T, \varphi \rangle| \leq C \sup_{|\alpha| \leq m} \|\partial^\alpha (\chi \varphi^\sharp)\|_{L^\infty} \leq \tilde{C} N_{d+1+m}(\varphi),$$

using the Leibniz formula and estimate (3.6.10). Hence $T \in \mathcal{S}'(\mathbb{R}^d)$.

The next step is to characterize the structure of $\hat{T} \in \mathcal{S}'(\mathbb{R}^d)$. Using the identity

$$\widehat{\tau_a T} = e^{-ia \cdot \xi} \hat{T},$$

and the \mathbb{Z}^d -periodicity of T , we have

$$\forall \gamma \in \mathbb{Z}^d, \quad e^{-i\gamma \cdot \xi} \hat{T} = \hat{T}.$$

Notice that the set where all the functions $\xi \mapsto (e^{-i\gamma \cdot \xi} - 1)$ vanish is $(2\pi\mathbb{Z})^d$, hence the support of \hat{T} is included into $(2\pi\mathbb{Z})^d$. Let us describe in more detail the structure of \hat{T} near each point $k \in (2\pi\mathbb{Z})^d$. Set

$$Q_k^* := k +] - 2\pi, 2\pi[^d.$$

Then, for $j = 1, \dots, d$, the functions

$$\xi \mapsto \frac{e^{i\xi_j} - 1}{\xi_j - k_j}$$

are C^∞ and do not cancel on Q_k^* . Consequently, on Q_k^* , the equations

$$e^{-i\xi_j} \hat{T} = \hat{T}, \quad j = 1, \dots, d$$

are equivalent to

$$(x_j - k_j) \hat{T} = 0, \quad j = 1, \dots, d,$$

and we know that this implies, by Proposition 2.3.11, that there exists $a_k \in \mathbb{C}$ such that

$$\hat{T}|_{Q_k^*} = a_k \delta_k.$$

Since the open sets Q_k^* cover \mathbb{R}^d , we infer, by the gluing principle, that

$$\hat{T} = \sum_{k \in (2\pi\mathbb{Z})^d} a_k \delta_k.$$

Furthermore, notice that the sequence (a_k) has moderate growth. Indeed, since \hat{T} is tempered, we have, if χ_0 is a Plateau function above 0 and compactly supported in $] - 2\pi, 2\pi[^d$,

$$|a_k| = |\langle T, \tau_k \chi_0 \rangle| \leq CN_p(\tau_k \chi_0) \leq B(1 + |k|)^p.$$

Finally, applying the inverse Fourier transform, we obtain

$$T = (2\pi)^d \sum_{k \in (2\pi\mathbb{Z})^d} a_k \check{\mathcal{F}}(\delta_k) = (2\pi)^d \sum_{k \in (2\pi\mathbb{Z})^d} a_k e_k.$$

This proves the first part of the theorem, with $c_k := (2\pi)^d a_k$. Furthermore, given $\ell \in (2\pi\mathbb{Z})^d$, and χ a test function such that $\chi^\# = 1$,

$$\langle T, \chi e_{-\ell} \rangle = \sum_{k \in (2\pi\mathbb{Z})^d} c_k \langle e_k, \chi e_{-\ell} \rangle = \sum_{k \in (2\pi\mathbb{Z})^d} c_k \langle 1, \chi e_{k-\ell} \rangle.$$

But, by formula (3.6.9),

$$\langle 1, \chi e_{k-\ell} \rangle = \int_{\mathbb{R}^d} \chi e_{k-\ell} = \int_Q \chi^\# e_{k-\ell} = \int_Q e_{k-\ell} = \begin{cases} 0 & \text{if } k \neq \ell \\ 1 & \text{if } k = \ell. \end{cases}$$

Finally, we obtain

$$\langle T, \chi e_{-\ell} \rangle = c_\ell,$$

as claimed, and this proves the uniqueness of the sequence (c_k) . \square

Remark 3.6.4 By deriving the Fourier decomposition (3.6.11), we have

$$c_k(\partial^\alpha T) = (ik)^\alpha c_k(T), \quad k \in (2\pi\mathbb{Z})^d.$$

Corollary 3.6.5 Let f be a locally integrable \mathbb{Z}^d -periodic function on \mathbb{R}^d . Then, in $\mathcal{S}'(\mathbb{R}^d)$,

$$(3.6.13) \quad f = \sum_{k \in (2\pi\mathbb{Z})^d} c_k(f) e^{ik \cdot x}, \quad c_k(f) := \int_Q f(x) e^{-ik \cdot x} dx, \quad k \in (2\pi\mathbb{Z})^d.$$

Furthermore, if f is locally square integrable, the above series is convergent in $L^2(Q)$, with

$$\|f\|_{L^2(Q)}^2 = \sum_{k \in (2\pi\mathbb{Z})^d} |c_k(f)|^2.$$

Finally, if f is smooth, the sequence $(c_k(f))$ is rapidly decreasing and the above series is convergent in \mathcal{C}^∞ .

Proof.— Let χ be a test function such that $\chi^\sharp = 1$. Then

$$c_k(f) = \langle f, \chi e_{-k} \rangle = \int_{\mathbb{R}^d} f(x) \chi(x) e^{-ik \cdot x} dx = \int_Q (f \chi e_{-k})^\sharp dx = \int_Q f e_{-k} dx.$$

Therefore the claim is a consequence of Theorem 3.6.3.

In the L^2 case, we observe that (e_k) is a orthonormal family of $L^2(Q)$ and that $c_k(f)$ is the inner product of f with e_k . From the identity (3.6.13), we observe that, if $c_k(f) = 0$ for every k , then $f = 0$. This implies that (e_k) is an orthonormal basis of $L^2(Q)$, and the assertion follows.

Finally, if f is smooth, for every α $\partial^\alpha f$ is locally square integrable, thus

$$\sum_k |k^\alpha c_k(f)|^2 = \sum_k |c_k(\partial^\alpha f)|^2 < \infty,$$

therefore $(c_k(f))$ is rapidly decreasing and consequently the series and all its derivatives converge uniformly.

□

Corollary 3.6.6 (The Poisson summation formula)

$$\sum_{\gamma \in \mathbb{Z}^d} \delta_\gamma = \sum_{k \in (2\pi\mathbb{Z})^d} e^{ik \cdot x}.$$

Proof.— Let χ be a test function such that $\chi^\sharp = 1$. Then

$$c_k \left(\sum_{\gamma \in \mathbb{Z}^d} \delta_\gamma \right) = \sum_{\gamma \in \mathbb{Z}^d} \langle \delta_\gamma, \chi e_{-k} \rangle = \sum_{\gamma \in \mathbb{Z}^d} \chi(\gamma) = \chi^\sharp(0) = 1.$$

Therefore the claim is a consequence of Theorem 3.6.3. □

As an application, let us now consider periodic solutions of PDE with constant coefficients.

Corollary 3.6.7 Let $p \in \mathbb{C}[X_1, \dots, X_d]$ and $P = p(\partial)$ be the corresponding differential operator.

- i) If $p(ik) \neq 0$ for every $k \in 2\pi\mathbb{Z}^d$, then the only \mathbb{Z}^d periodic solution u of $Pu = 0$ is $u = 0$.
- ii) If $p(ik) \neq 0$ for every $k \in 2\pi\mathbb{Z}^d \setminus \{0\}$, then every \mathbb{Z}^d -periodic solution of $Pu = 0$ is a constant.

As an illustration of Corollary 3.6.7, let us prove the following density theorem. Let $\omega = (\omega_1, \dots, \omega_d) \in \mathbb{R}^d$ be such that the components are linearly independent on \mathbb{Q} . Then, for every $x \in \mathbb{R}^d$, for every $\varepsilon > 0$, there exist $t \in \mathbb{R}$ and $\gamma \in \mathbb{Z}^d$ such that

$$|x - t\omega - \gamma| < \varepsilon.$$

Indeed, the function u defined on \mathbb{R}^d by

$$u(x) := \inf_{t \in \mathbb{R}} \inf_{\gamma \in \mathbb{Z}^d} |x - t\omega - \gamma|$$

is Lipschitz continuous, and is \mathbb{Z}^d periodic. Furthermore, for every $t \in \mathbb{R}$,

$$u(x + t\omega) = u(x).$$

This means

$$\forall \varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d), \quad \int_{\mathbb{R}^d} u(x + t\omega)\varphi(x) dx = \int_{\mathbb{R}^d} u(x)\varphi(x) dx.$$

Notice that the left hand side reads

$$\int_{\mathbb{R}^d} u(x + t\omega)\varphi(x) dx = \int_{\mathbb{R}^d} u(y)\varphi(y - t\omega) dy,$$

and, taking the derivative of this quantity with respect to t at $t = 0$, we obtain

$$- \int_{\mathbb{R}^d} u(y) \sum_{j=1}^d \omega_j \partial_j \varphi(y) dy.$$

In other words, in the sense of distributions on \mathbb{R}^d ,

$$\sum_{j=1}^d \omega_j \partial_j u = 0.$$

Since, by the assumption on the components of ω , the polynomial function

$$p(x) = \sum_{j=1}^d \omega_j x_j$$

does not vanish at $k \in 2\pi\mathbb{Z}^d \setminus \{0\}$, we infer that u is constant. Since clearly $u(0) = 0$, this completes the proof.

3.7 An application to probability theory

Recall that a probability measure on \mathbb{R}^d is a positive Borel measure μ on \mathbb{R}^d of total mass $\mu(\mathbb{R}^d) = 1$. Such a measure defines a tempered distribution through the formula

$$\forall \varphi \in \mathcal{S}(\mathbb{R}^d), \langle \mu, \varphi \rangle = \int_{\mathbb{R}^d} \varphi(x) d\mu(x).$$

Using Fubini's theorem, we observe that

$$\langle \hat{\mu}, \varphi \rangle = \int_{\mathbb{R}^d} \hat{\mu}(\xi) \varphi(\xi) d\xi,$$

where

$$\hat{\mu}(\xi) = \int_{\mathbb{R}^d} e^{-ix \cdot \xi} d\mu(x).$$

Notice that $|\hat{\mu}(\xi)| \leq 1$ and that the function $\hat{\mu}$ is continuous, thanks to the dominated convergence theorem. We now introduce a standard definition in probability theory.

Definition 3.7.1 We say that a sequence (μ_n) of probability measures converges tightly to a probability measure μ if, for every bounded continuous function f on \mathbb{R}^d ,

$$\int_{\mathbb{R}^d} f d\mu_n \rightarrow \int_{\mathbb{R}^d} f d\mu.$$

Before stating the main result, let us make the connection between tight convergence and convergence in \mathcal{S}' .

Proposition 3.7.2 Let (μ_n) be a sequence of probability measures on \mathbb{R}^d , and let μ be a probability measure on \mathbb{R}^d . Then μ_n converges tightly to μ if and only if μ_n converges to μ in $\mathcal{S}'(\mathbb{R}^d)$.

Proof.— Since every function in the Schwartz class is bounded and continuous, the only non trivial part to be proven is that the convergence in \mathcal{S}' of the probability measures μ_n to the probability measure μ implies the tight convergence of μ_n to μ . First of all, we prove that

$$\int_{\mathbb{R}^d} f d\mu_n \rightarrow \int_{\mathbb{R}^d} f d\mu,$$

if f is continuous and compactly supported. Indeed, by regularization, f can be uniformly approximated by a sequence of elements of the Schwartz space. The uniform bound on $\mu_n(\mathbb{R}^d) = 1$ then allows to extend the convergence on Schwartz test functions to continuous compactly supported test functions. Next, let f be bounded continuous function. Let χ be a plateau function over the unit ball of \mathbb{R}^d , and define, for every $R > 0$,

$$\chi_R(x) = \chi\left(\frac{x}{R}\right).$$

Then

$$\int_{\mathbb{R}^d} f d\mu_n - \int_{\mathbb{R}^d} f d\mu = \int_{\mathbb{R}^d} f \chi_R d\mu_n - \int_{\mathbb{R}^d} f \chi_R d\mu + \int_{\mathbb{R}^d} f(1 - \chi_R) d\mu_n - \int_{\mathbb{R}^d} f(1 - \chi_R) d\mu.$$

Since $f\chi_R$ is continuous and compactly supported, we have

$$\int_{\mathbb{R}^d} f\chi_R d\mu_n - \int_{\mathbb{R}^d} f\chi_R d\mu \xrightarrow{n \rightarrow \infty} 0.$$

Furthermore,

$$\left| \int_{\mathbb{R}^d} f(1 - \chi_R) d\mu_n \right| \leq \|f\|_{L^\infty} \int (1 - \chi_R) d\mu_n = \|f\|_{L^\infty} \left(1 - \int \chi_R d\mu_n \right) \xrightarrow{n \rightarrow \infty} \|f\|_{L^\infty} \left(1 - \int \chi_R d\mu \right).$$

Summing up, we have, for every $R > 0$,

$$\limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^d} f d\mu_n - \int_{\mathbb{R}^d} f d\mu \right| \leq 2\|f\|_{L^\infty} \left(1 - \int \chi_R d\mu \right).$$

Since μ is a probability measure, the right hand side tends to 0 as R tends to ∞ , by the dominated convergence theorem. Hence the left hand side, which is independent of R , is 0. \square

We now state the main result of this paragraph.

Theorem 3.7.3 (P. Lévy) Let (μ_n) be a sequence of probability measures on \mathbb{R}^d . The following statements are equivalent.

- i) μ_n converges tightly to some probability measure.
- ii) There exists a function $g : \mathbb{R}^d \rightarrow \mathbb{C}$, continuous at 0, such that,

$$\forall \xi \in \mathbb{R}^d, \hat{\mu}_n(\xi) \xrightarrow{n \rightarrow \infty} g(\xi).$$

Proof.— The first implication is trivial. Indeed, since, for every $\xi \in \mathbb{R}^d$, the function $f : x \mapsto e^{-ix \cdot \xi}$ is continuous and bounded, the tight convergence of μ_n to μ implies the convergence of $\hat{\mu}_n(\xi)$ to $\hat{\mu}(\xi)$.

Let us prove that (ii) implies (i). Since $\hat{\mu}_n$ and continuous and bounded by 1, we already know that g is measurable and bounded by 1, hence it defines a tempered distribution. Furthermore, the dominated convergence theorem implies that $\hat{\mu}_n \rightarrow g$ in \mathcal{S}' . Introducing $T = \mathcal{F}^{-1}(g)$, we infer that μ_n converges to T in \mathcal{S}' . Since μ_n is a positive measure, it is a positive distribution, and so is T . Hence $T = \mu$ is a positive measure.

It remains to prove that μ is a probability measure. Let χ be a plateau function over the unit ball, and, for every $R > 0$,

$$\chi_R(x) = \chi\left(\frac{x}{R}\right).$$

By Fatou's lemma,

$$\mu(\mathbb{R}^d) \leq \liminf_{R \rightarrow \infty} \int_{\mathbb{R}^d} \chi_R d\mu.$$

Since

$$\int_{\mathbb{R}^d} \chi_R d\mu = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} \chi_R d\mu_n \leq 1,$$

we first conclude that $\mu(\mathbb{R}^d) \leq 1$. Hence μ has finite total mass, and the dominated convergence theorem implies that

$$\mu(\mathbb{R}^d) = \lim_{R \rightarrow \infty} \int_{\mathbb{R}^d} \chi_R d\mu .$$

Let $\rho \in \mathcal{S}$ such that $\chi = \hat{\rho}$. Then $\chi_R = \hat{\rho}_R$, with

$$\rho_R(\xi) = R^d \rho(R\xi) .$$

Hence

$$\begin{aligned} \int_{\mathbb{R}^d} \chi_R d\mu &= \int_{\mathbb{R}^d} \hat{\rho}_R d\mu \\ &= \langle \mu, \hat{\rho}_R \rangle = \langle \hat{\mu}, \rho_R \rangle = \int_{\mathbb{R}^d} g(\xi) \rho_R(\xi) d\xi \\ &= \int_{\mathbb{R}^d} g\left(\frac{\xi}{R}\right) \rho(\xi) d\xi . \end{aligned}$$

Since g is continuous at $\xi = 0$, we infer, from the dominated convergence theorem,

$$\int_{\mathbb{R}^d} g\left(\frac{\xi}{R}\right) \rho(\xi) d\xi \xrightarrow{R \rightarrow \infty} g(0) \int_{\mathbb{R}^d} \rho(\xi) d\xi = g(0) \chi(0) = g(0) .$$

But $g(0) = \lim_{n \rightarrow \infty} \hat{\mu}_n(0) = 1$. Hence $\mu(\mathbb{R}^d) = 1$. □

Remark 3.7.4 Notice that the assumption of continuity of g at 0 is crucial, as shown by the following counterexample. Let $\rho \in L^1(\mathbb{R}^d)$ be nonnegative, with integral 1 on \mathbb{R}^d . For $n \geq 1$, consider the probability measure μ_n given by

$$d\mu_n = \rho_n(x) dx , \quad \rho_n(x) := \frac{1}{n^d} \rho\left(\frac{x}{n}\right) .$$

Then, by the Riemann–Lebesgue lemma,

$$\hat{\mu}_n(\xi) = \hat{\rho}(n\xi) \xrightarrow{n \rightarrow \infty} g(\xi) = \begin{cases} 0 & \text{if } \xi \neq 0 \\ 1 & \text{if } \xi = 0 . \end{cases}$$

Thus g is not continuous at 0. Indeed, one easily checks that μ_n converges to 0 in \mathcal{S}' , hence does not converge tightly.

We conclude by the classical application of P. Lévy's theorem to the central limit theorem.

Theorem 3.7.5 Let μ be a probability measure on \mathbb{R}^d such that

$$\int_{\mathbb{R}^d} |x|^2 d\mu(x) < \infty , \quad m = \int_{\mathbb{R}^d} x d\mu(x) .$$

Assume that no affine hyperplane H of \mathbb{R}^d passing through m satisfies $\mu(H) = 1$. Then the symmetric matrix $A = (a_{jk})_{1 \leq j, k \leq d}$ defined by

$$a_{jk} = \int_{\mathbb{R}^d} (x_j - m_j)(x_k - m_k) d\mu(x)$$

is positive definite and, for every bounded continuous function f on \mathbb{R}^d ,

$$\int_{(\mathbb{R}^d)^n} f\left(\frac{x_1 + \cdots + x_n - nm}{\sqrt{n}}\right) d\mu(x_1) \cdots d\mu(x_n) \xrightarrow{n \rightarrow \infty} \int_{\mathbb{R}^d} \frac{e^{-A^{-1}x \cdot x/2}}{(2\pi)^{d/2} \sqrt{\det A}} f(x) dx .$$

Proof.— For every integer $n \geq 1$, the formula

$$\int_{\mathbb{R}^d} f d\mu_n = \int_{(\mathbb{R}^d)^n} f\left(\frac{x_1 + \cdots + x_n - nm}{\sqrt{n}}\right) d\mu(x_1) \cdots d\mu(x_n)$$

defines a probability measure on \mathbb{R}^d , and we have to prove that μ_n converges tightly to the probability measure of density

$$G_A(x) = \frac{e^{-A^{-1}x \cdot x/2}}{(2\pi)^{d/2} \sqrt{\det A}}$$

with respect to the Lebesgue measure. Notice that, after diagonalising A and using the Fubini theorem, a linear change of variables and Proposition 3.2.3,

$$\widehat{G}_A(\xi) = e^{-A\xi \cdot \xi/2} .$$

Therefore, by P. Lévy's theorem, it is enough to prove that

$$\forall \xi \in \mathbb{R}^d, \hat{\mu}_n(\xi) \xrightarrow{n \rightarrow \infty} e^{-A\xi \cdot \xi/2} .$$

From the above definition of μ_n and the Fubini theorem, we have

$$\hat{\mu}_n(\xi) = \left[e^{i\xi \cdot m/\sqrt{n}} \hat{\mu}\left(\frac{\xi}{\sqrt{n}}\right) \right]^n .$$

Because of the assumption

$$\int_{\mathbb{R}^d} |x|^2 d\mu(x) < \infty ,$$

the function

$$\hat{\mu}(\xi) = \int_{\mathbb{R}^d} e^{-ix \cdot \xi} d\mu(x)$$

is C^2 and we can expand it near 0 by the Taylor formula, so that

$$\begin{aligned} e^{i\xi \cdot m/\sqrt{n}} \hat{\mu}\left(\frac{\xi}{\sqrt{n}}\right) &= \left(1 + i\frac{\xi \cdot m}{\sqrt{n}} - \frac{(\xi \cdot m)^2}{2n} + o\left(\frac{1}{n}\right)\right) \left(1 - i\frac{\xi \cdot m}{\sqrt{n}} - \frac{1}{2n} \int_{\mathbb{R}^d} (\xi \cdot x)^2 d\mu(x) + o\left(\frac{1}{n}\right)\right) \\ &= 1 - \frac{A\xi \cdot \xi}{2n} + o\left(\frac{1}{n}\right) . \end{aligned}$$

Consequently,

$$\hat{\mu}_n(\xi) = \left(1 - \frac{A\xi \cdot \xi}{2n} + o\left(\frac{1}{n}\right)\right)^n ,$$

which — by taking for instance the complex logarithm near 1 — converges to $\exp(-A\xi \cdot \xi/2)$ as $n \rightarrow \infty$. \square