

# Adiabatic theorems and linear response in the thermodynamic limit

Stefan Teufel

Fachbereich Mathematik, Universität Tübingen

Problèmes Spectraux en Physique Mathématique

Paris, January 18, 2021

## Motivation and plan of the talk

**Barry Simon:** Fifteen problems in mathematical physics (1984)

**4. Transport Theory:** *At some level, the fundamental difficulty of transport theory is that it is a steady state rather than equilibrium problem, so that the powerful formalism of equilibrium statistical mechanics is unavailable, and one does not have any way of precisely identifying the steady state and thereby computing things in it.*

⋮

Problem 4 B (Kubo Formula) *Either justify Kubo's formula in a quantum model, or else find an alternate theory of conductivity.*

# Motivation and plan of the talk

**Barry Simon:** Fifteen problems in mathematical physics (1984)

**4. Transport Theory:** *At some level, the fundamental difficulty of transport theory is that it is a steady state rather than equilibrium problem, so that the powerful formalism of equilibrium statistical mechanics is unavailable, and one does not have any way of precisely identifying the steady state and thereby computing things in it.*

⋮

Problem 4 B (Kubo Formula) *Either justify Kubo's formula in a quantum model, or else find an alternate theory of conductivity.*

**Plan of the talk:**

1. Linear response at zero temperature: setup and ideas

# Motivation and plan of the talk

**Barry Simon:** Fifteen problems in mathematical physics (1984)

**4. Transport Theory:** *At some level, the fundamental difficulty of transport theory is that it is a steady state rather than equilibrium problem, so that the powerful formalism of equilibrium statistical mechanics is unavailable, and one does not have any way of precisely identifying the steady state and thereby computing things in it.*

⋮

Problem 4 B (Kubo Formula) *Either justify Kubo's formula in a quantum model, or else find an alternate theory of conductivity.*

## Plan of the talk:

1. Linear response at zero temperature: setup and ideas
2. Adiabatic theorem for fermions in the thermodynamic limit with a gap in the bulk

## Linear response and Kubo's formula

In the context of **Hamiltonian quantum systems**, the linear response formalism answers the following question:

How does a system described by a Hamiltonian  $H_0$  that is initially in an equilibrium state  $\rho_0$  respond to a small static perturbation  $\varepsilon V$ ?

## Linear response and Kubo's formula

In the context of **Hamiltonian quantum systems**, the linear response formalism answers the following question:

How does a system described by a Hamiltonian  $H_0$  that is initially in an equilibrium state  $\rho_0$  respond to a small static perturbation  $\varepsilon V$ ?

Or somewhat more precisely:

What is the change

$$\text{tr}(\rho_\varepsilon A) - \text{tr}(\rho_0 A) = \varepsilon \cdot \sigma_A + o(\varepsilon)$$

of the expectation value of an observable  $A$  caused by the perturbation  $\varepsilon V$  at leading order in its strength  $\varepsilon$  ?

Here  $\rho_\varepsilon$  denotes the state of the system after the perturbation has been turned on and  $\sigma_A$  is called the linear response coefficient for  $A$ .

## Linear response and Kubo's formula

In the context of **Hamiltonian quantum systems**, the linear response formalism answers the following question:

How does a system described by a Hamiltonian  $H_0$  that is initially in an equilibrium state  $\rho_0$  respond to a small static perturbation  $\varepsilon V$ ?

Or somewhat more precisely:

What is the change

$$\text{tr}(\rho_\varepsilon A) - \text{tr}(\rho_0 A) = \varepsilon \cdot \sigma_A + o(\varepsilon)$$

of the expectation value of an observable  $A$  caused by the perturbation  $\varepsilon V$  at leading order in its strength  $\varepsilon$  ?

Here  $\rho_\varepsilon$  denotes the state of the system after the perturbation has been turned on and  $\sigma_A$  is called the linear response coefficient for  $A$ .

The answer clearly hinges on the problem of determining  $\rho_\varepsilon$ .

# Linear response and Kubo's formula

Modelling the switching process: Let

$$H_\varepsilon(t) := H_0 + f(t) \varepsilon V$$

with a switch function  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(t) = 0$  for  $t \leq -1$  and  $f(t) = 1$  for  $t \geq 0$ .

# Linear response and Kubo's formula

Modelling the switching process: Let

$$H_\varepsilon(t) := H_0 + f(t) \varepsilon V$$

with a switch function  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(t) = 0$  for  $t \leq -1$  and  $f(t) = 1$  for  $t \geq 0$ .

Let  $\rho(t)$  be the solution of the time-dependent Schrödinger equation

$$i \frac{d}{dt} \rho(t) = [H^\varepsilon(\eta t), \rho(t)]$$

with  $\rho(t) = \rho_0$  for  $t \leq -1/\eta$  and adiabatic parameter  $\eta \ll 1$ .

# Linear response and Kubo's formula

Modelling the switching process: Let

$$H_\varepsilon(t) := H_0 + f(t) \varepsilon V$$

with a switch function  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(t) = 0$  for  $t \leq -1$  and  $f(t) = 1$  for  $t \geq 0$ .

Let  $\rho(t)$  be the solution of the time-dependent Schrödinger equation

$$i \frac{d}{dt} \rho(t) = [H^\varepsilon(\eta t), \rho(t)]$$

with  $\rho(t) = \rho_0$  for  $t \leq -1/\eta$  and adiabatic parameter  $\eta \ll 1$ .

Then  $\rho(0)$ , or actually  $\rho(t)$  for any  $t \geq 0$ , is the natural candidate for the “state of the system after the perturbation has been turned on”.

## Linear response and Kubo's formula

Approximating  $\rho_\varepsilon := \rho(0)$  by first order time-dependent perturbation theory

$$\rho_\varepsilon = \rho_0 - \varepsilon i \int_{-\infty}^0 f(\eta t) e^{iH_0 t} [V, \rho_0] e^{-iH_0 t} dt + R^{\varepsilon, \eta, f},$$

one obtains

$$\text{tr}(\rho_\varepsilon A) - \text{tr}(\rho_0 A) = \varepsilon \tilde{\sigma}_A^{\eta, f} + \text{tr}(R^{\varepsilon, \eta, f} A)$$

with

$$\tilde{\sigma}_A^{\eta, f} = i \int_{-\infty}^0 f(\eta t) \left\langle [V, e^{-iH_0 t} A e^{iH_0 t}] \right\rangle_{\rho_0} dt.$$

## Linear response and Kubo's formula

Approximating  $\rho_\varepsilon := \rho(0)$  by first order time-dependent perturbation theory

$$\rho_\varepsilon = \rho_0 - \varepsilon i \int_{-\infty}^0 f(\eta t) e^{iH_0 t} [V, \rho_0] e^{-iH_0 t} dt + R^{\varepsilon, \eta, f},$$

one obtains

$$\text{tr}(\rho_\varepsilon A) - \text{tr}(\rho_0 A) = \varepsilon \tilde{\sigma}_A^{\eta, f} + \text{tr}(R^{\varepsilon, \eta, f} A)$$

with

$$\tilde{\sigma}_A^{\eta, f} = i \int_{-\infty}^0 f(\eta t) \left\langle [V, e^{-iH_0 t} A e^{iH_0 t}] \right\rangle_{\rho_0} dt.$$

One hopes to obtain a universal result independent of  $\eta$  and  $f$  in the adiabatic limit  $\eta \rightarrow 0$ .

## Linear response and Kubo's formula

Approximating  $\rho_\varepsilon := \rho(0)$  by first order time-dependent perturbation theory

$$\rho_\varepsilon = \rho_0 - \varepsilon i \int_{-\infty}^0 f(\eta t) e^{iH_0 t} [V, \rho_0] e^{-iH_0 t} dt + R^{\varepsilon, \eta, f},$$

one obtains

$$\text{tr}(\rho_\varepsilon A) - \text{tr}(\rho_0 A) = \varepsilon \tilde{\sigma}_A^{\eta, f} + \text{tr}(R^{\varepsilon, \eta, f} A)$$

with

$$\tilde{\sigma}_A^{\eta, f} = i \int_{-\infty}^0 f(\eta t) \left\langle [V, e^{-iH_0 t} A e^{iH_0 t}] \right\rangle_{\rho_0} dt.$$

One hopes to obtain a universal result independent of  $\eta$  and  $f$  in the adiabatic limit  $\eta \rightarrow 0$ . Choosing  $f = \exp$  this results in “Kubo's formula” for the linear response coefficient

$$\sigma_A^{\text{Kubo}} := \lim_{\eta \rightarrow 0} \tilde{\sigma}_A^{\eta, \exp} = \lim_{\eta \rightarrow 0} i \int_{-\infty}^0 e^{\eta t} \left\langle [V, e^{-iH_0 t} A e^{iH_0 t}] \right\rangle_{\rho_0} dt.$$

# Linear response and Kubo's formula

⋮

one obtains

$$\mathrm{tr}(\rho_\varepsilon A) - \mathrm{tr}(\rho_0 A) = \varepsilon \tilde{\sigma}_A^{\eta, f} + \mathrm{tr}(R^{\varepsilon, \eta, f} A)$$

One hopes to obtain a universal result independent of  $\eta$  and  $f$  in the adiabatic limit  $\eta \rightarrow 0$ . Choosing  $f = \exp$  this results in “Kubo's formula” for the linear response coefficient

$$\sigma_A^{\text{Kubo}} := \lim_{\eta \rightarrow 0} \tilde{\sigma}_A^{\eta, \exp} = \lim_{\eta \rightarrow 0} i \int_{-\infty}^0 e^{\eta t} \left\langle \left[ V, e^{-iH_0 t} A e^{iH_0 t} \right] \right\rangle_{\rho_0} dt.$$

# Linear response and Kubo's formula

⋮

one obtains

$$\mathrm{tr}(\rho_\varepsilon A) - \mathrm{tr}(\rho_0 A) = \varepsilon \tilde{\sigma}_A^{\eta, f} + \mathrm{tr}(R^{\varepsilon, \eta, f} A)$$

One hopes to obtain a universal result independent of  $\eta$  and  $f$  in the adiabatic limit  $\eta \rightarrow 0$ . Choosing  $f = \exp$  this results in “Kubo's formula” for the linear response coefficient

$$\sigma_A^{\mathrm{Kubo}} := \lim_{\eta \rightarrow 0} \tilde{\sigma}_A^{\eta, \exp} = \lim_{\eta \rightarrow 0} i \int_{-\infty}^0 e^{\eta t} \left\langle \left[ V, e^{-iH_0 t} A e^{iH_0 t} \right] \right\rangle_{\rho_0} dt.$$

“Justifying Kubo's formula” can mean two different things now:

- ▶ Show existence of the limit and compute  $\sigma_A^{\mathrm{Kubo}} = \lim_{\eta \rightarrow 0} \tilde{\sigma}_A^{\eta, \exp}$ .

# Linear response and Kubo's formula

⋮

one obtains

$$\mathrm{tr}(\rho_\varepsilon A) - \mathrm{tr}(\rho_0 A) = \varepsilon \tilde{\sigma}_A^{\eta, f} + \mathrm{tr}(R^{\varepsilon, \eta, f} A)$$

One hopes to obtain a universal result independent of  $\eta$  and  $f$  in the adiabatic limit  $\eta \rightarrow 0$ . Choosing  $f = \exp$  this results in “Kubo's formula” for the linear response coefficient

$$\sigma_A^{\mathrm{Kubo}} := \lim_{\eta \rightarrow 0} \tilde{\sigma}_A^{\eta, \exp} = \lim_{\eta \rightarrow 0} i \int_{-\infty}^0 e^{\eta t} \left\langle \left[ V, e^{-iH_0 t} A e^{iH_0 t} \right] \right\rangle_{\rho_0} dt.$$

“Justifying Kubo's formula” can mean two different things now:

- ▶ Show existence of the limit and compute  $\sigma_A^{\mathrm{Kubo}} = \lim_{\eta \rightarrow 0} \tilde{\sigma}_A^{\eta, \exp}$ .  
(e.g. **Bouclet, Germinet, Klein, Schenker JFA '05**)

# Linear response and Kubo's formula

⋮

one obtains

$$\mathrm{tr}(\rho_\varepsilon A) - \mathrm{tr}(\rho_0 A) = \varepsilon \tilde{\sigma}_A^{\eta, f} + \mathrm{tr}(R^{\varepsilon, \eta, f} A)$$

One hopes to obtain a universal result independent of  $\eta$  and  $f$  in the adiabatic limit  $\eta \rightarrow 0$ . Choosing  $f = \exp$  this results in “Kubo's formula” for the linear response coefficient

$$\sigma_A^{\mathrm{Kubo}} := \lim_{\eta \rightarrow 0} \tilde{\sigma}_A^{\eta, \exp} = \lim_{\eta \rightarrow 0} i \int_{-\infty}^0 e^{\eta t} \left\langle \left[ V, e^{-iH_0 t} A e^{iH_0 t} \right] \right\rangle_{\rho_0} dt.$$

“Justifying Kubo's formula” can mean two different things now:

- ▶ Show existence of the limit and compute  $\sigma_A^{\mathrm{Kubo}} = \lim_{\eta \rightarrow 0} \tilde{\sigma}_A^{\eta, \exp}$ .
- ▶ Show that  $\mathrm{tr}(R^{\varepsilon, \eta, f} A) = o(\varepsilon)$  uniformly in  $\eta$  and that  $\sigma_A^{\mathrm{Kubo}} = \lim_{\eta \rightarrow 0} \tilde{\sigma}_A^{\eta, f}$  for any switching function  $f$ .

# Linear response and Kubo's formula

Why should this work?

## Linear response and Kubo's formula

Why should this work? Two possible mechanisms:

- (a) During the switching process the state of the system remains (approximately) in an **instantaneous equilibrium state** of the perturbed Hamiltonian.

## Linear response and Kubo's formula

Why should this work? Two possible mechanisms:

- (a) During the switching process the state of the system remains (approximately) in an **instantaneous equilibrium state** of the perturbed Hamiltonian.
- (b) During the switching process the state of the system remains (approximately) in an **instantaneous non-equilibrium (almost) steady state** of the perturbed Hamiltonian.

## Linear response and Kubo's formula

Why should this work? Two possible mechanisms:

- (a) During the switching process the state of the system remains (approximately) in an **instantaneous equilibrium state** of the perturbed Hamiltonian.
- (b) During the switching process the state of the system remains (approximately) in an **instantaneous non-equilibrium (almost) steady state** of the perturbed Hamiltonian.


For a quantum system with Hamiltonian  $H_0$  starting in the gapped ground state  $\rho_0$  scenario (a) occurs, whenever the perturbation does not close the spectral gap.

Then, according to the **adiabatic theorem**, the state  $\rho(t)$  for times  $t \geq 0$ , i.e. when the perturbation is fully switched on, is


$$\rho_\varepsilon = \rho_0^\varepsilon + \mathcal{O}(\varepsilon),$$

where  $\rho_0^\varepsilon$  denotes the ground state of the perturbed Hamiltonian  $H_\varepsilon$ .  
(e.g. **Elgart, Schlein CPAM '04**; **Bachmann et al. CMP '18**)


## Example for (b): The Stark Hamiltonian



## Example for (b): The Stark Hamiltonian




## Example for (b): The Stark Hamiltonian




e.g. **Abou Salem, Fröhlich** *CMP* '07, **Elgart, Hagedorn** *CPAM* '11.


# Example for (b): An “Extended Stark Hamiltonian”



# Example for (b): An "Extended Stark Hamiltonian"



## Example for (b): An “Extended Stark Hamiltonian”



Nenciu *JMP* '02; Panati, Spohn, T. *CMP* '03.

## Interacting fermions on the lattice

First consider systems of interacting fermions on finite sets  $X \subset \mathbb{Z}^d$ .

## Interacting fermions on the lattice

First consider systems of interacting fermions on finite sets  $X \subset \mathbb{Z}^d$ .

The  $N$ -particle Hilbert space for such a system is

$$\mathfrak{h}_{X,N} := \ell^2(X, \mathbb{C}^s)^{\wedge N}$$

## Interacting fermions on the lattice

First consider systems of interacting fermions on finite sets  $X \subset \mathbb{Z}^d$ .

The  $N$ -particle Hilbert space for such a system is

$$\mathfrak{h}_{X,N} := \ell^2(X, \mathbb{C}^s)^{\wedge N}$$

and it is convenient to work on **Fock space**

$$\tilde{\mathfrak{F}}_X := \bigoplus_{N=0}^{s|X|} \mathfrak{h}_{X,N}.$$

## Interacting fermions on the lattice

First consider systems of interacting fermions on finite sets  $X \subset \mathbb{Z}^d$ .

The  $N$ -particle Hilbert space for such a system is

$$\mathfrak{h}_{X,N} := \ell^2(X, \mathbb{C}^s)^{\wedge N}$$

and it is convenient to work on **Fock space**  $\tilde{\mathfrak{F}}_X := \bigoplus_{N=0}^{|X|} \mathfrak{h}_{X,N}$ .

The algebra  $\mathcal{L}(\tilde{\mathfrak{F}}_X)$  of bounded operators on  $\tilde{\mathfrak{F}}_X$  is generated by the **fermionic creation and annihilation operators**  $a_{x,i}^*$  and  $a_{x,i}$ .

By  $\mathcal{A}_X \subset \mathcal{L}(\tilde{\mathfrak{F}}_X)$  we denote the sub-algebra of operators that commute with the number operator  $\mathfrak{n}^X := \sum_{x \in X} a_{x,i}^* a_{x,i}$ .

# Interacting fermions on the lattice

First consider systems of interacting fermions on finite sets  $X \subset \mathbb{Z}^d$ .

The  $N$ -particle Hilbert space for such a system is

$$\mathfrak{h}_{X,N} := \ell^2(X, \mathbb{C}^s)^{\wedge N}$$

and it is convenient to work on **Fock space**  $\mathfrak{F}_X := \bigoplus_{N=0}^{|X|} \mathfrak{h}_{X,N}$ .

The algebra  $\mathcal{L}(\mathfrak{F}_X)$  of bounded operators on  $\mathfrak{F}_X$  is generated by the **fermionic creation and annihilation operators**  $a_{x,i}^*$  and  $a_{x,i}$ .

By  $\mathcal{A}_X \subset \mathcal{L}(\mathfrak{F}_X)$  we denote the sub-algebra of operators that commute with the number operator  $\mathfrak{N}^X := \sum_{x \in X} a_{x,i}^* a_{x,i}$ .

A typical physical Hamiltonian is of the form

$$\begin{aligned} H_0^X &= \sum_{(x,y) \in X^2} a_{x,i}^* T_{ij}^X(x-y) a_{y,j} + \sum_{x \in X} a_{x,i}^* \phi_{ij}^X(x) a_{x,j} \\ &\quad + \sum_{(x,y) \in X^2} a_{x,i}^* a_{x,i} W^X(x-y) a_{y,j}^* a_{y,j} - \mu \mathfrak{N}^X. \end{aligned}$$

## Interacting fermions on the lattice

In order to describe infinite systems of interacting fermions one takes the **thermodynamic limit** of a sequence of finite systems e.g. on cubes  $\Lambda_k := \{-k, \dots, k\}^d \subset \mathbb{Z}^d$ ,  $k \in \mathbb{N}$ .

## Interacting fermions on the lattice

In order to describe infinite systems of interacting fermions one takes the **thermodynamic limit** of a sequence of finite systems e.g. on cubes  $\Lambda_k := \{-k, \dots, k\}^d \subset \mathbb{Z}^d$ ,  $k \in \mathbb{N}$ .

We consider a sequence of Hamiltonians that are **sums of local terms**,

$$H_0^{\Lambda_k} = \sum_{X \subset \Lambda_k} \phi^{\Lambda_k}(X),$$

where  $\phi^{\Lambda_k}(X) \in \mathcal{A}_X$  and  $\|\phi^{\Lambda_k}(X)\|$  is small if  $\text{diam } X$  is large.

## Interacting fermions on the lattice

In order to describe infinite systems of interacting fermions one takes the **thermodynamic limit** of a sequence of finite systems e.g. on cubes  $\Lambda_k := \{-k, \dots, k\}^d \subset \mathbb{Z}^d$ ,  $k \in \mathbb{N}$ .

We consider a sequence of Hamiltonians that are **sums of local terms**,

$$H_0^{\Lambda_k} = \sum_{X \subset \Lambda_k} \Phi^{\Lambda_k}(X),$$

where  $\Phi^{\Lambda_k}(X) \in \mathcal{A}_X$  and  $\|\Phi^{\Lambda_k}(X)\|$  is small if  $\text{diam } X$  is large.

A family  $B = \{B^{\Lambda_k}\}$  of self-adjoint operators  $B^{\Lambda_k}$  indexed by the domain  $\Lambda_k$  and possibly other parameters that is a **sum of local terms**,

$$B^{\Lambda_k} = \sum_{X \subset \Lambda_k} \Phi_B^{\Lambda_k}(X)$$

is called an **"SLT operator family"**. The map  $\Phi_B^{\Lambda_k} : \mathcal{P}(\Lambda_k) \rightarrow \mathcal{A}_{\Lambda_k}$  is called its **interaction**.

## Interacting fermions on the lattice

In order to describe infinite systems of interacting fermions one takes the **thermodynamic limit** of a sequence of finite systems e.g. on cubes  $\Lambda_k := \{-k, \dots, k\}^d \subset \mathbb{Z}^d$ ,  $k \in \mathbb{N}$ .

We consider a sequence of Hamiltonians that are **sums of local terms**,

$$H_0^{\Lambda_k} = \sum_{X \subset \Lambda_k} \Phi^{\Lambda_k}(X),$$

where  $\Phi^{\Lambda_k}(X) \in \mathcal{A}_X$  and  $\|\Phi^{\Lambda_k}(X)\|$  is small if  $\text{diam } X$  is large.

A family  $B = \{B^{\Lambda_k}\}$  of self-adjoint operators  $B^{\Lambda_k}$  indexed by the domain  $\Lambda_k$  and possibly other parameters that is a **sum of local terms**,

$$B^{\Lambda_k} = \sum_{X \subset \Lambda_k} \Phi_B^{\Lambda_k}(X)$$

is called an **"SLT operator family"**. The map  $\Phi_B^{\Lambda_k} : \mathcal{P}(\Lambda_k) \rightarrow \mathcal{A}_{\Lambda_k}$  is called its **interaction**. Typically

$$\|B^{\Lambda_k}\| \sim |\Lambda_k| = (2k+1)^d.$$

# Interacting fermions on the lattice

To quantify locality of SLT operators, one defines spaces  $\mathcal{B}_\zeta$  of SLT operators with norm

$$\|\Phi\|_\zeta := \sup_{k \in \mathbb{N}} \sup_{x, y \in \Lambda_k} \sum_{\{x, y\} \subset X \subset \Lambda_k} \frac{\|\Phi^{\Lambda_k}(X)\|}{\zeta(d^{\Lambda_k}(x, y))} =: \sup_{k \in \mathbb{N}} \|\Phi\|_{\zeta, \Lambda_k},$$

where  $\zeta : [0, \infty) \rightarrow (0, \infty)$  is a rapidly decaying function, e.g.  
 $\zeta(r) = e^{-ar}$

## Interacting fermions on the lattice

To quantify locality of SLT operators, one defines spaces  $\mathcal{B}_\zeta$  of SLT operators with norm

$$\|\Phi\|_\zeta := \sup_{k \in \mathbb{N}} \sup_{x, y \in \Lambda_k} \sum_{\{x, y\} \subset X \subset \Lambda_k} \frac{\|\Phi^{\Lambda_k}(X)\|}{\zeta(d^{\Lambda_k}(x, y))} =: \sup_{k \in \mathbb{N}} \|\Phi\|_{\zeta, \Lambda_k},$$

where  $\zeta : [0, \infty) \rightarrow (0, \infty)$  is a rapidly decaying function, e.g.  
 $\zeta(r) = e^{-ar}$

### Definition

One says that an SLT operator in  $\mathcal{B}_\zeta$  has a **thermodynamic limit**, if for all  $M \in \mathbb{N}$  and  $\delta > 0$  there exists  $K \geq M$  such that for all  $l, k \geq K$

$$\left\| \Phi^{\Lambda_l} - \Phi^{\Lambda_k} \right\|_{\zeta, \Lambda_M} \leq \delta.$$

# Interacting fermions on the lattice

To quantify locality of SLT operators, one defines spaces  $\mathcal{B}_\zeta$  of SLT operators with norm

$$\|\Phi\|_\zeta := \sup_{k \in \mathbb{N}} \sup_{x, y \in \Lambda_k} \sum_{\{x, y\} \subset X \subset \Lambda_k} \frac{\|\Phi^{\Lambda_k}(X)\|}{\zeta(d^{\Lambda_k}(x, y))} =: \sup_{k \in \mathbb{N}} \|\Phi\|_{\zeta, \Lambda_k},$$

where  $\zeta : [0, \infty) \rightarrow (0, \infty)$  is a rapidly decaying function, e.g.  
 $\zeta(r) = e^{-ar}$

## Definition

We say that an SLT operator in  $\mathcal{B}_\zeta$  has a **rapid thermodynamic limit** with exponent  $\gamma \in (0, 1)$ , if there exist  $\lambda, C > 0$  such that for all  $M \in \mathbb{N}$  and for all  $l, k \geq M + \lambda M^\gamma$

$$\left\| \Phi^{\Lambda_l} - \Phi^{\Lambda_k} \right\|_{\zeta, \Lambda_M} \leq C \zeta(M^\gamma) =: C \zeta_\gamma(M).$$

# Interacting fermions on the lattice

To quantify locality of SLT operators, one defines spaces  $\mathcal{B}_\zeta$  of SLT operators with norm

$$\|\Phi\|_\zeta := \sup_{k \in \mathbb{N}} \sup_{x, y \in \Lambda_k} \sum_{\{x, y\} \subset X \subset \Lambda_k} \frac{\|\Phi^{\Lambda_k}(X)\|}{\zeta(d^{\Lambda_k}(x, y))} =: \sup_{k \in \mathbb{N}} \|\Phi\|_{\zeta, \Lambda_k},$$

where  $\zeta : [0, \infty) \rightarrow (0, \infty)$  is a rapidly decaying function, e.g.  
 $\zeta(r) = e^{-ar}$

## Definition

We say that an SLT operator in  $\mathcal{B}_\zeta$  has a **rapid thermodynamic limit** with exponent  $\gamma \in (0, 1)$ , if there exist  $\lambda, C > 0$  such that for all  $M \in \mathbb{N}$  and for all  $l, k \geq M + \lambda M^\gamma$

$$\left\| \Phi^{\Lambda_l} - \Phi^{\Lambda_k} \right\|_{\zeta, \Lambda_M} \leq C \zeta(M^\gamma) =: C \zeta_\gamma(M).$$

**However, there is no limiting Hamiltonian for the infinite system!**

## Interacting fermions on the lattice

Since for  $Y \subset X$  we have  $\mathcal{A}_Y \subset \mathcal{A}_X$ , one can define the algebra of local observables as

$$\mathcal{A}_{\text{loc}} := \bigcup_{X \subset \mathbb{Z}^d, |X| < \infty} \mathcal{A}_X.$$

## Interacting fermions on the lattice

Since for  $Y \subset X$  we have  $\mathcal{A}_Y \subset \mathcal{A}_X$ , one can define the algebra of local observables as

$$\mathcal{A}_{\text{loc}} := \bigcup_{X \subset \mathbb{Z}^d, |X| < \infty} \mathcal{A}_X.$$

The quasi-local algebra is the  $C^*$ -algebra

$$\mathcal{A} := \overline{\mathcal{A}_{\text{loc}}}^{\|\cdot\|}$$

# Interacting fermions on the lattice

Since for  $Y \subset X$  we have  $\mathcal{A}_Y \subset \mathcal{A}_X$ , one can define the **algebra of local observables** as

$$\mathcal{A}_{\text{loc}} := \bigcup_{X \subset \mathbb{Z}^d, |X| < \infty} \mathcal{A}_X.$$

The **quasi-local algebra** is the  $C^*$ -algebra

$$\mathcal{A} := \overline{\mathcal{A}_{\text{loc}}}^{\|\cdot\|}$$

In order to regain control on the localisation properties of elements of  $\mathcal{A}$ , one defines sub-algebras  $\mathcal{D}_\zeta \subset \mathcal{A}$  with norm

$$\|B\|_\zeta := \|B\| + \sup_{k \in \mathbb{N}} \left( \frac{\|(1 - \mathbb{E}_{\Lambda_k})(B)\|}{\zeta(k)} \right) < \infty,$$

where  $\zeta : [0, \infty) \rightarrow (0, \infty)$  is again a rapidly decaying function and  $\mathbb{E}_{\Lambda_k} : \mathcal{A} \rightarrow \mathcal{A}_{\Lambda_k}$  denotes the **conditional expectation**.

# Interacting fermions on the lattice

## Proposition

Let  $H_0 \in \mathcal{B}_\zeta$  have a thermodynamic limit.

Then for any  $B \in \mathcal{A}_{\text{loc}}$  the limit

$$\mathfrak{U}_t(B) := \lim_{k \rightarrow \infty} e^{iH_0^{\wedge k} t} B e^{-iH_0^{\wedge k} t} \in \mathcal{A}$$

exists and defines a one-parameter family  $t \mapsto \mathfrak{U}_t$  of automorphisms of the algebra  $\mathcal{A}$  with densely defined generator  $\mathcal{L}_{H_0} : D(\mathcal{L}_{H_0}) \rightarrow \mathcal{A}$ .

# Interacting fermions on the lattice

## Proposition

Let  $H_0 \in \mathcal{B}_\zeta$  have a thermodynamic limit.

Then for any  $B \in \mathcal{A}_{\text{loc}}$  the limit

$$\mathfrak{U}_t(B) := \lim_{k \rightarrow \infty} e^{iH_0^{\wedge k} t} B e^{-iH_0^{\wedge k} t} \in \mathcal{A}$$

exists and defines a one-parameter family  $t \mapsto \mathfrak{U}_t$  of automorphisms of the algebra  $\mathcal{A}$  with densely defined generator  $\mathcal{L}_{H_0} : D(\mathcal{L}_{H_0}) \rightarrow \mathcal{A}$ .

Moreover, for suitable pairs  $f_1, f_2$  of rapidly decaying functions,

$$\mathfrak{U}_t : \mathcal{D}_{f_1} \rightarrow \mathcal{D}_{f_2}$$

is a bounded operator and  $\mathfrak{U}_t^{\wedge k} \circ \mathbb{E}_{\Lambda_k} \rightarrow \mathfrak{U}_t$  in norm.

# Interacting fermions on the lattice

## Proposition

Let  $H_0 \in \mathcal{B}_\zeta$  have a thermodynamic limit.

Then for any  $B \in \mathcal{A}_{\text{loc}}$  the limit

$$\mathfrak{U}_t(B) := \lim_{k \rightarrow \infty} e^{iH_0^{\wedge k} t} B e^{-iH_0^{\wedge k} t} \in \mathcal{A}$$

exists and defines a one-parameter family  $t \mapsto \mathfrak{U}_t$  of automorphisms of the algebra  $\mathcal{A}$  with densely defined generator  $\mathcal{L}_{H_0} : D(\mathcal{L}_{H_0}) \rightarrow \mathcal{A}$ .

Moreover, for suitable pairs  $f_1, f_2$  of rapidly decaying functions,

$$\mathfrak{U}_t : \mathcal{D}_{f_1} \rightarrow \mathcal{D}_{f_2}$$

is a bounded operator and  $\mathfrak{U}_t^{\wedge k} \circ \mathbb{E}_{\Lambda_k} \rightarrow \mathfrak{U}_t$  in norm.

Also the **Liouvillian**

$$\mathcal{L}_{H_0} : \mathcal{D}_{f_1} \rightarrow \mathcal{D}_{f_2}, \quad \mathcal{L}_{H_0}(B) := \lim_{k \rightarrow \infty} [H_0^{\wedge k}, \mathbb{E}_{\Lambda_k}(B)]$$

is a bounded operator and the convergence is in norm.

# Interacting fermions on the lattice

## Proposition

Let  $H_0 \in \mathcal{B}_\zeta$  have a thermodynamic limit.

Then for any  $B \in \mathcal{A}_{\text{loc}}$  the limit

$$\mathfrak{U}_t(B) := \lim_{k \rightarrow \infty} e^{iH_0^{\Lambda_k} t} B e^{-iH_0^{\Lambda_k} t} \in \mathcal{A}$$

exists and defines a one-parameter family  $t \mapsto \mathfrak{U}_t$  of automorphisms of the algebra  $\mathcal{A}$  with densely defined generator  $\mathcal{L}_{H_0} : D(\mathcal{L}_{H_0}) \rightarrow \mathcal{A}$ .

If  $H_0 \in \mathcal{B}_\zeta$  has a rapid thermodynamic limit with exponent  $\gamma \in (0, 1)$ , then there exist  $\lambda_1 > 0$ ,  $\lambda_2 \in (0, 1)$ , and  $C < \infty$ , such that for all  $l, k \in \mathbb{N}$  with  $l \geq k$ ,  $X \subset \Lambda_k$  and  $B \in \mathcal{A}_X$

$$\begin{aligned} \left\| \left( \mathfrak{U}_t^{\Lambda_l} - \mathfrak{U}_t^{\Lambda_k} \right) (B) \right\| &\leq C \|B\| \text{diam}(X)^{d+1} e^{2C_\zeta |t-s| \|\Phi_{H_0}\|_\zeta} |t-s| \\ &\quad \times \zeta_\gamma \left( \text{dist}^{\Lambda_l}(X, \Lambda_l \setminus \Lambda_{\max\{\lceil k - \lambda_1 k^\gamma \rceil, \lceil \lambda_2 \cdot k \rceil\}}) \right) \end{aligned}$$

## Adiabatic theorem

From now on we consider a time-dependent SLT Hamiltonian  $H_0(t) \in \mathcal{B}_{e^{-a}}$ ,  $t \in I \subset \mathbb{R}$ , possibly perturbed by a time-dependent operator  $\varepsilon V(t)$ , where  $V(t) = V_v(t) + H_1(t)$  is the sum of an SLT operator  $H_1(t)$  and a Lipschitz potential  $V_v(t)$ , i.e.

$$V_v^{\Lambda_k}(t) = \sum_{x \in \Lambda_k} v(x) a_x^* a_x.$$

## Adiabatic theorem

From now on we consider a time-dependent SLT Hamiltonian  $H_0(t) \in \mathcal{B}_{e^{-a}}$ ,  $t \in I \subset \mathbb{R}$ , possibly perturbed by a time-dependent operator  $\varepsilon V(t)$ , where  $V(t) = V_v(t) + H_1(t)$  is the sum of an SLT operator  $H_1(t)$  and a Lipschitz potential  $V_v(t)$ , i.e.

$$V_v^{\Lambda_k}(t) = \sum_{x \in \Lambda_k} v(x) a_x^* a_x.$$

Let

$$H_\varepsilon(t) := H_0(t) + \varepsilon V(t).$$

## Adiabatic theorem

From now on we consider a time-dependent SLT Hamiltonian  $H_0(t) \in \mathcal{B}_{e^{-a}}$ ,  $t \in I \subset \mathbb{R}$ , possibly perturbed by a time-dependent operator  $\varepsilon V(t)$ , where  $V(t) = V_v(t) + H_1(t)$  is the sum of an SLT operator  $H_1(t)$  and a Lipschitz potential  $V_v(t)$ , i.e.

$$V_v^{\Lambda_k}(t) = \sum_{x \in \Lambda_k} v(x) a_x^* a_x.$$

Let

$$H_\varepsilon(t) := H_0(t) + \varepsilon V(t).$$

Similar results as before hold for the corresponding **adiabatic evolution family**  $\mathfrak{U}_{t,t_0}^{\eta,\varepsilon}$  generated by the time-dependent Liouvillian  $\frac{1}{\eta} \mathcal{L}_{H_\varepsilon(t)}$  with adiabatic parameter  $\eta > 0$ , i.e.

$$\mathfrak{U}_{t,t_0}^{\eta,\varepsilon}(B) := \lim_{k \rightarrow \infty} \mathfrak{U}_{t,t_0}^{\eta,\varepsilon,\Lambda_k}(B) \in \mathcal{A}$$

# Adiabatic theorem

## Standard gap assumption

Assume that smallest eigenvalue  $E_0^{\Lambda_k}(t)$  (ground state) of  $H_0^{\Lambda_k}(t)$  is separated from the rest of the spectrum uniformly in the volume  $|\Lambda_k|$ ,

$$\inf_{\Lambda_k} \text{dist} \left( E_0^{\Lambda_k}(t), \sigma(H_0^{\Lambda_k}(t)) \setminus \{E_0^{\Lambda_k}(t)\} \right) =: g > 0.$$

# Adiabatic theorem

## Standard gap assumption

Assume that smallest eigenvalue  $E_0^{\Lambda_k}(t)$  (ground state) of  $H_0^{\Lambda_k}(t)$  is separated from the rest of the spectrum uniformly in the volume  $|\Lambda_k|$ ,

$$\inf_{\Lambda_k} \text{dist} \left( E_0^{\Lambda_k}(t), \sigma(H_0^{\Lambda_k}(t)) \setminus \{E_0^{\Lambda_k}(t)\} \right) =: g > 0.$$

## Examples

- ▶ Electrons in a Chern-trivial insulator, i.e. with the chemical potential  $\mu$  in a “band gap”.

# Adiabatic theorem

## Standard gap assumption

Assume that smallest eigenvalue  $E_0^{\Lambda_k}(t)$  (ground state) of  $H_0^{\Lambda_k}(t)$  is separated from the rest of the spectrum uniformly in the volume  $|\Lambda_k|$ ,

$$\inf_{\Lambda_k} \text{dist} \left( E_0^{\Lambda_k}(t), \sigma(H_0^{\Lambda_k}(t)) \setminus \{E_0^{\Lambda_k}(t)\} \right) =: g > 0.$$

## Examples

- ▶ Electrons in a Chern-trivial insulator, i.e. with the chemical potential  $\mu$  in a “band gap”.
- ▶ Electrons in a Chern-nontrivial insulator with periodic boundary conditions.

# Adiabatic theorem

## Standard gap assumption

Assume that smallest eigenvalue  $E_0^{\Lambda_k}(t)$  (ground state) of  $H_0^{\Lambda_k}(t)$  is separated from the rest of the spectrum uniformly in the volume  $|\Lambda_k|$ ,

$$\inf_{\Lambda_k} \text{dist} \left( E_0^{\Lambda_k}(t), \sigma(H_0^{\Lambda_k}(t)) \setminus \{E_0^{\Lambda_k}(t)\} \right) =: g > 0.$$

## Examples

- ▶ Electrons in a Chern-trivial insulator, i.e. with the chemical potential  $\mu$  in a “band gap”.
- ▶ Electrons in a Chern-nontrivial insulator with periodic boundary conditions.
- ▶ The filled Dirac sea.

# Adiabatic theorem

## Standard gap assumption

Assume that smallest eigenvalue  $E_0^{\Lambda_k}(t)$  (ground state) of  $H_0^{\Lambda_k}(t)$  is separated from the rest of the spectrum uniformly in the volume  $|\Lambda_k|$ ,

$$\inf_{\Lambda_k} \text{dist} \left( E_0^{\Lambda_k}(t), \sigma(H_0^{\Lambda_k}(t)) \setminus \{E_0^{\Lambda_k}(t)\} \right) =: g > 0.$$

Adiabatic theorems under the “**standard gap assumption**” in **finite volumes** with error estimates that are uniform in the volume were shown by **Bachmann, De Roeck, Fraas** CMP '18, **Monaco, T.**, RMP '19 and **T.**, CMP '20.

# Adiabatic theorem

## Standard gap assumption

Assume that smallest eigenvalue  $E_0^{\Lambda_k}(t)$  (ground state) of  $H_0^{\Lambda_k}(t)$  is separated from the rest of the spectrum uniformly in the volume  $|\Lambda_k|$ ,

$$\inf_{\Lambda_k} \text{dist} \left( E_0^{\Lambda_k}(t), \sigma(H_0^{\Lambda_k}(t)) \setminus \{E_0^{\Lambda_k}(t)\} \right) =: g > 0.$$

Adiabatic theorems under the “standard gap assumption” in finite volumes with error estimates that are uniform in the volume were shown by **Bachmann, De Roeck, Fraas** CMP '18, **Monaco, T.**, RMP '19 and **T.**, CMP '20.

In **Henheik, T.** '20 we prove an adiabatic theorem for  $\mathfrak{U}_{t,t_0}$  on  $\mathcal{A}$  and apply it to linear response.

## Adiabatic theorem with a gap in the bulk

**Motivation:** Response of Chern-nontrivial systems (e.g. quantum Hall systems), where the Hamiltonian has no spectral gap due to edge states.

## Adiabatic theorem with a gap in the bulk

**Motivation:** Response of Chern-nontrivial systems (e.g. quantum Hall systems), where the Hamiltonian has no spectral gap due to edge states.

**Idea:** Require the spectral gap only for the infinite system.

## Adiabatic theorem with a gap in the bulk

**Motivation:** Response of Chern-nontrivial systems (e.g. quantum Hall systems), where the Hamiltonian has no spectral gap due to edge states.

**Idea:** Require the spectral gap only for the infinite system.

**Problem:** There is no limiting Hamiltonian for the infinite system.

## Adiabatic theorem with a gap in the bulk

**Motivation:** Response of Chern-nontrivial systems (e.g. quantum Hall systems), where the Hamiltonian has no spectral gap due to edge states.

**Idea:** Require the spectral gap only for the infinite system.

**Problem:** There is no limiting Hamiltonian for the infinite system.

A state  $\rho$  on  $\mathcal{A}$  is called a  $\mathcal{L}_{H_0}$ -ground state, iff

$$\rho(B^* \mathcal{L}_{H_0}(B)) \geq 0 \quad \text{for all } B \in D(\mathcal{L}_{H_0}).$$

## Adiabatic theorem with a gap in the bulk

**Motivation:** Response of Chern-nontrivial systems (e.g. quantum Hall systems), where the Hamiltonian has no spectral gap due to edge states.

**Idea:** Require the spectral gap only for the infinite system.

**Problem:** There is no limiting Hamiltonian for the infinite system.

A state  $\rho$  on  $\mathcal{A}$  is called a  $\mathcal{L}_{H_0}$ -ground state, iff

$$\rho(B^* \mathcal{L}_{H_0}(B)) \geq 0 \quad \text{for all } B \in D(\mathcal{L}_{H_0}).$$

Let  $\rho$  be a  $\mathcal{L}_{H_0}$ -ground state and  $(\mathcal{H}_\rho, \pi_\rho, \Omega_\rho)$  be the corresponding GNS triple.

## Adiabatic theorem with a gap in the bulk

**Motivation:** Response of Chern-nontrivial systems (e.g. quantum Hall systems), where the Hamiltonian has no spectral gap due to edge states.

**Idea:** Require the spectral gap only for the infinite system.

**Problem:** There is no limiting Hamiltonian for the infinite system.

A state  $\rho$  on  $\mathcal{A}$  is called a  $\mathcal{L}_{H_0}$ -ground state, iff

$$\rho(B^* \mathcal{L}_{H_0}(B)) \geq 0 \quad \text{for all } B \in D(\mathcal{L}_{H_0}).$$

Let  $\rho$  be a  $\mathcal{L}_{H_0}$ -ground state and  $(\mathcal{H}_\rho, \pi_\rho, \Omega_\rho)$  be the corresponding GNS triple. Then there exists a unique densely defined, self-adjoint positive operator  $H_\rho \geq 0$  on  $\mathcal{H}_\rho$  satisfying

$$\pi_\rho(e^{is\mathcal{L}_{H_0}}(B)) = e^{isH_\rho} \pi_\rho(B) e^{-isH_\rho} \quad \text{and} \quad e^{-isH_\rho} \Omega_\rho = \Omega_\rho$$

for all  $B \in \mathcal{A}$  and  $s \in \mathbb{R}$ .

## Adiabatic theorem with a gap in the bulk

**Motivation:** Response of Chern-nontrivial systems (e.g. quantum Hall systems), where the Hamiltonian has no spectral gap due to edge states.

**Idea:** Require the spectral gap only for the infinite system.

**Problem:** There is no limiting Hamiltonian for the infinite system.

A state  $\rho$  on  $\mathcal{A}$  is called a  $\mathcal{L}_{H_0}$ -ground state, iff

$$\rho(B^* \mathcal{L}_{H_0}(B)) \geq 0 \quad \text{for all } B \in D(\mathcal{L}_{H_0}).$$

Let  $\rho$  be a  $\mathcal{L}_{H_0}$ -ground state and  $(\mathcal{H}_\rho, \pi_\rho, \Omega_\rho)$  be the corresponding GNS triple. Then there exists a unique densely defined, self-adjoint positive operator  $H_\rho \geq 0$  on  $\mathcal{H}_\rho$  satisfying

$$\pi_\rho(e^{is\mathcal{L}_{H_0}}(B)) = e^{isH_\rho} \pi_\rho(B) e^{-isH_\rho} \quad \text{and} \quad e^{-isH_\rho} \Omega_\rho = \Omega_\rho$$

for all  $B \in \mathcal{A}$  and  $s \in \mathbb{R}$ .

$H_\rho$  is called the **bulk Hamiltonian** associated with  $\rho$ .

# Adiabatic theorem with a gap in the bulk

## Gap assumption in the bulk (cf. Moon, Ogata, JFA '19)

There exists  $g > 0$  such that for each  $t \in I$  the Liouvillian  $\mathcal{L}_{H_0(t)}$  has a **unique ground state**  $\rho_t$  and

$$\sigma(H_{\rho_t}) \setminus \{0\} \subset [g, \infty).$$

# Adiabatic theorem with a gap in the bulk

## Adiabatic theorem

Let the Hamiltonian  $H_\varepsilon(t) = H_0(t) + \varepsilon V(t)$  satisfy the previous assumptions and denote by  $\mathcal{U}_{t,t_0}^{\varepsilon,\eta}$  the Heisenberg time-evolution it generates on  $\mathcal{A}$ .

# Adiabatic theorem with a gap in the bulk

## Adiabatic theorem

Let the Hamiltonian  $H_\varepsilon(t) = H_0(t) + \varepsilon V(t)$  satisfy the previous assumptions and denote by  $\mathfrak{U}_{t,t_0}^{\varepsilon,\eta}$  the Heisenberg time-evolution it generates on  $\mathcal{A}$ .

Then for any  $\varepsilon, \eta \in (0, 1]$  and  $t \in I$  there exists a near-identity automorphism  $\beta^{\varepsilon,\eta}(t)$  of  $\mathcal{A}$  such that the **super-adiabatic NEASS** defined by

$$\Pi_t^{\varepsilon,\eta} := \rho_t \circ \beta_t^{\varepsilon,\eta}$$

has the following properties:

# Adiabatic theorem with a gap in the bulk

## Adiabatic theorem

Let the Hamiltonian  $H_\varepsilon(t) = H_0(t) + \varepsilon V(t)$  satisfy the previous assumptions and denote by  $\mathfrak{U}_{t,t_0}^{\varepsilon,\eta}$  the Heisenberg time-evolution it generates on  $\mathcal{A}$ .

Then for any  $\varepsilon, \eta \in (0, 1]$  and  $t \in I$  there exists a near-identity automorphism  $\beta^{\varepsilon,\eta}(t)$  of  $\mathcal{A}$  such that the **super-adiabatic NEASS** defined by

$$\Pi_t^{\varepsilon,\eta} := \rho_t \circ \beta_t^{\varepsilon,\eta}$$

has the following properties:

**(1)** It almost **intertwines the time evolution**: For any  $n \in \mathbb{N}$  and any  $f \in \mathcal{S}$ , there exists a constant  $C_n$  such that for any  $t \in I$  and  $B \in \mathcal{D}_f$

$$\begin{aligned} & \left| \left( \Pi_{t_0}^{\varepsilon,\eta} \circ \mathfrak{U}_{t,t_0}^{\varepsilon,\eta} - \Pi_t^{\varepsilon,\eta} \right) (B) \right| \\ & \leq C_n \frac{\varepsilon^{n+1} + \eta^{n+1}}{\eta^{d+1}} \left( 1 + |t - t_0|^{d+1} \right) \|B\|_f. \end{aligned}$$

# Adiabatic theorem with a gap in the bulk

## Adiabatic theorem

Let the Hamiltonian  $H_\varepsilon(t) = H_0(t) + \varepsilon V(t)$  satisfy the previous assumptions and denote by  $\mathfrak{U}_{t,t_0}^{\varepsilon,\eta}$  the Heisenberg time-evolution it generates on  $\mathcal{A}$ .

Then for any  $\varepsilon, \eta \in (0, 1]$  and  $t \in I$  there exists a near-identity automorphism  $\beta^{\varepsilon,\eta}(t)$  of  $\mathcal{A}$  such that the **super-adiabatic NEASS** defined by

$$\Pi_t^{\varepsilon,\eta} := \rho_t \circ \beta_t^{\varepsilon,\eta}$$

has the following properties:

(2) It is **local in time**:  $\beta_t^{\varepsilon,\eta}$  depends only on  $H_\varepsilon$  and its time derivatives at time  $t$ .

# Adiabatic theorem with a gap in the bulk

## Adiabatic theorem

Let the Hamiltonian  $H_\varepsilon(t) = H_0(t) + \varepsilon V(t)$  satisfy the previous assumptions and denote by  $\mathfrak{U}_{t,t_0}^{\varepsilon,\eta}$  the Heisenberg time-evolution it generates on  $\mathcal{A}$ .

Then for any  $\varepsilon, \eta \in (0, 1]$  and  $t \in I$  there exists a near-identity automorphism  $\beta^{\varepsilon,\eta}(t)$  of  $\mathcal{A}$  such that the **super-adiabatic NEASS** defined by

$$\Pi_t^{\varepsilon,\eta} := \rho_t \circ \beta_t^{\varepsilon,\eta}$$

has the following properties:

(2) It is **local in time**:  $\beta_t^{\varepsilon,\eta}$  depends only on  $H_\varepsilon$  and its time derivatives at time  $t$ .

(3) It is **stationary** whenever the Hamiltonian is stationary: if  $H_\varepsilon$  is constant on an interval  $J \subset I$  then  $\Pi_t^{\varepsilon,\eta} = \Pi_t^{\varepsilon,0}$  is constant for  $t \in J$ .

# Adiabatic theorem with a gap in the bulk

## Adiabatic theorem

Let the Hamiltonian  $H_\varepsilon(t) = H_0(t) + \varepsilon V(t)$  satisfy the previous assumptions and denote by  $\mathfrak{U}_{t,t_0}^{\varepsilon,\eta}$  the Heisenberg time-evolution it generates on  $\mathcal{A}$ .

Then for any  $\varepsilon, \eta \in (0, 1]$  and  $t \in I$  there exists a near-identity automorphism  $\beta^{\varepsilon,\eta}(t)$  of  $\mathcal{A}$  such that the **super-adiabatic NEASS** defined by

$$\Pi_t^{\varepsilon,\eta} := \rho_t \circ \beta_t^{\varepsilon,\eta}$$

has the following properties:

- (2) It is **local in time**:  $\beta_t^{\varepsilon,\eta}$  depends only on  $H_\varepsilon$  and its time derivatives at time  $t$ .
- (3) It is **stationary** whenever the Hamiltonian is stationary: if  $H_\varepsilon$  is constant on an interval  $J \subset I$  then  $\Pi_t^{\varepsilon,\eta} = \Pi_t^{\varepsilon,0}$  is constant for  $t \in J$ .
- (4)  $\Pi_t^{\varepsilon,0}$  has an **explicit asymptotic expansion** in powers of  $\varepsilon$ .

# Adiabatic theorem with a gap in the bulk

## Adiabatic theorem

Let the Hamiltonian  $H_\varepsilon(t) = H_0(t) + \varepsilon V(t)$  satisfy the previous assumptions and denote by  $\mathfrak{U}_{t,t_0}^{\varepsilon,\eta}$  the Heisenberg time-evolution it generates on  $\mathcal{A}$ .

Then for any  $\varepsilon, \eta \in (0, 1]$  and  $t \in I$  there exists a near-identity automorphism  $\beta^{\varepsilon,\eta}(t)$  of  $\mathcal{A}$  such that the **super-adiabatic NEASS** defined by

$$\Pi_t^{\varepsilon,\eta} := \rho_t \circ \beta_t^{\varepsilon,\eta}$$

has the following properties:

**(5)** It equals the ground state of  $H_0$  whenever the perturbation vanishes and the Hamiltonian is stationary: if for some  $t \in I$  all time-derivatives of  $H_\varepsilon$  vanish at time  $t$  and  $V(t) = 0$ , then

$$\Pi_t^{\varepsilon,\eta} = \Pi_t^{\varepsilon,0} = \rho_t.$$

## Remark on time-scales

For  $\varepsilon \neq 0$ , the right hand side of

$$\begin{aligned} & |(\Pi_{t_0}^{\varepsilon, \eta} \circ \mathfrak{U}_{t, t_0}^{\varepsilon, \eta} - \Pi_t^{\varepsilon, \eta})(B)| \\ & \leq C_n \frac{\varepsilon^{n+1} + \eta^{n+1}}{\eta^{d+1}} \left(1 + |t - t_0|^{d+1}\right) \|B\|_f \end{aligned}$$

shows that the admissible adiabatic time scales  $\eta$  are coupled to the strength  $\varepsilon$  of the perturbation:

The adiabatic parameter  $\eta$  needs to be small, but not too small. The adiabatic switching must occur on time-scales that are fast compared to the life-time of the NEASS, i.e.  $\eta \gtrsim \varepsilon^m$  for some  $m \in \mathbb{N}$ .

## Remark on finite domains

Under an additional assumption on the rate of convergence in

$$\rho^{\Lambda_k} \rightarrow \rho$$

we show that a similar adiabatic theorem holds also for **finite systems** up to an additional error term that decays faster than any inverse polynomial in the system size.

## Remark on finite domains

Under an additional assumption on the rate of convergence in

$$\rho^{\Lambda_k} \rightarrow \rho$$

we show that a similar adiabatic theorem holds also for **finite systems** up to an additional error term that decays faster than any inverse polynomial in the system size.

There exists  $\lambda > 0$  such that for any  $n \in \mathbb{N}$  there exists a constant  $C_n$  and for any compact  $K \subset I$  and  $m \in \mathbb{N}$  there exists a constant  $\tilde{C}_{n,m,K}$  such that for all  $k \in \mathbb{N}$ , all finite  $X \subset \Lambda_k$ , all  $B \in \mathcal{A}_X$ , and all  $t, t_0 \in K$

$$\begin{aligned} & \left| \left( \Pi_{t_0}^{\varepsilon, \eta, \Lambda_k} \circ \mathcal{U}_{t, t_0}^{\varepsilon, \eta, \Lambda_k} - \Pi_t^{\varepsilon, \eta, \Lambda_k} \right) (B) \right| \\ & \leq C_n \frac{\varepsilon^{n+1} + \eta^{n+1}}{\eta^{d+1}} \left( 1 + |t - t_0|^{d+1} \right) \|B\| |X|^2 \\ & \quad + \tilde{C}_{n,m,K} \left( 1 + \eta \operatorname{dist}(X, \Gamma \setminus \Lambda_{\lfloor k - \lambda k \gamma \rfloor}) \right)^{-m} \|B\| \operatorname{diam}(X)^{2d}. \end{aligned}$$

## Elements of the proof: the local inverse of the Liouvillian

In finite volume  $\Lambda$ , where  $\rho$  is the ground state projection, the construction of the (stationary) NEASS proceeds as follows:

## Elements of the proof: the local inverse of the Liouvillian

In finite volume  $\Lambda$ , where  $\rho$  is the ground state projection, the construction of the (stationary) NEASS proceeds as follows:

With  $\Pi := e^{i\varepsilon S} \rho e^{-i\varepsilon S}$  and  $S := \sum_{\mu=1}^n \varepsilon^{\mu-1} A_{\mu}$  we have

$$[H, \Pi] = \left[ H_0 + \varepsilon V, e^{i\varepsilon S} \rho e^{-i\varepsilon S} \right]$$

## Elements of the proof: the local inverse of the Liouvillian

In finite volume  $\Lambda$ , where  $\rho$  is the ground state projection, the construction of the (stationary) NEASS proceeds as follows:

With  $\Pi := e^{i\varepsilon S} \rho e^{-i\varepsilon S}$  and  $S := \sum_{\mu=1}^n \varepsilon^{\mu-1} A_{\mu}$  we have

$$\begin{aligned} [H, \Pi] &= \left[ H_0 + \varepsilon V, e^{i\varepsilon S} \rho e^{-i\varepsilon S} \right] \\ &= i\varepsilon e^{i\varepsilon S} \left[ \mathcal{L}_{H_0}(A_1) - iV, \rho \right] e^{-i\varepsilon S} + \mathcal{O}(\varepsilon^2). \end{aligned}$$

## Elements of the proof: the local inverse of the Liouvillian

In finite volume  $\Lambda$ , where  $\rho$  is the ground state projection, the construction of the (stationary) NEASS proceeds as follows:

With  $\Pi := e^{i\varepsilon S} \rho e^{-i\varepsilon S}$  and  $S := \sum_{\mu=1}^n \varepsilon^{\mu-1} A_{\mu}$  we have

$$\begin{aligned} [H, \Pi] &= \left[ H_0 + \varepsilon V, e^{i\varepsilon S} \rho e^{-i\varepsilon S} \right] \\ &= i\varepsilon e^{i\varepsilon S} \left[ \mathcal{L}_{H_0}(A_1) - iV, \rho \right] e^{-i\varepsilon S} + \mathcal{O}(\varepsilon^2). \end{aligned}$$

We thus need to choose  $A_1$  such that

$$\left[ \mathcal{L}_{H_0}(A_1) - iV, \rho \right] \stackrel{!}{=} 0$$

## Elements of the proof: the local inverse of the Liouvillian

Assuming a spectral gap for  $H_0$ , **Bachmann, Michalakis, Nachtergaele, Sims**, CMP '12 (based on **Hastings, Wen**, PRB '05) constructed a linear map

$$\mathcal{I}_{H_0, g}^\Lambda : \mathcal{A}_\Lambda \rightarrow \mathcal{A}_\Lambda$$

that maps SLT operators to SLT operators and satisfies

$$\mathcal{I}_{H_0, g}^\Lambda |_{\mathcal{A}_\Lambda^{\text{OD}}} = i \mathcal{L}_{H_0}^{-1} |_{\mathcal{A}_\Lambda^{\text{OD}}} .$$

## Elements of the proof: the local inverse of the Liouvillian

Assuming a spectral gap for  $H_0$ , **Bachmann, Michalakis, Nachtergaele, Sims**, CMP '12 (based on **Hastings, Wen**, PRB '05) constructed a linear map

$$\mathcal{I}_{H_0, g}^\wedge : \mathcal{A}_\Lambda \rightarrow \mathcal{A}_\Lambda$$

that maps SLT operators to SLT operators and satisfies

$$\mathcal{I}_{H_0, g}^\wedge |_{\mathcal{A}_\Lambda^{\text{OD}}} = i \mathcal{L}_{H_0}^{-1} |_{\mathcal{A}_\Lambda^{\text{OD}}}.$$

Thus, for all  $A \in \mathcal{A}_\Lambda$ ,

$$\left[ \mathcal{L}_{H_0} \circ \mathcal{I}_{H_0, g}^\wedge(A) - iA, \rho \right] = 0$$

and one can choose  $A_1 = \mathcal{I}_{H_0, g}^\wedge(V)$ .

## Elements of the proof: the local inverse of the Liouvillian

Assuming a spectral gap for  $H_0$ , **Bachmann, Michalakis, Nachtergaele, Sims**, CMP '12 (based on **Hastings, Wen**, PRB '05) constructed a linear map

$$\mathcal{I}_{H_0, g}^\Lambda : \mathcal{A}_\Lambda \rightarrow \mathcal{A}_\Lambda$$

that maps SLT operators to SLT operators and satisfies

$$\mathcal{I}_{H_0, g}^\Lambda |_{\mathcal{A}_\Lambda^{\text{OD}}} = i \mathcal{L}_{H_0}^{-1} |_{\mathcal{A}_\Lambda^{\text{OD}}}.$$

Thus, for all  $A \in \mathcal{A}_\Lambda$ ,

$$\left[ \mathcal{L}_{H_0} \circ \mathcal{I}_{H_0, g}^\Lambda(A) - iA, \rho \right] = 0$$

$$\Leftrightarrow \rho \left( \left[ \mathcal{L}_{H_0} \circ \mathcal{I}_{H_0, g}^\Lambda(A) - iA, B \right] \right) = 0 \quad \forall B \in \mathcal{A}_\Lambda$$

and one can choose  $A_1 = \mathcal{I}_{H_0, g}^\Lambda(V)$ .

# Elements of the proof: the local inverse of the Liouvillian

Based on techniques of **Moon, Ogata**, *JFA* '19 we show that

$$\mathcal{I}_{H_0, g} := \lim_{k \rightarrow \infty} \mathcal{I}_{H_0, g}^{\wedge k}$$

exists as a bounded operator from  $\mathcal{D}_{f_1}$  to  $\mathcal{D}_{f_2}$  and satisfies:

## Lemma

Let  $H_0$  have a gap in the bulk.

Then for all  $A \in \mathcal{A}$  with  $\mathcal{I}_{H_0, g}(A) \in D(\mathcal{L}_{H_0})$  and all  $B \in D(\mathcal{L}_{H_0})$

$$\rho([\mathcal{L}_{H_0} \circ \mathcal{I}_{H_0, g}(A) - iA, B]) = 0.$$

# Elements of the proof: the local inverse of the Liouvillian

Based on techniques of **Moon, Ogata**, *JFA* '19 we show that

$$\mathcal{I}_{H_0, g} := \lim_{k \rightarrow \infty} \mathcal{I}_{H_0, g}^{\wedge k}$$

exists as a bounded operator from  $\mathcal{D}_{f_1}$  to  $\mathcal{D}_{f_2}$  and satisfies:

## Lemma

Let  $H_0$  have a gap in the bulk.

Then for all  $A \in \mathcal{A}$  with  $\mathcal{I}_{H_0, g}(A) \in D(\mathcal{L}_{H_0})$  and all  $B \in D(\mathcal{L}_{H_0})$

$$\rho([\mathcal{L}_{H_0} \circ \mathcal{I}_{H_0, g}(A) - iA, B]) = 0.$$

**Problem:**  $V \notin \mathcal{A}$ .

# Elements of the proof: the local inverse of the Liouvillian

Based on techniques of **Moon, Ogata**, *JFA* '19 we show that

$$\mathcal{I}_{H_0, g} := \lim_{k \rightarrow \infty} \mathcal{I}_{H_0, g}^{\wedge k}$$

exists as a bounded operator from  $\mathcal{D}_{f_1}$  to  $\mathcal{D}_{f_2}$  and satisfies:

## Lemma

Let  $H_0$  have a gap in the bulk.

Then for all  $A \in \mathcal{A}$  with  $\mathcal{I}_{H_0, g}(A) \in D(\mathcal{L}_{H_0})$  and all  $B \in D(\mathcal{L}_{H_0})$

$$\rho([\mathcal{L}_{H_0} \circ \mathcal{I}_{H_0, g}(A) - iA, B]) = 0.$$

**Problem:**  $V \notin \mathcal{A}$ .

**Solution:** Take the thermodynamic limit for  $H_0$  and the perturbation  $V$  independently.

## Concluding remarks

- ▶ Proving uniqueness of the ground state  $\rho$  of  $\mathcal{L}_{H_0}$  and “fast convergence” of  $\rho^{\wedge k} \rightarrow \rho$ , e.g.

$$|(\rho - \rho^{\wedge})(B)| \leq C_n \|B\| \text{dist}(X, \partial\Lambda)^{-n}$$

for all  $B \in \mathcal{A}_X$ , are difficult problems that have not yet been achieved for interacting fermionic systems.

## Concluding remarks

- ▶ Proving uniqueness of the ground state  $\rho$  of  $\mathcal{L}_{H_0}$  and “fast convergence” of  $\rho^{\wedge k} \rightarrow \rho$ , e.g.

$$|(\rho - \rho^{\wedge})(B)| \leq C_n \|B\| \text{dist}(X, \partial\Lambda)^{-n}$$

for all  $B \in \mathcal{A}_X$ , are difficult problems that have not yet been achieved for interacting fermionic systems.

For weakly interacting spin systems such a result has been shown by **Yarotzky**, JSP '05.

## Concluding remarks

- ▶ Proving uniqueness of the ground state  $\rho$  of  $\mathcal{L}_{H_0}$  and “fast convergence” of  $\rho^{\wedge k} \rightarrow \rho$ , e.g.

$$|(\rho - \rho^{\wedge})(B)| \leq C_n \|B\| \text{dist}(X, \partial\Lambda)^{-n}$$

for all  $B \in \mathcal{A}_X$ , are difficult problems that have not yet been achieved for interacting fermionic systems.

For weakly interacting spin systems such a result has been shown by **Yarotzky**, JSP '05.

For non-interacting fermionic systems this can be shown, but I know no reference.

## Concluding remarks

- ▶ Proving uniqueness of the ground state  $\rho$  of  $\mathcal{L}_{H_0}$  and “fast convergence” of  $\rho^{\wedge k} \rightarrow \rho$ , e.g.

$$|(\rho - \rho^{\wedge})(B)| \leq C_n \|B\| \text{dist}(X, \partial\Lambda)^{-n}$$

for all  $B \in \mathcal{A}_X$ , are difficult problems that have not yet been achieved for interacting fermionic systems.

For weakly interacting spin systems such a result has been shown by **Yarotzky**, JSP '05.

For non-interacting fermionic systems this can be shown, but I know no reference.

- ▶ A similar justification of linear response for systems with mobility gap instead of spectral gap is a difficult open problem even for non-interacting systems.

## References



J. Henheik and S.T.

Justifying Kubo's formula for gapped systems at zero temperature:  
A brief review and some new results.

Reviews in Mathematical Physics, Online First (2020).

# References



J. Henheik and S.T.

Justifying Kubo's formula for gapped systems at zero temperature:  
A brief review and some new results.

Reviews in Mathematical Physics, Online First (2020).



J. Henheik and S.T.

Adiabatic theorem in the thermodynamic limit. Part I:  
Systems with a uniform gap

ArXiv 2012.15238 (2020).

# References



J. Henheik and S.T.

Justifying Kubo's formula for gapped systems at zero temperature:  
A brief review and some new results.

Reviews in Mathematical Physics, Online First (2020).



J. Henheik and S.T.

Adiabatic theorem in the thermodynamic limit. Part I:  
Systems with a uniform gap

ArXiv 2012.15238 (2020).



J. Henheik and S.T.

Adiabatic theorem in the thermodynamic limit. Part II:  
Systems with a gap in the bulk.

ArXiv 2012.15239 (2020).

## References



J. Henheik and S.T.

Justifying Kubo's formula for gapped systems at zero temperature:  
A brief review and some new results.

Reviews in Mathematical Physics, Online First (2020).



J. Henheik and S.T.

Adiabatic theorem in the thermodynamic limit. Part I:  
Systems with a uniform gap

ArXiv 2012.15238 (2020).



J. Henheik and S.T.

Adiabatic theorem in the thermodynamic limit. Part II:  
Systems with a gap in the bulk.

ArXiv 2012.15239 (2020).



S.T.

Non-equilibrium almost-stationary states and linear response for  
gapped quantum systems.

Communications in Mathematical Physics 373:621–653 (2020).

## More references



S. Bachmann, W. de Roeck, and M. Fraas.

The adiabatic theorem and linear response theory for extended quantum systems.

Communications in Mathematical Physics, 361:997–1027 (2018).

## More references



S. Bachmann, W. de Roeck, and M. Fraas.

The adiabatic theorem and linear response theory for extended quantum systems.

Communications in Mathematical Physics, 361:997–1027 (2018).



A. Moon and Y. Ogata.

Automorphic equivalence within gapped phases in the bulk.

Journal of Functional Analysis 108422 (2019).

## More references



S. Bachmann, W. de Roeck, and M. Fraas.

The adiabatic theorem and linear response theory for extended quantum systems.

Communications in Mathematical Physics, 361:997–1027 (2018).



A. Moon and Y. Ogata.

Automorphic equivalence within gapped phases in the bulk.

Journal of Functional Analysis 108422 (2019).



S. Bachmann, S. Michalakis, B. Nachtergaele, and R. Sims.

Automorphic equivalence within gapped phases of quantum lattice systems.

Communications in Mathematical Physics 309:835–871 (2012).

## More references



S. Bachmann, W. de Roeck, and M. Fraas.  
The adiabatic theorem and linear response theory for extended quantum systems.  
Communications in Mathematical Physics, 361:997–1027 (2018).



A. Moon and Y. Ogata.  
Automorphic equivalence within gapped phases in the bulk.  
Journal of Functional Analysis 108422 (2019).



S. Bachmann, S. Michalakis, B. Nachtergaele, and R. Sims.  
Automorphic equivalence within gapped phases of quantum lattice systems.  
Communications in Mathematical Physics 309:835–871 (2012).



D. Yarotzky  
Uniqueness of the ground state in weak perturbations of non-interacting gapped quantum lattice systems..  
Journal of Statistical Physics 118:119–144 (2005).

## More references



S. Bachmann, W. de Roeck, and M. Fraas.  
The adiabatic theorem and linear response theory for extended quantum systems.  
Communications in Mathematical Physics, 361:997–1027 (2018).



A. Moon and Y. Ogata.  
Automorphic equivalence within gapped phases in the bulk.  
Journal of Functional Analysis 108422 (2019).



S. Bachmann, S. Michalakis, B. Nachtergaele, and R. Sims.  
Automorphic equivalence within gapped phases of quantum lattice systems.  
Communications in Mathematical Physics 309:835–871 (2012).



D. Yarotzky  
Uniqueness of the ground state in weak perturbations of non-interacting gapped quantum lattice systems..  
Journal of Statistical Physics 118:119–144 (2005).



G. Panati, H. Spohn, and S.T.  
Effective dynamics for Bloch electrons: Peierls substitution and beyond  
Communications in Mathematical Physics 242:547–578 (2003).

## More references



S. Bachmann, W. de Roeck, and M. Fraas.  
The adiabatic theorem and linear response theory for extended quantum systems.  
Communications in Mathematical Physics, 361:997–1027 (2018).



A. Moon and Y. Ogata.  
Automorphic equivalence within gapped phases in the bulk.  
Journal of Functional Analysis 108422 (2019).

**Thanks for your attention!**



D. Yarotzky  
Uniqueness of the ground state in weak perturbations of non-interacting gapped quantum lattice systems..  
Journal of Statistical Physics 118:119–144 (2005).



G. Panati, H. Spohn, and S.T.  
Effective dynamics for Bloch electrons: Peierls substitution and beyond  
Communications in Mathematical Physics 242:547–578 (2003).